

BLOAT GUARD- poloxalene powder
Phibro Animal Health

Bloat Guard®
(poloxalene)

TYPE A MEDICATED ARTICLE

For use in the manufacture of feed

For prevention of legume (alfalfa, clover) and wheat pasture bloat in cattle

Active Ingredient

Poloxalene 53%

(Equivalent to 240 g/lb)

Inert Ingredients:

Ethoxyquin (a preservative), 0.064%; butylated hydroxytoluene (a preservative), 0.20%; Verxite ® (nonnutritive), approximately 47%

CAUTION:

For use in manufacturing medicated animal feeds only.

CAUTION: Certain components of animal feeds, including medicated premixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety.

Precautions such as the following should be considered: dust masks or respirators and protective clothing should be worn; dust arresting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions—seek prompt medical treatment if such reactions are suspected.

STORE AT ROOM TEMPERATURE

Mixing Directions

Bloat Guard Type A Medicated Article must be thoroughly and evenly distributed into feedstuffs. A mixture of Bloat Guard and 1 common feed ingredient should be made before final mixing when less than 25 lb of Bloat Guard is to be blended into a ton of feed.

Possible Mixing Ratios

Each lb of Bloat Guard contains 240 g of poloxalene, the active drug ingredient. The following table illustrates various amounts of Bloat Guard which may be added to feeds. The resulting concentration of poloxalene is also shown.

lb of Bloat Guard per ton of Type C Medicated Feed	g of poloxalene per lb of Type C Medicated Feed	% w/w of poloxalene in Type C Medicated Feed
8.3 lb	1	0.22
16.6 lb	2	0.44

33.2 lb	4	0.88
50.0 lb	6	1.32
83.3 lb	10	2.20
100.0 lb	12	2.64
208.3 lb	25	5.50
416.6 lb	50	11.00
625.0 lb	75	16.50
833.3 lb	100	22.00

For example: 50 lb of Bloat Guard, when added to 1 ton of feed, results in a final concentration of 6 g poloxalene per lb of Type C Medicated Feed (1.32% by weight).

Feeding Directions

The dosage of Bloat Guard is proportional to body weight and also depends upon the severity of the bloat-producing conditions. The normal dosage of Bloat Guard (1 g of poloxalene per 100 lb of body weight) is recommended for cattle under moderate bloat-producing conditions. For cattle under severe bloat-producing conditions, the normal dosage should be doubled (2 g of poloxalene per 100 lb of body weight).

Repeat the feeding of Bloat Guard when animals are exposed to bloat-producing conditions more than 12 hours from the last feeding of Bloat Guard, but do not exceed the double dosage of Bloat Guard in any 24-hour period. Animals should be fed the recommended amounts of Bloat Guard starting 2 or 3 days before they are exposed to bloat-producing conditions.

Caution: It is essential that each animal consumes the total recommended dosage of Bloat Guard daily for adequate protection.

Warning:

The normal life of this product is at least 24 months. However, when the product is subjected to extreme temperatures (38°C/100°F) for long periods of time (6 months), spontaneous combustion may occur. The product is not combustible unless it develops a strong, irritating odor; if this occurs, flush with water and discard immediately.

NOT FOR HUMAN USE

RESTRICTED DRUG (CALIFORNIA) – USE ONLY AS DIRECTED

Follow Directions on Back Panel

Bloat Guard® is a Phibro Animal Health registered trademark for Poloxalene

Phibro Animal Health, Inc., Ridgefield Park, NJ 07660.

Net Weight 50 lb (22.7 kg)

Approved by FDA under NADA #032-704

8818000

101-9029-09

Verxite® is a registered trademark of W.R. Grace & Company

Bloat Guard®
(poloxalene)

Phibro
ANIMAL HEALTH CORPORATION



Bloat Guard®

(poloxalene)

TYPE A
MEDICATED ARTICLE

For use in the manufacture of feed
For prevention of legume (alfalfa, clover) and
wheat pasture bloat in cattle

Active Ingredient
Poloxalene..... 53%
(Equivalent to 240 g/lb)

Inert Ingredients:
Ethoxyquin (a preservative), 0.064%; butylated hydroxytoluene (a preservative),
0.20%; Verzie® (nonnutritive), approximately 47%

CAUTION: For use in manufacturing medicated animal feeds only.

CAUTION: Certain components of animal feeds, including medicated premixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety.

Precautions such as the following should be considered: dust masks or respirators and protective clothing should be worn; dust arresting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions—seek prompt medical treatment if such reactions are suspected.

Follow Directions on Back Panel

Bloat Guard® is a Phibro Animal Health Corporation registered trademark for Poloxalene
Phibro Animal Health Corporation, Teaneck, NJ 07666

Approved by FDA under NADA # 032-704
8818000
101-9029-09

Net Weight 50 lb (22.7 kg)

Phibro
ANIMAL HEALTH CORPORATION

Phibro
ANIMAL HEALTH CORPORATION

Bloat Guard®
(poloxalene)

8818000
101-9029-09



Bloat Guard®

(poloxalene)

TYPE A
MEDICATED ARTICLE



Mixing Directions

Bloat Guard Type A Medicated Article must be thoroughly and evenly distributed into feed-stuffs. A mixture of Bloat Guard and 1 common feed ingredient should be made before final mixing when less than 25 lb of Bloat Guard is to be blended into a ton of feed.

Possible Mixing Ratios

Each lb of Bloat Guard contains 240 g of poloxalene, the active drug ingredient. The following table illustrates various amounts of Bloat Guard which may be added to feeds. The resulting concentration of poloxalene is also shown.

lb of Bloat Guard per ton of Type C Medicated Feed	g of poloxalene per lb of Type C Medicated Feed	% w/w of poloxalene in Type C Medicated Feed
8.3 lb	1	0.22
16.6 lb	2	0.44
33.2 lb	4	0.88
50.0 lb	6	1.32
83.3 lb	10	2.20
100.0 lb	12	2.64
200.0 lb	25	5.50
416.6 lb	50	11.00
625.0 lb	75	16.50
833.3 lb	100	22.00

For example: 50 lb of Bloat Guard, when added to 1 ton of feed, results in a final concentration of 6 g poloxalene per lb of Type C Medicated Feed (1.32% by weight).

Feeding Directions

The dosage of Bloat Guard is proportional to body weight and also depends upon the severity of the bloat-producing conditions. The normal dosage of Bloat Guard (1 g of poloxalene per 100 lb of body weight) is recommended for cattle under moderate bloat-producing conditions. For cattle under severe bloat-producing conditions, the normal dosage should be doubled (2 g of poloxalene per 100 lb of body weight).

Repeat the feeding of Bloat Guard when animals are exposed to bloat-producing conditions more than 12 hours from the last feeding of Bloat Guard, but do not exceed the double dosage of Bloat Guard in any 24-hour period. Animals should be fed the recommended amounts of Bloat Guard starting 2 or 3 days before they are exposed to bloat-producing conditions.

Caution: It is essential that each animal consumes the total recommended dosage of Bloat Guard daily for adequate protection.

Warning: The normal life of this product is at least 24 months. However, when the product is subjected to extreme temperatures (38°C/100°F) for long periods of time (6 months), spontaneous combustion may occur. The product is not combustible unless it develops a strong, irritating odor; if this occurs, flush with water and discard immediately.

Ventis® is a registered trademark of W.R. Grace & Company.
Store At Or Below 25°C (77°F), Excursions Permitted Up To 40°C (104°F)

Not For Human Use

Restricted Drug (California) - Use Only as Directed

8818000
101-9029-09

Distributed by:
Phibro Animal Health Corporation
Teaneck, NJ 07666

BLOAT GUARD

poloxalene powder

Product Information

Product Type

OTC TYPE A MEDICATED ARTICLE ANIMAL
DRUG

Item Code
(Source)

NDC:66104-
2301

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLOXALENE (UNII: V8B3K56SW0) (POLOXALENE - UNII:V8B3K56SW0)	POLOXALENE	240 g in 0.45 kg

Inactive Ingredients

Ingredient Name

Strength

BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)

ETHOXYQUIN (UNII: 9T1410R4OR)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66104-2301-0	22.7 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA032704	03/10/1966	

Labeler - Phibro Animal Health (006989008)

Registrant - Phibro Animal Health (006989008)

Revised: 6/2019

Phibro Animal Health