SINUS RELIEF PRESSURE, PAIN AND COUGH- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated VALU MERCHANDISERS COMPANY

1199-BST-2021-1118

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves:
 - headache
 - nasal congestion
 - minor aches and pains
 - sinus congestion and pressure
 - cough
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- diabetes
- liver disease
- heart disease
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

store between 20-25°C (68-77°F) in a dry place

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

COMPARE TO THE ACTIVE INGREDIENTS IN MUCIINEX® SINUS-MAX® PRESSURE, PAIN & COUGH†

Best Choice®

MAXIMUM STRENGTH

SINUS RELIEF

Pressure, Pain & Cough

ACTUAL SIZE

Relieves Sinus Pressure & Congestion

Relieves Headache, Controls Cough

Thins & Loosens Mucus

Pain Reliever - Acetaminophen

Cough Suppressant - Dextromethorphan HBr

Expectorant - Guaifenesin

Nasal Decongestant - Phenylephrine HCl

For Ages 12+

20 CAPLETS

PROUDLY DISTRIBUTED BY: VALU MERCHANDISERS, CO. 5000 KANSAS AVE

www.bestchoicebrand.com

100% Guaranteed Best Choice,

Maximum Strength Per 4-Hour Dose

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

† This product is not manufactured or distributed by Reckitt Benckleer, distributor of Mucinex® Sinus-Max® Pressure, Pain & Cough.

1-844-705-4384 Guestions or comments?

Drug Facts (continued)

dioxide, com starch, croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate,

glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

maltodextrin, microcrystalline cellulose, polyeth

Pressure, Pain & Coug SINUS RELIEF

Drug Facts

persistent or chronic cough such as occurs with

When using this product do not use more than direct Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

 nervousness, dizziness, or sleeplessness occu ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days Stop use and ask a doctor if

Active ingredients (in each caplet) Purpose . Nasal decongestan

 nasal congestion
 minor aches and pains Uses ... temporarily relieves: Guaifenesin 200 mg....... Phenylephrine HCl 5 mg...

sinus congestion and pressure

■ cough

■ promotes nasal and/or sinus drainage m helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs

fever gets worse or lasts more than 3 days
 redness or swelling is present

headache that lasts These could be signs of a serious condition. If pregnant or breast-feeding, ask a health

cough comes back or occurs with rash or

new symptoms occur

Liver warning: This product contains acetaminopher Severe liver damage may occur if you take more than 12 caplets in 24 hours, which is the

more productive

with other drugs containing acetaminophen
 s or more alcoholic drinks daily while using this product

recommended dose (overdose) may cause liver damage. In case of overdose, get medical help contact a Poison Control Center right away

Keep out of reach of children. Overdose warning: Taking more than the

critical for adults as well as for children even if

do not notice any signs or sympto

(1-800-222-1222). Quick medical attention

Directions a do not take more than

 do not take more than 12 caplets in any directed (see Overdose warning)

24-hour period

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

skin reddening
 blisters

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

with any other drug containing acetaminophen

(prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

adults and children 12 years of age and over-take 2 caplets every 4 hours

children under 12 years of age: do not use

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

■ store between 20-25°C (68-77°F) in a dry place ■ retain outer package for complete product

Ask a doctor before use if you have a diabetes or pharmacist before taking this product.

 Inver disease
 Insight blood pressure
 Introuble urinating due to an enlarged prostate gland cough that occurs with too much phiegm (mucus)



MAXIMUM STRENGTH

SINUS RELIEF Pressure, Pain & Cough

COMPARE TO THE ACTIVE INGREDIENTS IN MUCINEX® SINUS-MAX® PRESSURE, PAIN & COUGHT

MAXIMUM STRENGTH

SINUS RELIEF Pressure, Pain & Cough

Relieves Sinus Pressure & Congestion, Relieves Headache, Controls Cough Thins & Loosens Mucus

Pain Reliever - Acetaminophen Cough Suppressant - Dextromethorphan HBr Expectorant - Guaifenesin Nasal Decongestant – Phenylephrine HCI

For Ages 12+

20 CAPLETS



SINUS RELIEF PRESSURE, PAIN AND COUGH

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-399
--------------	----------------	--------------------	---------------

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients			
Ingredient Name	Strength		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)			
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)			
STARCH, CORN (UNII: O8232NY3SJ)			
STEARIC ACID (UNII: 4ELV7Z65AP)			
TALC (UNII: 7SEV7J4R1U)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)			
CROSPOVIDONE (UNII: 2S7830E561)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
ALUMINUM OXIDE (UNII: LMI26O6933)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MALTODEXTRIN (UNII: 7CVR7L4A2D)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			

Product Characteristics				
Color	red	Score	no score	
Shape	OVAL	Size	19mm	
Flavor		Imprint Code	AAA;1165	
Contains				

Packaging			
# Hom Codo	Packago Doscription	Marketing Start	Marketing End

#	item Code	гаскаде резсприон	Date	Date
1	NDC:63941- 399-02	2 in 1 CARTON	03/01/2021	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/01/2021	

Labeler - VALU MERCHANDISERS COMPANY (868703513)

Revised: 10/2024 VALU MERCHANDISERS COMPANY