

# **MUCUS DM- dextromethorphan hydrobromide guaifenesin liquid**

## **Rite Aid Corporation**

-----

### **Drug Facts**

#### **Active ingredients (in each 20 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

#### **Purpose**

Cough suppressant

Expectorant

#### **Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily relieves
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep

#### **Warnings**

##### **Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

##### **Ask a doctor before use if you have**

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

##### **When using this product,**

**do not use more than directed.**

##### **Stop use and ask a doctor if**

cough lasts more than 7 days, comes back, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other device
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

**Other information**

- **each 20 mL contains: sodium 20 mg**
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients**

citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

**Principal Display Panel**

\*Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® DM Max

MAXIMUM STRENGTH

mucus relief

DM max

dextromethorphan HBr 20 mg

guaifenesin 400 mg

cough suppressant

expectorant

controls cough

relieves chest congestion

thin & loosens mucus

4 hour dosing

for ages 12+

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® DM Max

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**DISTRIBUTED BY: RITE AID**

**30 HUNTER LANE**

**CAMP HILL, PA 17011**

**Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST**

**Package Label**

**RITE AID**  
PHARMACY

\*Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® DM Max

**MAXIMUM STRENGTH**  
**mucus relief**  
**DM max**

dextromethorphan HBr 20 mg  
guaifenesin 400 mg

cough suppressant  
expectorant

controls cough  
relieves chest congestion  
thins & loosens mucus  
4 hour dosing  
for ages 12+

**6 FL OZ (177 mL)**

DISTRIBUTED BY: RITE AID  
30 HUNTER LANE  
CAMP HILL, PA 17011

IF YOU'RE NOT SATISFIED,  
WE'LL HAPPILY REFUND  
YOUR MONEY.

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

ITEM 367224    PLD-4408A    LB003881



0 11822 37224 4

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® DM Max.

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

PLD-D409A  
LB004108

**Drug Facts**

Active ingredients (in each 20 mL)	Purposes
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep

**Warnings**

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Drug Facts (continued)**

**Ask a doctor before use if you have**

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**When using this product, do not use more than directed.**

**Stop use and ask a doctor** if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Drug Facts (continued)**

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

**Other information**

- each 20 mL contains: sodium 20 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Drug Facts (continued)**

**Inactive ingredients**

citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

PEEL CORNER FOR MORE DRUG FACTS

**RITE AID Maximum Strength Mucus Relief DM Max**

**MUCUS DM**

dextromethorphan hydrobromide guaifenesin liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-5050	
<b>Route of Administration</b>	ORAL			
<b>Active Ingredient/Active Moiety</b>				
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)		GUAIFENESIN	400 mg in 20 mL	
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>			<b>Strength</b>	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYL GALLATE (UNII: 8D4SNN7V92)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
XANTHAN GUM (UNII: TTV12P4NEE)				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:11822-5050-6	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2019	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
OTC Monograph Drug	M012	09/30/2019		

**Labeler** - Rite Aid Corporation (014578892)