

**ROCKLATAN- netarsudil and latanoprost ophthalmic solution,
0.02%/0.005% solution/ drops
Alcon Laboratories, Inc.**

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ROCKLATAN® safely and effectively. See full prescribing information for ROCKLATAN®.

**ROCKLATAN® (netarsudil and latanoprost ophthalmic solution) 0.02%/0.005%, for topical ophthalmic use
Initial U.S. Approval: 2019**

-----**RECENT MAJOR CHANGES**-----

Warnings and Precautions, Epithelial Corneal Edema (5.1)

01/2025

-----**INDICATIONS AND USAGE**-----

ROCKLATAN® is a fixed dose combination of a Rho kinase inhibitor and a prostaglandin F_{2α} analogue indicated for the reduction of elevated intraocular pressure (IOP) in patients with open-angle glaucoma or ocular hypertension. (1)

-----**DOSAGE AND ADMINISTRATION**-----

One drop in the affected eye(s) once daily in the evening. (2)

-----**DOSAGE FORMS AND STRENGTHS**-----

Ophthalmic solution containing netarsudil 0.02% and latanoprost 0.005%. (3)

-----**CONTRAINDICATIONS**-----

None. (4)

-----**WARNINGS AND PRECAUTIONS**-----

- Pigmentation: Pigmentation of the iris, periorbital tissue (eyelid) and eyelashes can occur. Iris pigmentation likely to be permanent. (5.2)
- Eyelash Changes: Gradual change to eyelashes including increased length, thickness and number of lashes. Usually reversible. (5.3)

-----**ADVERSE REACTIONS**-----

The most common adverse reaction is conjunctival hyperemia (59%). Other common adverse reactions were: instillation site pain (20%), corneal verticillata (15%), and conjunctival hemorrhage (11%). (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Alcon Laboratories, Inc. at 1 800 757 9195, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----**DRUG INTERACTIONS**-----

Thimerosal: *In vitro* studies have shown that precipitation can occur when eye drops containing thimerosal are mixed with ROCKLATAN®. If such drugs are used, they should be administered at least 5 minutes apart. (7)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 1/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1. INDICATIONS AND USAGE**
- 2. DOSAGE AND ADMINISTRATION**
- 3. DOSAGE FORMS AND STRENGTHS**

4. CONTRAINDICATIONS

5. WARNINGS AND PRECAUTIONS

- 5.1 Epithelial Corneal Edema
- 5.2 Pigmentation
- 5.3 Eyelash Changes
- 5.4 Intraocular Inflammation
- 5.5 Macular Edema
- 5.6 Herpetic Keratitis
- 5.7 Bacterial Keratitis
- 5.8 Use with Contact Lenses

6. ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7. DRUG INTERACTIONS

8. USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

11. DESCRIPTION

12. CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13. NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14. CLINICAL STUDIES

16. HOW SUPPLIED/STORAGE AND HANDLING

17. PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1. INDICATIONS AND USAGE

ROCKLATAN is a fixed dose combination of a Rho kinase inhibitor and a prostaglandin F_{2α} analogue indicated for the reduction of elevated intraocular pressure (IOP) in patients with open-angle glaucoma or ocular hypertension.

2. DOSAGE AND ADMINISTRATION

The recommended dosage is one drop in the affected eye(s) once daily in the evening. If one dose is missed, treatment should continue with the next dose in the evening. The dosage of ROCKLATAN should not exceed once daily.

ROCKLATAN may be used concomitantly with other topical ophthalmic drug products to lower IOP. If more than one topical ophthalmic drug is being used, the drugs should be

administered at least five (5) minutes apart.

3. DOSAGE FORMS AND STRENGTHS

Ophthalmic solution containing netarsudil 0.02% (0.2 mg/mL) and latanoprost 0.005% (0.05 mg/mL).

4. CONTRAINDICATIONS

None.

5. WARNINGS AND PRECAUTIONS

5.1 Epithelial Corneal Edema

ROCKLATAN contains netarsudil which has been associated with Epithelial corneal edema, described as honeycomb or bullous, and has been reported in some patients with pre-existing corneal stromal edema or following ocular procedures that could affect corneal endothelial function. Epithelial corneal edema typically resolves upon discontinuation of ROCKLATAN. Advise patients to notify their physician if they experience eye pain or decreased vision while using ROCKLATAN [*see Adverse Reactions (6.2) and Patient Counselling Information (17)*].

5.2 Pigmentation

ROCKLATAN contains latanoprost which has been reported to cause changes to pigmented tissues. The most frequently reported changes have been increased pigmentation of the iris, periorbital tissue (eyelid), and eyelashes. Pigmentation is expected to increase as long as latanoprost is administered.

The pigmentation change is due to increased melanin content in the melanocytes rather than to an increase in the number of melanocytes. After discontinuation of latanoprost, pigmentation of the iris is likely to be permanent, while pigmentation of the periorbital tissue and eyelash changes have been reported to be reversible in some patients. Patients who receive treatment should be informed of the possibility of increased pigmentation. Beyond 5 years the effects of increased pigmentation are not known.

Iris color change may not be noticeable for several months to years. Typically, the brown pigmentation around the pupil spreads concentrically towards the periphery of the iris and the entire iris or parts of the iris become more brownish. Neither nevi nor freckles of the iris appear to be affected by treatment. While treatment with ROCKLATAN can be continued in patients who develop noticeably increased iris pigmentation, these patients should be examined regularly [*see Patient Counseling Information (17)*].

5.3 Eyelash Changes

ROCKLATAN contains latanoprost which may gradually change eyelashes and vellus hair in the treated eye; these changes include increased length, thickness, pigmentation, the number of lashes or hairs, and misdirected growth of eyelashes. Eyelash changes are usually reversible upon discontinuation of treatment [*see Patient Counseling Information (17)*].

5.4 Intraocular Inflammation

ROCKLATAN contains latanoprost which should be used with caution in patients with a history of intraocular inflammation (iritis/uveitis) and should generally not be used in patients with active intraocular inflammation because it may exacerbate inflammation.

5.5 Macular Edema

Macular edema, including cystoid macular edema, has been reported during treatment with latanoprost. ROCKLATAN should be used with caution in aphakic patients, in pseudophakic patients with a torn posterior lens capsule, or in patients with known risk factors for macular edema.

5.6 Herpetic Keratitis

Reactivation of Herpes Simplex keratitis has been reported during treatment with latanoprost. ROCKLATAN should be used with caution in patients with a history of herpetic keratitis. ROCKLATAN should be avoided in cases of active herpes simplex keratitis because it may exacerbate inflammation.

5.7 Bacterial Keratitis

There have been reports of bacterial keratitis associated with the use of multiple-dose containers of topical ophthalmic products. These containers had been inadvertently contaminated by patients who, in most cases, had a concurrent corneal disease or a disruption of the ocular epithelial surface [*see Patient Counseling Information (17)*].

5.8 Use with Contact Lenses

Contact lenses should be removed prior to the administration of ROCKLATAN and may be reinserted 15 minutes after administration.

6. ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The most common ocular adverse reaction observed in controlled clinical studies with ROCKLATAN was conjunctival hyperemia which was reported in 59% of patients. Five percent of patients discontinued therapy due to conjunctival hyperemia. Other common ocular adverse reactions reported were: instillation site pain (20%), corneal verticillata (15%), and conjunctival hemorrhage (11%). Eye pruritus, visual acuity reduced, increased lacrimation, instillation site discomfort, and blurred vision were reported in 5-8% of patients.

Other adverse reactions from clinical trials that have been reported with the individual components and not listed above include:

Netarsudil 0.02%

Instillation site erythema, corneal staining, increased lacrimation, and erythema of eyelid.

Latanoprost 0.005%

Foreign body sensation, punctate keratitis, burning and stinging, itching, increased pigmentation of the iris, excessive tearing, eyelid discomfort, dry eye, eye pain, eyelid margin crusting, erythema of the eyelid, upper respiratory tract infection/nasopharyngitis/influenza, photophobia, eyelid edema, myalgia/arthralgia/back pain, and rash/allergic reactions.

6.2 Postmarketing Experience

The following adverse reactions have been identified during postmarketing use of ROCKLATAN. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Netarsudil 0.02%

Eye disorders: Epithelial corneal edema has been reported in some patients with pre-existing corneal stromal edema or ocular procedures that could affect corneal endothelial function. [see *Warnings and Precautions (5.1)*].

7. DRUG INTERACTIONS

In vitro drug interaction studies have shown that precipitation can occur when eye drops containing thimerosal are mixed with ROCKLATAN. If such drugs are used, they should be administered at least five (5) minutes apart.

The combined use of two or more prostaglandins or prostaglandin analogs including latanoprost ophthalmic solution 0.005% is not recommended. It has been shown that administration of these prostaglandin drug products more than once daily may decrease the IOP lowering effect or cause paradoxical elevations in IOP.

8. USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no adequate and well-controlled studies of ROCKLATAN ophthalmic solution or its pharmacologically active ingredients (netarsudil and latanoprost) in pregnant women to inform any drug associated risk. However, systemic exposure to netarsudil from ocular administration is low [see *Clinical Pharmacology (12.3)*].

Reproduction studies of latanoprost showed embryofetal lethality in rabbits. No embryofetal lethality was observed at a dose approximately 15 times higher than the recommended human ophthalmic dose (RHOD). Intravenous administration of netarsudil to pregnant rats and rabbits during organogenesis did not produce adverse embryofetal effects at clinically relevant systemic exposures. ROCKLATAN should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Data

Animal Data

Netarsudil administered daily by intravenous injection to rats during organogenesis caused abortions and embryofetal lethality at doses ≥ 0.3 mg/kg/day (126-fold the plasma exposure at the RHOD, based on C_{max}). The no-observed-adverse-effect-level (NOAEL) for embryofetal development toxicity was 0.1 mg/kg/day (40-fold the plasma exposure at the RHOD, based on C_{max}).

Netarsudil administered daily by intravenous injection to rabbits during organogenesis caused embryofetal lethality and decreased fetal weight at 5 mg/kg/day (1480-fold the plasma exposure at the RHOD, based on C_{max}). Malformations were observed at ≥ 3 mg/kg/day (1330-fold the plasma exposure at the RHOD, based on C_{max}), including thoracogastroschisis, umbilical hernia and absent intermediate lung lobe. The NOAEL for embryofetal development toxicity was 0.5 mg/kg/day (214-fold the plasma exposure at the RHOD, based on C_{max}).

Reproduction studies have been performed with latanoprost in rats and rabbits. In 4 of 16 pregnant rabbits, no viable fetuses were present at a dose that was approximately 80 times higher than the RHOD. Latanoprost did not produce embryofetal lethality in rabbits at a dose approximately 15 times higher than the RHOD.

8.2 Lactation

Risk Summary

There are no data on the presence of netarsudil or latanoprost in human milk, the effects on the breastfed infant, or the effects on milk production. However, systemic exposure to netarsudil following topical ocular administration is low, and it is not known whether measurable levels of netarsudil would be present in maternal milk following topical ocular administration.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for ROCKLATAN and any potential adverse effects on the breast-fed child from ROCKLATAN.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.


8.5 Geriatric Use

No overall differences in safety or effectiveness have been observed between elderly and other adult patients.

11. DESCRIPTION


ROCKLATAN (netarsudil and latanoprost ophthalmic solution) 0.02%/0.005% is a fixed dose combination of a Rho kinase inhibitor and a prostaglandin $F_{2\alpha}$ analogue.

The chemical name of netarsudil dimesylate is: (S)-4-(3-amino-1-(isoquinolin-6-yl-amino)-1-oxopropan-2-yl)benzyl 2,4-dimethylbenzoate dimesylate. Its molecular formula is $C_{30}H_{35}N_3O_9S_2$ and its chemical structure is:



Netarsudil mesylate is a light yellow to white powder that is freely soluble in water, soluble in methanol, sparingly soluble in dimethyl formamide, and practically insoluble in dichloromethane and heptane.

The chemical name of latanoprost is: isopropyl-(Z)-7[1R,2R,3R,5S] 3,5-dihydroxy-2-[(3R)-3-hydroxy-5-phenylpentyl]cyclopentyl]-5-heptenoate. Its molecular formula is $C_{26}H_{40}O_5$ and its chemical structure is:



Latanoprost is a colorless to slightly yellow oil that is very soluble in acetonitrile and freely soluble in acetone, ethanol, ethyl acetate, isopropanol, methanol, and octanol. It is practically insoluble in water.

ROCKLATAN (netarsudil and latanoprost ophthalmic solution) 0.02%/0.005% is supplied as a sterile, isotonic, buffered aqueous solution of netarsudil mesylate and latanoprost with a pH of approximately 5 and an osmolality of approximately 295 mOsmol/kg. Each mL of ROCKLATAN contains 0.20 mg of netarsudil (equivalent to 0.28 mg of netarsudil dimesylate) and 0.05 mg latanoprost. Benzalkonium chloride, 0.02%, is added as a preservative. The inactive ingredients are: boric acid, mannitol, sodium hydroxide to adjust pH, and water for injection.

12. CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

ROCKLATAN is comprised of two components: netarsudil and latanoprost. Each of these two components decreases elevated IOP. Elevated IOP represents a major risk factor

for glaucomatous field loss. The higher the level of IOP, the greater the likelihood of optic nerve damage and glaucomatous visual field loss.

ROCKLATAN is believed to reduce IOP by increasing the outflow of aqueous humor.

12.3 Pharmacokinetics

Absorption

The systemic exposures of netarsudil and its active metabolite, AR-13503, were evaluated in 18 healthy subjects after topical ocular administration of netarsudil ophthalmic solution 0.02% once daily (1 drop bilaterally in the morning) for 8 days. There were no quantifiable plasma concentrations of netarsudil (lower limit of quantitation (LLOQ) 0.100 ng/mL) post dose on Day 1 and Day 8. Only 1 plasma concentration at 0.11 ng/mL for the active metabolite was observed for 1 subject on Day 8 at 8 hours post-dose.

Distribution

The distribution volume in humans is 0.16 ± 0.02 L/kg. Latanoprost is absorbed through the cornea where the isopropyl ester prodrug is hydrolyzed to the acid form to become biologically active. The acid of latanoprost can be measured in aqueous humor during the first 4 hours, and in plasma only during the first hour after local administration.

Metabolism

After topical ocular dosing, netarsudil is metabolized by esterases in the eye to an active metabolite, AR-13503.

Latanoprost, an isopropyl ester prodrug, is hydrolyzed by esterases in the cornea to the biologically active acid. The active acid of latanoprost reaching the systemic circulation is primarily metabolized by the liver to the 1,2-dinor and 1,2,3,4-tetranor metabolites via fatty acid β -oxidation.

Excretion

The elimination of the acid of latanoprost from human plasma is rapid ($t_{1/2} = 17$ min) after both intravenous and topical administration. Systemic clearance is approximately 7 mL/min/kg. Following hepatic β -oxidation, the metabolites are mainly eliminated via the kidneys. Approximately 88% and 98% of the administered dose are recovered in the urine after topical and intravenous dosing, respectively.

13. NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term studies in animals have not been performed to evaluate the carcinogenic potential of netarsudil.

Latanoprost was not carcinogenic in either mice or rats when administered by oral gavage at doses of up to 170 mcg/kg/day (approximately 2800 times the RHOD) for up to 20 and 24 months, respectively.

Mutagenesis

Netarsudil was not mutagenic in the Ames test, in the mouse lymphoma test, or in the *in vivo* rat micronucleus test.

Latanoprost was not mutagenic in bacteria, in mouse lymphoma, or in mouse micronucleus tests. Chromosome aberrations were observed *in vitro* with human lymphocytes. Additional *in vitro* and *in vivo* studies on unscheduled DNA synthesis in rats were negative.

Impairment of Fertility

Studies to evaluate the effects of netarsudil on male or female fertility in animals have not been performed.


Latanoprost has not been found to have effects on male or female fertility in animal studies.

14. CLINICAL STUDIES

ROCKLATAN (netarsudil and latanoprost ophthalmic solution) 0.02%/0.005% was evaluated in 2 randomized and controlled clinical trials, namely PG324-CS301 (NCT 02558400, referred to as Study 301) and PG324-CS302 (NCT 02674854, referred to as Study 302) in patients with open-angle glaucoma and ocular hypertension. Studies 301 and 302 enrolled subjects with IOP < 36 mmHg and compared IOP lowering effect of ROCKLATAN dosed once daily to individually administered netarsudil 0.02% once daily and latanoprost 0.005% once daily. The treatment duration was 12 months for Study 301 and 3 months for Study 302.


The average IOP lowering effect of ROCKLATAN was 1 to 3 mmHg greater than monotherapy with either netarsudil 0.02% or latanoprost 0.005% throughout 3 months (Figures 1 and 2). In Study 301 IOP reductions were maintained throughout 12 months.

Figure 1: Study 301 Mean IOP (mmHg) by Treatment Group and Treatment Difference in Mean IOP



The least square mean IOP at each post-baseline time point was derived using an analysis of covariance adjusted for baseline IOP and based on observed data for all randomized subjects (238 in Rocklatan group, 244 in netarsudil group, 236 in latanoprost group).

Figure 2: Study 302 Mean IOP (mmHg) by Treatment Group and Treatment Difference in Mean IOP



The least square mean IOP at each post-baseline time point was derived using an analysis of covariance adjusted for baseline IOP and based on observed data for all randomized subjects (245 in Rocklatan group, 255 in netarsudil group, 250 in latanoprost group).

16. HOW SUPPLIED/STORAGE AND HANDLING

ROCKLATAN (netarsudil and latanoprost ophthalmic solution) 0.02%/0.005% is supplied sterile in clear low density polyethylene bottles with opaque white polyethylene dropper tips and white polypropylene screw caps.

2.5 mL fill in a 4 mL container - NDC # 70727-529-25

Storage: Protect from light. Until opened, store at 2°C to 8°C (36°F to 46°F). After opening, the product may be kept at 2°C to 25°C (36°F to 77°F) for up to 6 weeks. If after opening the product is kept refrigerated at 2°C to 8°C (36°F to 46°F), then the product can be used until the expiration date stamped on the bottle. During shipment, the bottle may be maintained at temperatures up to 40°C (104°F) for a period not exceeding 14 days.

17. PATIENT COUNSELING INFORMATION

Potential for Pigmentation

Advise patients about the potential for increased brown pigmentation of the iris, which may be permanent. Inform patients about the possibility of eyelid skin darkening, which may be reversible after discontinuation of ROCKLATAN [see *Warnings and Precautions (5.2)*].

Potential for Eyelash Changes

Inform patients of the possibility of eyelash and vellus hair changes in the treated eye during treatment with ROCKLATAN. These changes may result in a disparity between eyes in length, thickness, pigmentation, number of eyelashes or vellus hairs, and/or direction of eyelash growth. Eyelash changes are usually reversible upon discontinuation of treatment.

Handling the Container

Instruct patients to avoid allowing the tip of the dispensing container to contact the eye, surrounding structures, fingers, or any other surface in order to minimize contamination of the solution. Serious damage to the eye and subsequent loss of vision may result from using contaminated solutions [see *Warnings and Precautions (5.7)*].

When to Seek Physician Advice

Advise patients that if they develop an intercurrent ocular condition (e.g., trauma, or infection, or decreased vision with or without eye pain), have ocular surgery, or develop any ocular reactions, particularly conjunctivitis and eyelid reactions, they should immediately seek their physician's advice concerning the continued use of ROCKLATAN.

Use with Contact Lenses

Advise patients that ROCKLATAN contains benzalkonium chloride, which may be absorbed by soft contact lenses. Contact lenses should be removed prior to instillation of ROCKLATAN and may be reinserted 15 minutes following its administration.

Use with Other Ophthalmic Drugs

If more than 1 topical ophthalmic drug is being used, the drugs should be administered at least 5 minutes between applications.

Missed Dose

Advise patients that if one dose is missed, treatment should continue with the next dose in the evening.

© 2024-2025 Alcon Inc.

U.S. Pat.: www.alconpatents.com

ALCON LABORATORIES, Inc., Fort Worth, Texas 76134

1-800-757-9195

Alcon.medinfo@alcon.com

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

NDC 70727-529-25

rocklatan[®]

(netarsudil and latanoprost
ophthalmic solution) 0.02%/0.005%

For topical application
in the eye

Sterile
Rx only

2.5 mL

Usual dosage:

One drop in the affected
eye(s) once daily in the
evening.

Each ml Rocklatan[®]
ophthalmic solution contains:

Active: netrasudil mesylate
0.285 mg

latanoprost 0.050 mg

Preservative:

benzalkonium chloride
0.20 mg

Inactives: mannitol, boric
acid, sodium hydroxide to
adjust pH and water
for injection.

Manufactured for:
ALCON LABORATORIES, INC.
Fort Worth, TX 76134

U.S. Pat.: www.alconpatents.com

PROTECT FROM LIGHT

Until opened, store at
2°C - 8°C (36°F - 46°F).

After opening, the product
may be kept at 2°C - 25°C
(36°F - 77°F) for up to
6 weeks.

If after opening the product is
kept refrigerated at 2°C - 8°C
(36°F - 46°F), then the
product can be used until the

expiration date stamped on the bottle. During shipment, the bottle may be maintained at temperatures up to 40°C (104°F) for a period not exceeding 14 days.



NDC 70727-529-25

rocklatan®

(netarsudil and latanoprost
ophthalmic solution) 0.02%/0.005%

Sterile 2.5 mL Rx only

For topical application in the eye.

See carton
for storage conditions.
Manufactured for:
Alcon Laboratories, Inc.
Fort Worth, TX 76134

LOT:
EXP:


NDC 70727-529-25

rocklatan®
(netarsudil and latanoprost
ophthalmic solution) 0.02%/0.005%

Sterile 2.5 mL Rx only **Alcon** 41083

For topical application in the eye.
See carton for
storage conditions.
Manufactured for:
Alcon Laboratories, Inc.
Fort Worth, TX 76134

(01)00370727529253

NDC 70727-529-99

SAMPLE - Not for Resale

rocklatan®

(netarsudil and latanoprost
ophthalmic solution) 0.02%/0.005%

For topical application
in the eye

Sterile
Rx only
2.5 mL

Usual dosage:

One drop in the affected
eye(s) once daily in the
evening.

Each ml Rocklatan®
ophthalmic solution contains:

Active: netrasudil mesylate
0.285 mg
latanoprost 0.050 mg

Preservative:

benzalkonium chloride

0.20 mg

Inactives: mannitol, boric acid, sodium hydroxide to adjust pH and water for injection.

Manufactured for:

ALCON LABORATORIES, INC.

Fort Worth, TX 76134

U.S. Pat.: www.alconpatents.com

PROTECT FROM LIGHT

Until opened, store at 2°C - 8°C (36°F - 46°F).

After opening, the product may be kept at 2°C - 25°C (36°F - 77°F) for up to 6 weeks.

If after opening the product is kept refrigerated at 2°C - 8°C (36°F - 46°F), then the product can be used until the expiration date stamped on the bottle. During shipment, the bottle may be maintained at temperatures up to 40°C (104°F) for a period not exceeding 14 days.



NDC 70727-529-99 SAMPLE - Not for Resale


rocklatan®
 (netarsudil and latanoprost
 ophthalmic solution) 0.02%/0.005%

Sterile 2.5 mL Rx only

For topical application in the eye.

See carton for
storage conditions.
Manufactured for:
Alcon Laboratories, Inc.
Fort Worth, TX 76134

LOT:
EXP:



NDC 70727-529-99 **SAMPLE – Not for Resale** For topical application in the eye.
rocklatan[®]
(netarsudil and latanoprost
ophthalmic solution) 0.02%/0.005%
Sterile 2.5 mL Rx only **Alcon**

See carton for
storage conditions.
Manufactured for:
Alcon Laboratories, Inc.
Fort Worth, TX 76134
41084

NDC 70727-529-25


rocklatan[®]
(netarsudil and latanoprost
ophthalmic solution) 0.02%/0.005%

**For topical application
in the eye**

Sterile
Rx only

2.5 mL

3 2 4 4



OCR-B 6pt Font (1.4 mm minimum character height); all UpperCase.

2D code and human readable data will be printed on-line (during packaging). Size shown here is only a visual representation of the production carton. Not to size. For reference only proofing printed information.

NDC 70727-529-99


rocklatan®
(netarsudil and latanoprost
ophthalmic solution) 0.02%/0.005%

**For topical application
in the eye**

Sterile
Rx only

2.5 mL
SAMPLE - Not for Resale

3 2 4 3



3 2 4 3

NDC 70727-529-25

rocklatan®
(netarsudil and latanoprost
ophthalmic solution) 0.02%/0.005%

For topical application

in the eye

Sterile
Rx only

2.5 mL

Alcon

Usual dosage:

One drop in the affected eye(s) once daily in the evening.

Each ml Rocklatan[®] ophthalmic solution contains:

Active: netrasudil mesylate 0.285 mg
latanoprost 0.050 mg

Preservative:

benzalkonium chloride 0.20 mg

Inactives: mannitol, boric acid, sodium hydroxide to adjust pH and water for injection.

ALCON LABORATORIES, INC.
Fort Worth, TX 76134 USA

U.S. Pat.: www.alconpatents.com

PROTECT FROM LIGHT

Until opened, store at 2°C - 8°C (36°F - 46°F).

After opening, the product may be kept at 2°C - 25°C (36°F - 77°F) for up to 6 weeks.

If after opening the product is kept refrigerated at 2°C - 8°C (36°F - 46°F), then the product can be used until the expiration date stamped on the bottle. During shipment, the bottle may be maintained at temperatures up to 40°C (104°F) for a period not exceeding 14 days.


300068395-1224

SN:

LOT:

EXP.

GTIN: 00370727529253



PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

NDC 70727-529-25 For topical application in the eye.

rocklatan®
 (netarsudil and latanoprost
 ophthalmic solution) 0.02%/0.005%

Sterile 2.5 mL Rx only Alcon

See carton for storage conditions.

Alcon Laboratories, Inc.
 Fort Worth, TX 76134 USA

300068406
 1224



PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

NDC 70727-529-99

SAMPLE - Not for Resale

rocklatan[®]

(netarsudil and latanoprost
ophthalmic solution) 0.02%/0.005%

For topical application
in the eye

Sterile

Rx only **2.5 mL**

Usual dosage:

One drop in the affected
eye(s) once daily in the
evening.

Each ml Rocklatan[®]
ophthalmic solution contains:

Active: netrasudil mesylate
0.285 mg

latanoprost 0.050 mg

Preservative:

benzalkonium chloride
0.20 mg

Inactives: mannitol, boric
acid, sodium hydroxide to
adjust pH and water for
injection.

ALCON LABORATORIES, INC.
Fort Worth, TX 76134 USA


U.S. Pat.: www.alconpatents.com

PROTECT FROM LIGHT

**Until opened, store at 2°C - 8°C (36°F - 46°F).
After opening, the product may be kept at 2°C - 25°C (36°F - 77°F) for up to 6 weeks.**

If after opening the product is kept refrigerated at 2°C - 8°C (36°F - 46°F), then the product can be used until the expiration date stamped on the bottle. During shipment, the bottle may be maintained at temperatures up to 40°C (104°F) for a period not exceeding 14 days.

300068396-1124



PACKAGE/LABEL PRINCIPAL DISPLAY

NDC 70727-529-99 SAMPLE - Not for Resale

rocklatan®
 (netarsudil and latanoprost
 ophthalmic solution) 0.02%/0.005%

Sterile 2.5 mL Rx only Alcon

For topical application in the eye.

See carton for
 storage conditions.

Alcon Laboratories, Inc.

Fort Worth, TX 76134 USA

LOT: EXP.:

300068407
1124



ROCKLATAN

netarsudil and latanoprost ophthalmic solution, 0.02%/0.005% solution/ drops

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70727-529
Route of Administration	OPHTHALMIC, TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NETARSUDIL MESYLATE (UNII: VL756B1K0U) (NETARSUDIL - UNII:W6I5QDT7QI)	NETARSUDIL	0.20 mg in 1 mL
Latanoprost (UNII: 6Z5B6HVF6O) (Latanoprost Acid - UNII:EJ85341990)	Latanoprost	0.05 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
MANNITOL (UNII: 3OWL53L36A)	
BORIC ACID (UNII: R57ZHV85D4)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70727-	1 in 1 CARTON	03/13/2010	

1	529-25	1 in 1 CARTON	03/12/2019	
1		2.5 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:70727-529-99	1 in 1 CARTON	03/12/2019	
2		2.5 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA208259	03/12/2019	

Labeler - Alcon Laboratories, Inc. (008018525)

Registrant - Alcon Laboratories, Inc. (008018525)

Establishment

Name	Address	ID/FEI	Business Operations
Alcon Research LLC		007672236	manufacture(70727-529)

Revised: 1/2026

Alcon Laboratories, Inc.