

ED BRON GP- guaifenesin and phenylephrine liquid
EDWARDS PHARMACEUTICALS, INC.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).

ED BRON GP

Drug Facts

Active Ingredients (in each 5 mL teaspoonful)	Purpose
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal Decongestant

Uses

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive

Warnings

Do not exceed recommended dosage.

Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Do not take this product, unless directed by a doctor, if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland

Do not take this product for persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema, or where cough is accompanied by excessive phlegm (mucus) unless directed by a doctor.

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition.
- new symptoms occur

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

Directions

Do not exceed recommended dosage.

Adults and children 12 years of age and over:	2 teaspoonfuls (10 mL) every 4 hours, not to exceed 12 teaspoonfuls in 24 hours, or as directed by a doctor.
Children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 4 hours, not to exceed 6 teaspoonfuls in 24 hours, or as directed by a doctor.
Children under 6 years of age:	Consult a doctor.

Other information

Store at 59° - 86° F (15° - 30° C) [see USP for Controlled Room Temperature].

Inactive ingredients

Citric Acid, Methyl Paraben, Orange Flavor, Potassium Citrate, Potassium Sorbate, Propyl Paraben, Propylene Glycol, Purified Water, Sorbitol Solution 70%, and Sucralose.

Question? Comments?

Call 1-800-543-9560

PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label

NDC 0485-0208-16

ED Bron GP Liquid

Expectorant • Nasal Decongestant

Sugar Free • Alcohol Free • Dye Free

Each teaspoonful (5 mL)

for oral administration contains:

Guaifenesin 100 mg

Phenylephrine HCl 5 mg

Orange Flavor

Tamper evident by foil seal under cap.

Do not use if foil seal is broken or missing.

Manufactured for:

EDWARDS Pharmaceuticals, Inc.

Paoli, PA 19301

16oz. (473 mL)

 NDC 0485-0208-16 ED Bron GP Liquid Expectorant • Nasal Decongestant Sugar Free • Alcohol Free • Dye Free Each teaspoonful (5 mL) for oral administration contains: Guaifenesin 100 mg Phenylephrine HCl 5 mg Orange Flavor Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing. Manufactured for: EDWARDS Pharmaceuticals, Inc. Paoli, PA 19301 16oz. (473 mL)	Drug Facts Active Ingredients (in each 5 mL teaspoonful) Guaifenesin 100 mg Expectorant Phenylephrine HCl 5 mg Nasal Decongestant Purpose Uses temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other respiratory allergies: ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes ■ nasal congestion ■ reduces swelling of nasal passages ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive Warnings Do not exceed recommended dosage. Do not use this product ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Do not take this product, unless directed by a doctor, if you have ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ difficulty in urination due to enlargement of the prostate gland Do not take this product for persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema, or where cough is accompanied by excessive phlegm (mucus) unless directed by a doctor. Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or	Drug Facts (continued) persistent headache. These could be signs of a serious condition. ■ new symptoms occur If pregnant or breastfeeding , ask a health professional before use. Keep out of reach of children. In case of accidental overdose, seek professional help or contact a Poison Control Center immediately. Directions Do not exceed recommended dosage. Adults and children 12 years of age and over: 2 teaspoonfuls (10 mL) every 4 hours, not to exceed 12 teaspoonfuls in 24 hours, or as directed by a doctor. Children 6 to under 12 years of age: 1 teaspoonful (5 mL) every 4 hours, not to exceed 6 teaspoonfuls in 24 hours, or as directed by a doctor. Children under 6 years of age: Consult a doctor. Other information Store at 59° - 86° F (15° - 30° C) [see USP for Controlled Room Temperature]. Inactive ingredients Citric Acid, Methyl Paraben, Orange Flavor, Potassium Citrate, Potassium Sorbate, Propyl Paraben, Propylene Glycol, Purified Water, Sorbitol Solution 70%, and Sucralose. Question? Comments? Call 1-800-543-9560 Rev. 01/13
	 N 3 04850 20816 7	

ED BRON GP

guaifenesin and phenylephrine liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0485-0208
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	ORANGE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0485-0208-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/01/2012	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/01/2012	

Revised: 10/2025

EDWARDS PHARMACEUTICALS, INC.