TUSSIN MULTI SYMPTOM COLD CF- dextromethorphan hbr guaifenesin phenylephrine hcl liquid QUALITY CHOICE (Chain Drug Marketing Association)

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr 20 mg
Guaifenesin 200 mg
Phenylephrine HCl 10 mg

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - o cough due to minor throat and bronchial irritation

Warnings

Do not use

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- thyroid
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

taking any other oral nasal decongestant or stimulant.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adult and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

Other information

• store between 20-25°C (68°-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Questions or comments?

Call 1-800-935-2362 Monday-Friday 9AM-5PM EST

Principal display panel

*Compare to the Active Ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

Adult

Tussin CF

Multi-Symptom Cold

Dextromethorphan HBr

Cough suppressant

Guaifenesin

Expectorant

Phenylephrine HCl

Nasal decongestant

Relieves:

- Cough
- Mucus
- Nasal congestion

For ages 12 years & over

Alcohol Free

Non-Drowsy

FL OZ (mL)

Dosing Cup Included

*This product is not manufactured or distributed by Haleon group of companies, distributors of Robitussin® Peak Cold Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed by CDMA, Inc.

Novi, MI 48375

Package Label

Drug Facts (continued)

Directions

- do not take more than 6 doses in any 24-hour period measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
 mL = milliliter
 this adult product is not intended for use in children under
- 12 years of age adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years; do not use

Other information

■ store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Questions or comments? Call 1-800-935-2362 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by Haleon group of companies, distributor of Robitussin® Peak Cold Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRIN SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MIS





Dosing Cup Included



NDC 83324-155-04



*Compare to the Active Ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

Adult Tussin CF

Multi-Symptom Cold

Dextromethorphan HBr Cough Suppressant

Guaifenesin

Expectorant

Phenylephrine HCI

Nasal Decongestant

Relieves:

Cough

Mucus

Nasal Congestion

For Ages 12 & Over Alcohol Free Non-Drowsy

4 FL OZ (118 mL)



NDC 83324-155-04

*Compare to the Active Ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

Adult Tussin **CF**

Multi-Symptom Cold

Dextromethorphan HBr

Cough Suppressant

Guaifenesin

Expectorant

Phenylephrine HCI

Nasal Decongestant

Relieves:

Cough

Mucus

Nasal Congestion

For Ages 12 & Over Alcohol Free Non-Drowsy

4 FL OZ (118 mL)



PLD-F322C FC009142

Lot No.:

QUALITY CHOICE Adult Multi-Symptom Cold

TUSSIN MULTI SYMPTOM COLD CF

dextromethorphan hbr guaifenesin phenylephrine hcl liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:83324-155

Route of Administration ORAL

Active Ingredient/Active Moiety Ingredient Name **Basis of Strength** Strength **DEXTROMETHORPHAN HYDROBROMIDE** (UNII: 9D2RTI9KYH) **DEXTROMETHORPHAN** 20 mg **HYDROBROMIDE** in 10 mL (DEXTROMETHORPHAN - UNII:7355X3ROTS) 200 mg GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) **GUAIFENES IN** in 10 mL

Drug Facts

Active ingredients Purposes

(in each 10 mL)

extromethorphan HBr 20 mg. .Cough suppressar Guaifenesin 200 mg Phenylephrine HCl 10 mg.. Expectorant ...Nasal decongestan

USES

The helps loosen phiegm (mucus) and thin bronchial secretions to drain bronchial tubes

Temporarily relieves these symptoms occurring with a cold

The masal congestion

Cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this

Ask a doctor before use if you have

- Ask a occurretore see it you have

 lear disease length line of pressure

 diabetes thyroid disease

 trouble urinating due to an enlarged prostate gland

 cough that cocurs with too much philegm (mucus)

 cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any othe oral nasal decongestant or stimulant.

When using this product, do not use more than directed.

Stop use and ask a doctor if

nervousness, dizziness, or sleeplessness occur

- symptoms do not get better within 7 days or are accompanied by
- cough lasts more than 7 days, comes back, or is accompanied fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before

use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222)

PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE -	PHENYLEPHRINE	10 mg
UNII:1WS297W6MV)	HYDROCHLORIDE	in 10 mL

Inactive Ingredients				
Ingredient Name	Strength			
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
LACTIC ACID (UNII: 33X04XA5AT)				
MENTHOL (UNII: L7T10EIP3A)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				

Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:83324- 155-04	1 in 1 BOX	07/31/2024		
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	07/31/2024		

Labeler - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)

Revised: 5/2024 QUALITY CHOICE (Chain Drug Marketing Association)