

**TUSSIN MULTI SYMPTOM COLD CF- dextromethorphan hbr guaifenesin phenylephrine hcl liquid**  
**QUALITY CHOICE (Chain Drug Marketing Association)**

-----

**Drug Facts**

**Active ingredients (in each 10 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 200 mg

Phenylephrine HCl 10 mg

**Purposes**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold
  - nasal congestion
  - cough due to minor throat and bronchial irritation

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are**

taking any other oral nasal decongestant or stimulant.

**When using this product,  
do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adult and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

**Other information**

- store between 20-25°C (68°-77°F). Do not refrigerate.

**Inactive ingredients**

citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

**Questions or comments?**

Call **1-800-935-2362** Monday-Friday 9AM-5PM EST

**Principal display panel**

\*Compare to the Active Ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

**Adult**

**Tussin CF**

## **Multi-Symptom Cold**

Dextromethorphan HBr

Cough Suppressant

Guaifenesin

Expectorant

Phenylephrine HCl

Nasal Decongestant

Relieves:

- Cough
- Mucus
- Nasal congestion

For ages 12 years & over

Alcohol Free

FL OZ (mL)

Dosing Cup Included

\*This product is not manufactured or distributed by Haleon group of companies, distributor of Robitussin® Peak Cold Multi-Symptom Cold CF.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Distributed by CDMA, Inc.

Novi, MI 48375

[www.qualitychoice.com](http://www.qualitychoice.com)

Questions: 800-935-2362

**Package Label**

**Drug Facts (continued)**

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

**Other information**

- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

**Questions or comments?**  
Call 1-800-935-2362 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by Haleon group of companies, distributor of Robitussin® Peak Cold Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

**PARENTS:**  
Learn about how medicine works  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



Dosing Cup Included



Distributed by CDMA, Inc.  
Novi, MI 48375  
[www.qualitychoice.com](http://www.qualitychoice.com)  
Questions: 800-935-2362

NDC 83324-155-04



\*Compare to the Active Ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

## Adult Tussin CF

### Multi-Symptom Cold

Dextromethorphan HBr  
Cough Suppressant  
Guaifenesin  
Expectorant  
Phenylephrine HCl  
Nasal Decongestant

Relieves:  
Cough  
Mucus  
Nasal Congestion

For Ages 12 & Over  
Alcohol Free

4 FL OZ (118 mL)

NDC 83324-155-04



\*Compare to the Active Ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

## Adult Tussin CF

### Multi-Symptom Cold

Dextromethorphan HBr  
Cough Suppressant  
Guaifenesin  
Expectorant  
Phenylephrine HCl  
Nasal Decongestant

Relieves:  
Cough  
Mucus  
Nasal Congestion

For Ages 12 & Over  
Alcohol Free

4 FL OZ (118 mL)

**Drug Facts**

**Active ingredients (in each 10 mL)**

Dextromethorphan HBr 20 mg.....Cough suppressant  
Guaifenesin 200 mg.....Expectorant  
Phenylephrine HCl 10 mg.....Nasal decongestant

**Purposes**

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold
- nasal congestion
- cough due to minor throat and bronchial irritation

**Warnings**

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- high blood pressure
- thyroid disease

**Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.**

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.



6 35515 95706 0

PLD-G322D FC009657

Lot No.:

Exp. Date:

## Quality Choice Adult Tussin CF Multi-Symptom Cold

## TUSSIN MULTI SYMPTOM COLD CF

dextromethorphan hbr guaifenesin phenylephrine hcl liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83324-155
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL

<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL	
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>			<b>Strength</b>	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
LACTIC ACID (UNII: 33X04XA5AT)				
MENTHOL (UNII: L7T10EIP3A)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:83324-155-04	1 in 1 BOX	07/31/2024	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
OTC Monograph Drug	M012	07/31/2024		

**Labeler** - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)

Revised: 1/2026

QUALITY CHOICE (Chain Drug Marketing Association)