

STOMACH RELIEF REGULAR STRENGTH- bismuth subsalicylate liquid
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)

Drug Facts

Active ingredient (in each 30 mL dose)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including
 - heartburn
 - indigestion
 - nausea
 - gas
 - fullness
 - belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL (1 dose) every 1/2 hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
 - 30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
 - do not exceed 8 doses (240 mL) in 24 hours
 - use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL contains:** sodium 11 mg
- **each 30 mL contains:** salicylate 206 mg
- low sodium
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?

Call **1-888-309-9030**

Principal Display Panel

Compare to the active ingredient of Pepto-Bismol®*

Regular Strength Stomach Relief

Bismuth Subsalicylate, 525 mg

Upset Stomach Reliever/Antidiarrheal

Relieves:

- Heartburn
- Indigestion
- Nausea
- Upset stomach
- Diarrhea

Alcohol Free

Sugar Free

Cherry Flavor

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY:

OLD EAST MAIN CO.

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

Package Label

DG health Compare to the active ingredient of Pepto-Bismol**

Regular Strength Stomach Relief

Bismuth Subsalicylate, 525 mg
Upset Stomach Reliever/
Antidiarrheal

Relieves:

- Heartburn
- Indigestion
- Nausea
- Upset stomach
- Diarrhea

Alcohol Free
Sugar Free

Cherry Flavor

8 FL OZ (237 mL)

PLD-A493A LB009623



Drug Facts

Active ingredient (in each 30 mL dose)
Bismuth subsalicylate 525 mg.....Upset stomach reliever/Antidiarrheal

Uses relieves ■ diarrhea ■ travelers' diarrhea ■ upset stomach due to overindulgence in food and drink, including ■ heartburn ■ gas ■ indigestion ■ nausea ■ fullness ■ belching

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY Product of Germany
OLD EAST MAIN CO.
100 MISSION RIDGE
GOODLETTSVILLE,
TN 37072

100% Satisfaction Guaranteed! (888) 309-9030

A0209



3 49580 73608 5
PLD-A493A LB009968

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Contains salicylate. Do not take if you are:

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product
a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical

Drug Facts (continued)

help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over ■ 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- each 30 mL contains: sodium 11 mg
- each 30 mL contains: salicylate 206 mg
- low sodium
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?
Call 1-888-309-9030

Dollar General Regular Strength Stomach Relief

STOMACH RELIEF REGULAR STRENGTH

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-436
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KOOR)	
XANTHAN GUM (UNII: TTV12P4NEE)	

SUCRALOSE (UNII: 96K6UQ3ZD4)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-436-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/29/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	11/29/2019	

Labeler - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 8/2024

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)