

**STOMACH RELIEF REGULAR STRENGTH- bismuth subsalicylate liquid**  
**Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

-----  
**Drug Facts**

**Active ingredient (in each 30 mL dose)**

Bismuth subsalicylate 525 mg

**Purpose**

Upset stomach reliever/Antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - fullness
  - belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use**

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

## **Ask a doctor or pharmacist before use if you are**

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

## **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur.

## **Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## **Directions**

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over
  - 30 mL (1 dose) every 1/2 or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
  - 30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
  - do not exceed 8 doses (240 mL) in 24 hours
  - use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

## **Other information**

- **each 30 mL contains:** sodium 11 mg
- **each 30 mL contains:** salicylate 206 mg
- low sodium
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

## **Inactive ingredients**

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

## **Questions or comments?**

Call **1-888-309-9030**

## **Principal Display Panel**

Compare to the active ingredient of Pepto-Bismol®\*

### **Regular Strength Stomach Relief**

Bismuth subsalicylate 525 mg

Upset Stomach Reliever/Antidiarrheal

Relieves:

- Heartburn
- Indigestion
- Nausea
- Upset stomach
- Diarrhea

Alcohol Free

Sugar Free

FL OZ (mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company, Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

**TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY: DOLGENCORP, LLC

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

## **Package Label**

**Drug Facts (continued)**

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** ■ mL = milliliter ■ shake well before using  
 ■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product ■ adults and children 12 years and over ■ 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years of age; ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 30 mL contains: sodium 11 mg ■ each 30 mL contains: salicylate 206 mg ■ low sodium ■ keep tightly closed ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)

**Inactive ingredients** benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

**Questions or comments?** Call 1-888-309-9030

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

\*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

**100% Satisfaction Guaranteed!**  
(888) 309-9030

DISTRIBUTED BY  
DOLGENCORP, LLC  
100 MISSION RIDGE  
GOODLETTSVILLE,  
TN 37072

**Compare to the active ingredient of Pepto-Bismol®\***

## Regular Strength Stomach Relief

**Bismuth Subsalicylate, 525 mg  
Upset Stomach Reliever/  
Antidiarrheal**

Relieves:

- Heartburn
- Indigestion
- Nausea
- Upset stomach
- Diarrhea

Alcohol Free  
Sugar Free

**8 FL OZ (237 mL)**

Cherry Flavor

**Drug Facts**

**Active ingredient (in each 30 mL dose)** Purpose  
 Bismuth subsalicylate 525 mg.....Upset stomach reliever/Antidiarrheal

**Uses** relieves ■ diarrhea ■ travelers' diarrhea ■ upset stomach due to overindulgence in food and drink, including ■ heartburn ■ gas ■ indigestion ■ nausea ■ fullness ■ belching

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert: Contains salicylate. Do not take if you are:**  
 ■ allergic to salicylates (including aspirin)  
 ■ taking other salicylate products

**Do not use** if you have  
 ■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have**  
 ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for** ■ diabetes ■ gout ■ arthritis ■ anticoagulation (thinning the blood)

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if**  
 ■ symptoms get worse or last more than 2 days  
 ■ ringing in the ears or loss of hearing occurs  
 ■ diarrhea lasts more than 2 days

### DOLLAR GENERAL HEALTH Regular Strength Stomach Relief

## STOMACH RELIEF REGULAR STRENGTH

bismuth subsalicylate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-436
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 7671P0Y5NH)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-436-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/29/2019	
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008		11/29/2019	

**Labeler** - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 12/2023

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)