

**ASPIRIN 81 MG- aspirin tablet, coated**  
**Reliable 1 Laboratories LLC**

-----  
**Aspirin 81mg EC**

***Active ingredient (in each tablet)***

Aspirin 81 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

temporarily relieves minor aches and pains

**Because of its delayed action, this product will not produce fast relief of headaches or other symptoms needing immediate relief.**

ask your doctor about other uses for enteric-coated 81 mg Aspirin

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be early sign of Reye's syndrome, a rare but serious illness.

**Allergy Alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

- take more or for longer time than directed

**Do not use:** if you are allergic to aspirin or any other pain reliver/fever reducer

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis

- taking any other drug

- under a doctor's care for any serious condition

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.

- you are experiance any of the following signs of stomach bleeding: feel faint, have bloody or black stools, vomit blood, have stomach pain that does not get better

- pain gets worse or lasts more than 10 days

- redness or swelling is present

- new symptoms occur

- ringing in the ears or a loss of hearing occurs.

these could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor becwause it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### ***Directions***

- Drink a full glass of water with each dose

- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

- children under 12 years: consult a doctor

### ***Other information***

- store at 25°C (77°F) excursions premitted between 15°-30°C (59°-86°F)

- Use by expiration date on package

### ***Inactive ingredients***

corn starch, pregelatinized starch, povidone. microcrystalline cellulose, colloidal silicon dioxide, stearic acid, methacrylic acid and ethyl acrylate copolymer, talc, titanium dioxide, triethyl citrate, colloidal silicon dioxide, sodium bicarbonate, sodium lauryl sulfate, d&c

yellow #10, hypromellose, triacetin.

### Ask a doctor before use if

stomach bleeding warning applies to you

you have a history of stomach problems, such as heartburn

you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

you are taking a diuretic

you have asthma

you have not been drinking fluids

you have lost a lot of fluid due to vomiting or diarrhea

### Drug Facts

NDC 69168-066-02

Reliable-1

Adult Low Strength

Pain Reliever

Aspirin USP 81 mg (NSAID)

Enteric Coated

120 tablets

Compare to Active Ingredient in Aspirin Regimen BAYER®

NDC 69168-066-02

**Reliable-1**  
LABORATORIES

ADULT LOW STRENGTH

**Pain Reliever**  
**Aspirin USP**  
ENTERIC COATED 81 MG (NSAID)

120 Tablets

†Compare to Active Ingredient in Aspirin Regimen BAYER®

Reliable-1 Laboratories is not affiliated with the owner of the trademark BAYER®. Rev. 02  
Distributed by: Reliable-1 Laboratories LLC  
Valley Stream, NY 11580  
www.reliable1labs.com  
516-341-0666

3 69618 06602 1

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\* ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains  
**Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.** ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer

**Drug Facts** continued on back of label

PEEL HERE FOR MORE DRUG FACTS

LOT #: \_\_\_\_\_  
EXP. DATE: \_\_\_\_\_

**UNVARNISHED AREA**

### Drug Facts (continued)

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint.
  - have bloody or black stools.
  - vomit blood.
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days.
- redness or swelling is present.
- new symptoms occur.
- ringing in the ears or a loss of hearing occurs. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredients** colloidal silicon dioxide, corn starch, D&C yellow #10, hypromellose, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, povidone, pregelatinized starch, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate.

STOP PEEING



UNVARNISHED AREA

PEEL HERE FOR MORE DRUG FACTS

LOT #: 3

EXP. DATE:

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.

Drug Facts continued on back of label

Ask a doctor before use if

- stomach bleeding warning applies to you

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

NDC 69618-066-10



ADULT LOW STRENGTH

Pain Reliever  
Aspirin USP

ENTERIC COATED 81 MG (NSAID)

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

1000 Tablets

†Compare to Active Ingredient in Aspirin Regimen BAYER®

### Drug Facts (continued)

**Ask a doctor before use if**

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint.
  - have bloody or black stools.
  - vomit blood.
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days.
- redness or swelling is present.
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredients** Colloidal silicon dioxide, corn starch, D&C yellow #10, hypromellose, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, povidone, pregelatinized starch, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate.

STOP PEEING

# ASPIRIN 81 MG

aspirin tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69618-066
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
POVIDONE (UNII: FZ989GH94E)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TRACETIN (UNII: XHX3C3X673)	

## Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	5mm
<b>Flavor</b>		<b>Imprint Code</b>	17;S
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69618-066-10	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2020	
2	NDC:69618-066-02	120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2020	

**Drug Facts** (continued)

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma ■ you have not been drinking fluids ■ you have lost a lot of fluid due to vomiting or diarrhea

**Ask a doctor or pharmacist before use if you are** ■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug ■ under a doctor's care for any serious condition

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ● feel faint ● have bloody or black stools ● vomit blood ● have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present ■ new symptoms occur ■ ringing in the ears or a loss of hearing occurs. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.


**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

**Other information** ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

**Inactive ingredients** corn starch, pregelatinized starch, povidone, microcrystalline cellulose, colloidal silicon dioxide, stearic acid, methacrylic acid and ethyl acrylate copolymer, talc, titanium dioxide, triethyl citrate, colloidal anhydrous silica, sodium bicarbonate, sodium lauryl sulfate, d&c yellow # 10

**STOP PEELING**



1Reliable-1 Laboratories is not affiliated with the owner of the trademark BAYER®. Distributed by: **Reliable-1 Laboratories LLC** Rev. 00 Valley Stream, NY 11580 www.reliable1labs.com 516-341-0666



**Drug Facts**

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Active ingredient (in each tablet)	Purpose
Aspirin 81 mg (NSAID)*	Pain reliever
*nonsteroidal anti-inflammatory drug	

**Uses** ■ temporarily relieves minor aches and pains. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.** ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

**Warnings**


**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer

**Drug Facts** continued on back of label



LOT #:

EXP. DATE:

**UNVARNISHED AREA**

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/01/2020	

**Labeler** - Reliable 1 Laboratories LLC (079718111)

**Registrant** - Reliable 1 Laboratories LLC (079718111)

Revised: 1/2025

Reliable 1 Laboratories LLC