



binding to sterols in the cell membrane of susceptible *Candida* species with a resultant change in membrane permeability allowing leakage of intracellular components. Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

## **INDICATIONS AND USAGE**

Nystatin Oral Suspension is indicated for the treatment of candidiasis in the oral cavity.

## **CONTRAINDICATIONS**

The preparation is contraindicated in patients with a history of hypersensitivity to any of its components.

## **PRECAUTIONS**

### **General**

This medication is not to be used for the treatment of systemic mycoses.

Discontinue treatment if sensitization or irritation is reported during use.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

No long-term animal studies have been performed to evaluate carcinogenic potential. There also have been no studies to determine mutagenicity or whether this medication affects fertility in males or females.

### **Pregnancy**

Teratogenic Effects Category C

Animal reproduction studies have not been conducted with nystatin oral suspension. It is also not known whether nystatin oral suspension can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Nystatin oral suspension should be given to a pregnant woman only if clearly needed.

### **Nursing Mothers**

It is not known whether nystatin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when nystatin is administered to a nursing woman.

### **Pediatric Use**

See **DOSAGE AND ADMINISTRATION**.

## **ADVERSE REACTIONS**

Nystatin is well tolerated even with prolonged therapy. Oral irritation and sensitization have been reported (See **PRECAUTIONS: General**).

*Gastrointestinal:* Diarrhea (including one case of bloody diarrhea), nausea, vomiting, gastrointestinal upset/disturbances.

*Dermatologic:* Rash, including urticaria has been reported rarely.

Stevens-Johnson syndrome has been reported very rarely.

*Other:* Tachycardia, bronchospasm, facial swelling, and non-specific myalgia have also been reported.

## **OVERDOSAGE**

Oral doses of nystatin in excess of five million units daily have caused nausea and gastrointestinal upset. There have been no reports of serious toxic effects or superinfections (See **CLINICAL PHARMACOLOGY, Pharmacokinetics**).

## **DOSAGE AND ADMINISTRATION**

**INFANTS:** 2 mL (200,000 units) four times daily (in infants and young children, use dropper to place one-half of dose in each side of mouth and avoid feeding for 5 to 10 minutes).

**NOTE:** Limited clinical studies in premature and low birth weight infants indicate that 1 mL four times daily is effective.

**CHILDREN AND ADULTS:** 4 to 6 mL (400,000 to 600,000 units) four times daily (one-half of dose in each side of mouth). The preparation should be retained in the mouth as long as possible before swallowing.

Continue treatment for at least 48 hours after perioral symptoms have disappeared and cultures demonstrate eradication of *Candida albicans*.

## **HOW SUPPLIED**

Nystatin Oral Suspension, USP, 100,000 USP Nystatin Units per mL, is available as a cherry-mint flavored, light creamy yellow, ready-to-use suspension in 473 mL bottles.

It is supplied as follows:

NDC 64950-372-47     473 mL

## **Storage**

Store at 20°C to 25°C (68°F to 77°F) with excursions permitted between 15°C to 30°C (59°F to 86°F) [See USP Controlled Room Temperature]; avoid freezing.

Distributed by:  
Genus Lifesciences Inc.  
Allentown, PA 18102

Rev. 12/25

**PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label**

NDC 64950-372-47

Nystatin Oral  
Suspension, USP

100,000 units per mL

SHAKE WELL BEFORE USING

EACH mL CONTAINS: 100,000 USP  
Nystatin Units in a vehicle containing  
50% sucrose. Not more than 1% alcohol  
by volume.

USUAL DOSAGE FOR INFANTS:  
2 mL (200,000 units) four times daily  
(1 mL in each side of mouth).

USUAL DOSAGE FOR CHILDREN  
AND ADULTS: See package insert.

DISPENSE in tight, light-resistant  
container as defined in the USP.

STORE at 20°C to 25°C (68°F to  
77°F) with excursions permitted  
between 15°C to 30°C (59°F to 86°F)  
[See USP Controlled Room Temperature];  
avoid freezing.

1 Pint (473 mL)

Rx Only

Genus™  
Lifesciences Inc.

NDC 64950-372-47

# Nystatin Oral Suspension, USP

100,000 units per mL

SHAKE WELL BEFORE USING

Distributed by:  
Genus Lifesciences Inc.  
Allentown, PA 18102

Rev. 12/25



GTIN 00364950372472



NDC 64950-372-47

# Nystatin Oral Suspension, USP

100,000 units per mL

SHAKE WELL BEFORE USING

**EACH mL CONTAINS:** 100,000 USP Nystatin Units in a vehicle containing 50% sucrose. Not more than 1% alcohol by volume.

**USUAL DOSAGE FOR INFANTS:** 2 mL (200,000 units) four times daily (1 mL in each side of mouth).

**USUAL DOSAGE FOR CHILDREN AND ADULTS:** See package insert.

**DISPENSE** in tight, light-resistant container as defined in the USP.

**STORE** at 20°C to 25°C (68°F to 77°F) with excursions permitted between 15°C to 30°C (59°F to 86°F) [See USP Controlled Room Temperature]; avoid freezing.

1 Pint (473 mL) Rx Only



## NYSTATIN

nystatin suspension

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:64950-372
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
nystatin (UNII: BDF1O1C72E) (nystatin - UNII:BDF1O1C72E)	nystatin	100000 [USP'U] in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

<b>alcohol</b> (UNII: 3K9958V90M)	
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)	
<b>Glycerin</b> (UNII: PDC6A3C00X)	
<b>Methylparaben</b> (UNII: A2I8C7HI9T)	
<b>ETHYL MALTOL</b> (UNII: L6Q8K29L05)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>Saccharin Sodium</b> (UNII: SB8ZUX40TY)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>Sucrose</b> (UNII: C151H8M554)	
<b>CHERRY</b> (UNII: BUC5I9595W)	
<b>MINT</b> (UNII: FV98Z8GITP)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	

### Product Characteristics

<b>Color</b>	YELLOW	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY, MINT (CHERRY MINT)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:64950-372-47	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/24/2026	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065148	03/24/2026	

**Labeler** - Genus Lifesciences Inc. (113290444)

### Establishment

Name	Address	ID/FEI	Business Operations
TriRx Huntsville Pharmaceutical Services		117090286	MANUFACTURE(64950-372) , PACK(64950-372) , LABEL(64950-372) , ANALYSIS(64950-372)

Revised: 3/2026

Genus Lifesciences Inc.