

BUDESONIDE- budesonide spray, metered
CVS Pharmacy

Drug Facts

Active ingredient (in each spray)

Budesonide (glucocorticoid) 32 mcg

Purpose

Nasal allergy symptom reliever

Uses

Temporarily relieves these symptoms of hay fever or other upper respiratory allergies:

- nasal congestion
- runny nose
- itchy nose
- sneezing

Warnings

Do not use

- in children under 6 years of age
- if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed
- are using a steroid medicine for asthma, allergies or skin rash
- have an eye infection
- have or had glaucoma or cataracts

When using this product

- the growth rate of some children may be slower
- some symptoms may get better on the first day of treatment. It may take up to two weeks of daily use to feel the most symptom relief.
- do not share this bottle with anyone else as this may spread germs
- remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if

- you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
- you have or develop symptoms of an infection such as persistent fever

- you have any change in vision
- you have severe or frequent nosebleeds

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Direction

Read insert (inside package) on how to:

- get a new bottle ready (primed) before first use
- prime bottle again if not used for two days
- use the spray
- clean the spray nozzle

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

Adults and children 12 years of age and older	<ul style="list-style-type: none"> • once daily, spray 2 times into each nostril while sniffing gently • once your allergy symptoms improve, reduce to 1 spray in each nostril per day
---	--

CHILDREN 6 TO UNDER 12 YEARS OF AGE

- the growth rate of some children may be slower while using this product. Talk to your child's doctor if yours child needs to use the spray for longer than two months a year

children 6 to under 12 years of age	<ul style="list-style-type: none"> • an adult should supervise use • once daily, spray 1 times into each nostril while sniffing gently • if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day
children under 6 years of age	<ul style="list-style-type: none"> • do not use

- **do not use more than directed**
- if you forget a dose, do not double the next dose
- do not spray into eyes or mouth
- if allergy symptoms do not improve after two weeks, stop using and talk to a doctor
- do not use for the common cold
- shake well before each use

Other information

- **do not use if carton is seal are broken**
- keep the package and insert. They contain important information.
- store upright at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Do not freeze. Protect from light.

Inactive ingredients

carboxymethyl cellulose sodium, dextrose anhydrous, edetate disodium dihydrate, hydrochloric acid (for pH adjustment), microcrystalline cellulose, polysorbate 80, potassium sorbate, purified water

Questions or comments?

call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

ALLERGY RELIEF

Nasal Spray

bedesonide (Glucocorticoid) 32 mcg per Spray

24-Hour Relief of: Nasal congestion, Runny nose, Itchy nose, Sneezing

Original Prescription Strength

STORE UPRIGHT

METERED SPRAYS

FL OZ (mL)

IMPORTANT: Read all product information before using. Keep package and insert. They contain important information.

TAMPER EVIDENT: DO NOT USE IF CARTON SEALS ARE BROKEN.

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

Package label



CVS Nasal Spray



Drug Facts

Active ingredient (in each spray)
Budesonide (glucocorticoid)
32 mcg

Purpose
Nasal allergy symptom reliever

Uses
Temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 ■ nasal congestion
 ■ runny nose
 ■ sneezing
 ■ itchy nose

Warnings
 Do not use
 ■ in children under 6 years of age
 ■ if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you
 ■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed
 ■ are using a steroid medicine for asthma, allergies or skin rash
 ■ have an eye infection
 ■ have or had glaucoma or cataracts

When using this product
 ■ the growth rate of some children may be slower
 ■ some symptoms may get better on the first day of treatment. It may take up to two weeks of daily use to feel the most symptom relief.
 ■ do not share this bottle with anyone else as this may spread germs
 ■ remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if
 ■ you have, or come into contact with someone who has chickenpox, measles or tuberculosis
 ■ you have or develop symptoms of an infection such as persistent fever
 ■ you have any change in vision
 ■ you have severe or frequent nosebleeds

Drug Facts (continued)
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-232-1222) right away.

Directions
 Read insert (inside package) on how to:
 ■ get a new bottle ready (primed) before first use
 ■ prime bottle again if not used for two days
 ■ use the spray
 ■ clean the spray nozzle

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER
 adults and children 12 years of age and older
 ■ once daily, spray 2 times into each nostril while sniffing gently
 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 6 TO UNDER 12 YEARS OF AGE
 ■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.
 children 6 to under 12 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently
 ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 4 TO UNDER 6 YEARS OF AGE
 ■ do not use
 children under 6 years of age

Drug Facts (continued)

- do not use more than directed
- if you forget a dose, do not double the next dose
- do not spray into eyes or mouth
- if allergy symptoms do not improve after two weeks, stop using and talk to a doctor
- do not use for the common cold
- shake well before each use

Other information

- do not use if carton seals are broken
- keep the package and insert. They contain important information.
- store upright at 20° to 25°C (68° to 77°F) (see USP Controlled Room Temperature). Do not freeze. Protect from light.

Inactive ingredients
 carbomer/ethyl cellulose sodium, dextrose anhydrous, edate disodium dihydrate, hydrochloric acid (for pH adjustment), microcrystalline cellulose, polyorbital 80, potassium sorbate, purified water

Questions or comments?
 Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

IMPORTANT: Read all product information before using. Keep package and insert. They contain important information.

TAMPER EVIDENT: DO NOT USE IF CARTON SEALS ARE BROKEN.

Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02896
 © 2025 CVS Pharmacy
 CVS.com® 1-800-SHOP CVS
 V-34667
 100% money back guaranteed.
 CVS.com/retampcyl

NDC 69842-001-01

ALLERGY RELIEF
Nasal Spray
 Budesonide (Glucocorticoid)
 32 mcg per Spray
 Nasal Allergy Symptom Reliever

24-Hour Relief of: Nasal congestion, Runny nose, Itchy nose, Sneezing
 Original Prescription Strength

120 METERED SPRAYS | **STORE UPRIGHT**
 0.285 FL OZ (8.43 mL)

package contains one bottle actual size

PLS-NBA / 02843

#179283



ING2217-0

CVS Allergy Relief

BUDESONIDE			
budesonide spray, metered			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-001
Route of Administration	NASAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	BUDESONIDE (UNII: Q3OKS62Q6X) (Budesonide - UNII:Q3OKS62Q6X)	BUDESONIDE	32 ug
Inactive Ingredients			

Ingredient Name	Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
DEXTROSE, UNSPECIFIED FORM (UNII: IY9XDZ35W2)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-001-01	1 in 1 BOTTLE, SPRAY	07/01/2025	
1		120 in 1 BOTTLE, SPRAY; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078949	07/01/2025	

Labeler - CVS Pharmacy (062312574)

Revised: 1/2026

CVS Pharmacy