

CORID- amprolium powder, for solution
Huvepharma, Inc

CORID®

(amprolium)

20% Soluble Powder Coccidiostat

An aid in the treatment and prevention of bovine coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves

Net Contents: 10 oz. (283.5 grams)

Active Ingredient: amprolium.....20%

Distributed by:

Huvepharma, Inc.

Peachtree City, GA 30269

For questions, please call 1-877-994-4883

Approved by FDA under NADA # 033-165

IN DRINKING WATER

Mix CORID daily with fresh drinking water.

Stock solution for water proportioners may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add CORID 20% Soluble Powder to drinking water at the rate of 4 oz/50 gal. One 10 oz pack to 125 gallons of water. For use in automatic water proportioners that meter 1 fl oz of stock solution per gallon of drinking water, add 1 bag (10 oz) in 1 gallon of water to make stock solution. At the usual rate of water consumption, this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Treatment should be on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CORID 20% Soluble Powder to drinking water at the rate of 4 oz/100 gal. One 10 oz pack to 250 gallons of water. For use in automatic water proportioners that meter 1 fl oz of stock solution per gallon of drinking water, add 1/2 bag (5 oz) in 1 gallon of water to make stock solution. At usual rates of water consumption, this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

AS A DRENCH

CORID drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add 3 oz CORID 20% Soluble Powder to 1 qt (32 fl oz) of water and, with a dose syringe, give 1 fl oz of this

solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Treatment should be on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 oz of CORID 20% Soluble Powder to 1 qt (32 fl oz) of water and, with a dose syringe, give 1 fl oz of this solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

Lot No & Exp Date

CORID® (amprolium) 20% Soluble Powder - FOR ANIMAL USE ONLY

INDICATION: An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves.

CAUTION: For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

USE DIRECTIONS: There are **200 mg** of amprolium in every **1 gram** of CORID 20% Soluble Powder.

5-Day Treatment Protocol

Daily Dosage: 10 mg amprolium/kg (10 mg per 2.2 lb body weight)
1 oz = 3 1/2 measuring tablespoonsful

21-Day Prevention Protocol

Daily Dosage: 5 mg amprolium/kg (5 mg per 2.2 lb body weight)
1 oz = 28.35 grams

RESIDUE WARNING: Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

NOT FOR HUMAN USE. Restricted Drug (California)- Use only as directed. Keep this and all drugs out of the reach of children.

STORAGE: Store between 5° - 25°C (41° - 77°F) with brief excursions to 40°C (104°F). Avoid high humidity storage conditions.

Contact Huvepharma Inc. at 1-877-994-4883 or <http://www.huvepharma.us>. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

® **CORID is a registered trademark of Huvepharma EOOD.**



OBSERVE LABEL DIRECTIONS

IN DRINKING WATER

Mix CORID daily with fresh drinking water. Stock solution for water proportioners may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add CORID 20% Soluble Powder to drinking water at the rate of 4 oz/50 gal. One 10 oz pack to 125 gallons of water. For use in automatic water proportioners that meter 1 fl oz of stock solution per gallon of drinking water, add 1 bag (10 oz) in 1 gallon of water to make stock solution. At the usual rate of water consumption, this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Treatment should be on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CORID 20% Soluble Powder to drinking water at the rate of 4 oz/100 gal. One 10 oz pack to 250 gallons of water. For use in automatic water proportioners that meter 1 fl oz of stock solution per gallon of drinking water, add 1/2 bag (5 oz) in 1 gallon of water to make stock solution. At usual rates of water consumption, this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

AS A DRENCH

CORID drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add 3 oz CORID 20% Soluble Powder to 1 qt (32 fl oz) of water and, with a dose syringe, give 1 fl oz of this solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Treatment should be on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 oz of CORID 20% Soluble Powder to 1 qt (32 fl oz) of water and, with a dose syringe, give 1 fl oz of this solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

Lot No & Exp Date ▼

CORID[®] (amprolium)

20% Soluble Powder Coccidiostat

An aid in the treatment and prevention of bovine coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves

Net Contents: 10 oz. (283.5 grams)

Active Ingredient: amprolium.....20%

Distributed by:

Huvepharma, Inc.
Peachtree City, GA 30269

For questions, please call 1-877-994-4883

Approved by FDA under NADA # 033-165



CORID[®] (amprolium) 20% Soluble Powder – FOR ANIMAL USE ONLY

INDICATION: An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves.

CAUTION: For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

USE DIRECTIONS: There are 200 mg of amprolium in every 1 gram of CORID 20% Soluble Powder.

5-Day Treatment Protocol	21-Day Prevention Protocol
Daily Dosage: 10 mg amprolium/kg (10 mg per 2.2 lb body weight)	Daily Dosage: 5 mg amprolium/kg (5 mg per 2.2 lb body weight)
1 oz = 3 1/2 measuring tablespoons	1 oz = 28.35 grams

RESIDUE WARNING: Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

NOT FOR HUMAN USE. Restricted Drug (California) - Use only as directed. Keep this and all drugs out of the reach of children.

STORAGE: Store between 5° - 25°C (41° - 77°F) with brief excursions to 40°C (104°F). Avoid high humidity storage conditions.

Contact Huvepharma Inc. at 1-877-994-4883 or <http://www.huvepharma.us>. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VEIS or <http://www.fda.gov/reportanimalae>.

© CORID is a registered trademark of Huvepharma EOOD.



KCP-S00015
Version 02
Rev 03-2022

CORID

amprolium powder, for solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:23243-2309
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
amprolium ion (UNII: H2T307KMZR) (amprolium ion - UNII:H2T307KMZR)	amprolium ion	200 mg in 1 g

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-2309-3	283.5 g in 1 POUCH		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
NADA	NADA033165	12/12/2017	

Labeler - Huvepharma, Inc (619153559)

Registrant - Huvepharma EOOD (552691651)

Establishment

Name	Address	ID/FEI	Business Operations
Huvepharma Inc		883128204	analysis, pack, label, manufacture, medicated animal feed manufacture

Revised: 12/2022

Huvepharma, Inc