

**PREVENT HYDROGEN PEROXIDE 3 PERCENT- hydrogen peroxide liquid**  
**Eco-Med Pharmaceuticals, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Prevent+ Hydrogen Peroxide 3% USP**

**DRUG FACTS**

**ACTIVE INGREDIENT**

Hydrogen Peroxide 3%

**PURPOSE:**

First Aid Antiseptic/Oral debriding agent

**USES:**

First aid to help prevent the risk of infection in minor cuts, scrapes and burns. Aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth.

**WARNINGS:**

For external use only.

**Do not use**

in the eyes or apply over large areas of the body longer than one week. Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns.

**When using this product:**

Do not get into eyes. Do not apply over large areas of the body. Do not use longer than one week unless directed by a doctor .

**STOP USE AND ASK A DOCTOR IF:**

Condition persists or gets worse.

If sore mouth symptoms do not improve in 7 days.

Irritation, pain or redness persists or worsens swelling, rash or fever develops.

**Keep out of reach of children.**

In case of ingestion, get medical help or contact a Poison Control Center right away.

**DIRECTIONS:**

For first aid antiseptic use: Clean affected area, apply small amount of product on affected area 1-3 times a day. May be covered with a sterile bandage, If bandaged, let dry first, For oral debriding agent (oral rinse): Adults and children 2 years of age and over: Mix with an equal amount of water, swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. Use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor. Children under 12 years

should be supervised in the use of this product. Children under 2 years of age: consult a dentist or doctor

**OTHER INFORMATION:**

Keep tightly closed and in a dark place at room temperature. Do not shake bottle. Hold away from face when opening.

**INACTIVE INGREDIENT:**

Purified water.

**Questions:**

1-844-432-6633 Mon- Fri 8am - 4 pm (EST) Email: [info@eco-med.com](mailto:info@eco-med.com)

**Package Labeling:**

HYDROGEN PEROXIDE



# prevent

Hydrogen Peroxide  
Antiseptic Topical Solution  
3% USP

12x 500ml/16.9 fl.oz.

FOR EXTERNAL USE ONLY  
KEEP OUT OF REACH OF CHILDREN

Manufactured by



151 Carlingview Drive, Unit 14  
Etobicoke, ON, Canada M9W 5S4

NDC ????



**DRUG FACTS**

**ACTIVE INGREDIENT:** Hydrogen Peroxide 3%  
**PURPOSE:** First Aid Antiseptic/Oral debriding agent

**USES:**  
First aid to help prevent the risk of infection in minor cuts, scrapes and burns. Aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth.

**WARNINGS:**  
For external use only. Do not use in the eyes or apply over large areas of the body longer than one week. Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns. When using this product: Do not get into eyes. Do not apply over large areas of the body. Do not use longer than one week unless directed by a doctor.

**STOP USE AND ASK A DOCTOR IF:**  
• Condition persists or gets worse.  
• If sore mouth symptoms do not improve in 7 days.  
• Irritation, pain or redness persists or worsens swelling, rash or fever develops.  
Keep out of reach of children. In case of ingestion, get medical help or contact a Poison Control Center right away.

**DIRECTIONS:**  
For first aid antiseptic use: Clean affected area, apply small amount of product on affected area 1-3 times a day. May be covered with a sterile bandage. If bandaged, let dry first. For oral debriding agent (oral rinse): Adults and children 2 years of age and over: Mix with an equal amount of water, swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. Use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor. Children under 12 years should be supervised in the use of this product. Children under 2 years of age: consult a dentist or doctor

**OTHER INFORMATION:**  
Keep tightly closed and in a dark place at room temperature. Do not shake bottle. Hold away from face when opening.

**INACTIVE INGREDIENT:** Purified water.

**Questions :** 1-844-432-6633 Mon-Fri 8am - 4pm (EST) Email: info@eco-med.com

Reorder: 70HPB500

EM54-BC12

FRONT

BACK

# HYDROGEN PEROXIDE<sup>+</sup> prevent

## Hydrogen Peroxide Antiseptic Topical Solution

3% USP

500ml/16.9 fl.oz

FOR EXTERNAL USE ONLY  
KEEP OUT OF REACH OF CHILDREN

Manufactured by

**eco+med**  
PHARMACEUTICAL

151 Carlingview Drive, Unit 14  
Etobicoke, ON. Canada M9W 5S4

NDC ?????

**DRUG FACTS**

**ACTIVE INGREDIENT:** Hydrogen Peroxide 3%.....  
**PURPOSE:** First Aid Antiseptic/Oral debriding agent

**USES:**  
First aid to help prevent the risk of infection in minor cuts, scrapes and burns. Aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth.

**WARNINGS:**  
For external use only. Do not use in the eyes or apply over large areas of the body longer than one week. Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns. When using this product: Do not get into eyes. Do not apply over large areas of the body. Do not use longer than one week unless directed by a doctor.

**STOP USE AND ASK A DOCTOR IF:**

- Condition persists or gets worse.
- If sore mouth symptoms do not improve in 7 days.
- Irritation, pain or redness persists or worsens swelling, rash or fever develops.

Keep out of reach of children. In case of ingestion, get medical help or contact a Poison Control Center right away.

**DIRECTIONS:**  
For first aid antiseptic use: Clean affected area, apply small amount of product on affected area 1-3 times a day. May be covered with a sterile bandage. If bandaged, let dry first. For oral debriding agent (oral rinse): Adults and children 2 years of age and over: Mix with an equal amount of water, swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. Use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor. Children under 12 years should be supervised in the use of this product. Children under 2 years of age: consult a dentist or doctor

**OTHER INFORMATION:**  
Keep tightly closed and in a dark place at room temperature. Do not shake bottle. Hold away from face when opening.

**INACTIVE INGREDIENT:** Purified water.**Questions :** 1-844-432-6633 Mon- Fri 8am - 4pm (EST) Email: info@eco-med.com

Reorder: 70HPB500  
EM54-B

**PREVENT HYDROGEN PEROXIDE 3 PERCENT**

hydrogen peroxide liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:78835-008
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	0.03 mL in 1 mL

**Inactive Ingredients**

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:78835-008-01	12 in 1 BOX	10/26/2020	
1		500 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	10/26/2020	

**Labeler** - Eco-Med Pharmaceuticals, Inc. (249183435)

Revised: 10/2020

Eco-Med Pharmaceuticals, Inc.