

**JD ANTISEPTIC ALCOHOL 70% - alcohol liquid**  
**IWANNA CORP SAS**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**ANTISEPTIC ALCOHOL 70%**

Rubbing Alcohol

First Aid Antiseptic

Agent Antibacterial

Tamper evident: do not use if printed safety seal under cap is broken or missing

WARNING FLAMMABLE

Keep away from heat, spark, electrical, fire or flame

**Active Ingredient(s)**

Ethyl Alcohol 70% v/v. Purpose: First Aid Antiseptic

**Purpose**

First Aid Antiseptic

**Use**

First Aid to help prevent the risk of infection in minor cuts, scrapes and burns.

**Warnings**

FOR EXTERNAL USE ONLY

If taken internally, serious gastric disturbances will result

FLAMMABLE, KEEP AWAY FROM FIRE OR FLAME

USE ONLY IN A WELL VENTILATED AREA; FUMES MAY BE TOXIC

ASK A DOCTOR BEFORE USE IF YOU HAVE Deep or puncture wounds, animal bites or serious burns

**Do not use**

- 

**When using this product**

Do not get into the eyes

Do not apply over large areas of the body

Do not use longer than one week unless directed by a Doctor

.

## Keep out of reach of children

In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.(1800-222-1222)

## Stop Use and ask Doctor

if condition persists or gets worse

## Directions

- Clean affected area
- Apply a small amount of this product on the area 1-3 times daily
- If bandaged, let dry first
- May be covered with a sterile bandage

## Other information

- Store at controlled room temperature
- Will produce serious gastric disturbances if taken internally

## Inactive ingredients

purified water, polihexanide hydrochloride

## J&D

### Ethyl Alcohol 70% - Principal Label

**Drug Facts**

Active Ingredients	Purpose
Ethyl Alcohol (70% v/v)	First aid antiseptic

**Uses**  
first aid to help prevent the risk of infection in minor cuts, scrapes, and burns.

**WARNINGS**  
**FOR EXTERNAL USE ONLY**  
If taken internally, serious gastric disturbances will result.  
**FLAMMABLE, KEEP AWAY FROM FIRE OR FLAME.**  
**USE ONLY IN A WELL-VENTILATED AREA; FUMES MAY BE TOXIC.**

**ASK A DOCTOR BEFORE USE IF YOU HAVE** Deep or puncture wounds, Animal bites, or serious burns.

**WHEN USING THIS PRODUCT** ■Do not get into eyes ■Do not apply over large areas of the body ■Do not use longer than one week unless directed by a doctor

**STOP USE AND ASK A DOCTOR IF** condition persists or gets worse.

**KEEP OUT OF REACH OF CHILDREN.** In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately (1-800-222-1222)

**DIRECTIONS** ■Clean affected area ■Apply small amount of this product on the area 1-3 times daily ■If bandaged, let dry first ■May be covered with a sterile bandage

**OTHER INFORMATION** ■Store at controlled room temperature ■will produce serious gastric disturbances if taken internally

**INACTIVE INGREDIENT** purified water, polihexanide hydrochloride

**J&D ETHYL ALCOHOL 70%**

**RUBBING ALCOHOL  
FIRST AID ANTISEPTIC  
AGENT ANTIBACTERIAL**

**16.9 OZ. (500mL)**

**WARNING  
FLAMMABLE**  
Keep away from heat, spark,  
Electrical, fire or flame.

**NDC: 75525-004-50**

Lot: IWC101120  
Mfg.Dt. 11/2020  
Exp.Dt. 10/22

**NOT TESTED  
ON ANIMALS**

**MADE IN COLOMBIA**

Distributed by:  
**J&D INTERNATIONAL IMPORT &  
EXPORT  
RIDGEFIELD, NJ 07657**

010000001000000

IWSC029

**TAMPER EVIDENT: DO NOT USE IF PRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

## J&D

### Ethyl Alcohol 70% - 500 ml

500 mL

NDC: 75525-004-50

Drug Facts	
<b>Active Ingredients</b>	<b>Purpose</b>
Ethyl Alcohol (70% v/v)	First aid antiseptic
<b>Uses</b>	
first aid to help prevent the risk of infection in minor cuts, scrapes, and burns.	
<b>WARNINGS</b>	
<b>FOR EXTERNAL USE ONLY</b>	
If taken internally, serious gastric disturbances will result.	
<b>FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.</b>	
<b>USE ONLY IN A WELL-VENTILATED AREA; FUMES MAY BE TOXIC</b>	
<b>ASK A DOCTOR BEFORE USE IF YOU HAVE</b> Deep or puncture wounds, Animal bites, or serious burns.	
<b>WHEN USING THIS PRODUCT</b> ■Do not get into eyes ■Do not apply over large areas of the body ■Do not use longer than one week unless directed by a doctor	
<b>STOP USE AND ASK A DOCTOR IF</b> condition persists or gets worse.	
<b>KEEP OUT OF REACH OF CHILDREN.</b> In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately (1-800-222-1222)	
<b>DIRECTIONS</b> ■Clean affected area ■Apply small amount of this product on the area 1-3 times daily ■If bandaged, let dry first ■May be covered with a sterile bandage	
<b>OTHER INFORMATION</b> ■Store at controlled room temperature ■will produce serious gastric disturbances if taken internally	
<b>INACTIVE INGREDIENT</b> purified water, polihexanide hydrochloride	



**ETHYL**  
**ALCOHOL**  
**70%**



**WARNING**  
**FLAMMABLE**

Keep away from heat, spark, Electrical, fire or flame.

**RUBBING ALCOHOL**  
**FIRST AID ANTISEPTIC**  
**AGENT ANTIBACTERIAL**

**16.9 OZ. (500mL)**

NDC: 75525-004-50

Lot: IWC101120  
Mfg.Dt. 11/2020  
Exp.Dt. 10/22



MADE IN COLOMBIA

Distributed by:  
**J&D INTERNATIONAL IMPORT & EXPORT**  
RIDGEFIELD, NJ 07657



IWSC029

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**J&D**  
**Ethyl Alcohol 70% - 1000 ml**

1000 mL

NDC: 75525-004-10

Drug Facts	
<b>Active Ingredients</b>	<b>Purpose</b>
Ethyl Alcohol (70% v/v)	First aid antiseptic
<b>Uses</b>	
first aid to help prevent the risk of infection in minor cuts, scrapes, and burns.	
<b>WARNINGS</b>	
<b>FOR EXTERNAL USE ONLY</b>	
If taken internally, serious gastric disturbances will result.	
<b>FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.</b>	
<b>USE ONLY IN A WELL-VENTILATED AREA; FUMES MAY BE TOXIC</b>	
<b>ASK A DOCTOR BEFORE USE IF YOU HAVE</b> Deep or puncture wounds, Animal bites, or serious burns.	
<b>WHEN USING THIS PRODUCT</b> ■Do not get into eyes ■Do not apply over large areas of the body ■Do not use longer than one week unless directed by a doctor	
<b>STOP USE AND ASK A DOCTOR IF</b> condition persists or gets worse.	
<b>KEEP OUT OF REACH OF CHILDREN.</b> In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately (1-800-222-1222)	
<b>DIRECTIONS</b> ■Clean affected area ■Apply small amount of this product on the area 1-3 times daily ■If bandaged, let dry first ■May be covered with a sterile bandage	
<b>OTHER INFORMATION</b> ■Store at controlled room temperature ■will produce serious gastric disturbances if taken internally	
<b>INACTIVE INGREDIENT</b> purified water, polihexanide hydrochloride	



**ETHYL**  
**ALCOHOL**  
**70%**



**WARNING**  
**FLAMMABLE**

Keep away from heat, spark, Electrical, fire or flame.

**RUBBING ALCOHOL**  
**FIRST AID ANTISEPTIC**  
**AGENT ANTIBACTERIAL**

**33.8 OZ. (1,000mL)**

NDC: 75525-004-10

Lot: IWC101120  
Mfg.Dt. 11/2020  
Exp.Dt. 10/22



MADE IN COLOMBIA

Distributed by:  
**J&D INTERNATIONAL IMPORT & EXPORT**  
RIDGEFIELD, NJ 07657



IWSC030

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**J&D**  
**Ethyl Alcohol 70% - 1 Gal - 3800 ml**

1 Gal - 3800 ml

NDC: 75525-004-01

Drug Facts	
<b>Active Ingredients</b>	<b>Purpose</b>
Ethyl Alcohol (70% v/v)	First aid antiseptic
<b>Uses</b>	
first aid to help prevent the risk of infection in minor cuts, scrapes, and burns.	
<b>WARNINGS</b>	
<b>FOR EXTERNAL USE ONLY</b>	
If taken internally, serious gastric disturbances will result.	
<b>FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.</b>	
<b>USE ONLY IN A WELL-VENTILATED AREA; FUMES MAY BE TOXIC</b>	
<b>ASK A DOCTOR BEFORE USE IF YOU HAVE</b> Deep or puncture wounds, Animal bites, or serious burns.	
<b>WHEN USING THIS PRODUCT</b> ■Do not get into eyes ■Do not apply over large areas of the body ■Do not use longer than one week unless directed by a doctor	
<b>STOP USE AND ASK A DOCTOR IF</b> condition persists or gets worse.	
<b>KEEP OUT OF REACH OF CHILDREN.</b> In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately (1-800-222-1222)	
<b>DIRECTIONS</b> ■Clean affected area ■Apply small amount of this product on the area 1-3 times daily ■If bandaged, let dry first ■May be covered with a sterile bandage	
<b>OTHER INFORMATION</b> ■Store at controlled room temperature ■will produce serious gastric disturbances if taken internally	
<b>INACTIVE INGREDIENT</b> purified water, polihexanide hydrochloride	

# ETHYL ALCOHOL 70%

**WARNING FLAMMABLE**  
Keep away from heat, spark, Electrical, fire or flame.

**RUBBING ALCOHOL  
FIRST AID ANTISEPTIC  
AGENT ANTIBACTERIAL**

**1 gal (3.800 mL)**

NDC: 75525-004-01

Lot: IWC101120  
Mfg.Dt. 11/2020  
Exp.Dt. 10/22

MADE IN COLOMBIA

Distributed by:  
**J&D INTERNATIONAL IMPORT & EXPORT**  
RIDGEFIELD, NJ 07657

IWSC031

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**J&D**  
**Ethyl Alcohol 70% - 5 Gal- 18.9 L**  
 5 Gal- 18.9 L  
 NDC: 75525-004-05

Drug Facts	
<b>Active Ingredients</b>	<b>Purpose</b>
Ethyl Alcohol (70% v/v)	First aid antiseptic
<b>Uses</b>	
first aid to help prevent the risk of infection in minor cuts, scrapes, and burns.	
<b>WARNINGS</b>	
<b>FOR EXTERNAL USE ONLY</b>	
If taken internally, serious gastric disturbances will result.	
<b>FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.</b>	
<b>USE ONLY IN A WELL-VENTILATED AREA; FUMES MAY BE TOXIC</b>	
<b>ASK A DOCTOR BEFORE USE IF YOU HAVE</b> Deep or puncture wounds, Animal bites, or serious burns.	
<b>WHEN USING THIS PRODUCT</b> ■Do not get into eyes ■Do not apply over large areas of the body ■Do not use longer than one week unless directed by a doctor	
<b>STOP USE AND ASK A DOCTOR IF</b> condition persists or gets worse.	
<b>KEEP OUT OF REACH OF CHILDREN.</b> In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately (1-800-222-1222)	
<b>DIRECTIONS</b> ■Clean affected area ■Apply small amount of this product on the area 1-3 times daily ■If bandaged, let dry first ■May be covered with a sterile bandage	
<b>OTHER INFORMATION</b> ■Store at controlled room temperature ■will produce serious gastric disturbances if taken internally	
<b>INACTIVE INGREDIENT</b> purified water, polihexanide hydrochloride	

# ETHYL ALCOHOL 70%

**WARNING FLAMMABLE**  
Keep away from heat, spark, Electrical, fire or flame.

**RUBBING ALCOHOL  
FIRST AID ANTISEPTIC  
AGENT ANTIBACTERIAL**

**5 gal (18.9 L)**

NDC: 75525-004-05

Lot: IWC101120  
Mfg.Dt. 11/2020  
Exp.Dt. 10/22

MADE IN COLOMBIA

Distributed by:  
**J&D INTERNATIONAL IMPORT & EXPORT**  
RIDGEFIELD, NJ 07657


IWSC032

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**J&D**  
**Ethyl Alcohol 70% - 55 Gal- 208 L**  
 55 Gal- 208 L  
 NDC: 75525-004-55

**Drug Facts**

<b>Active Ingredients</b> Ethyl Alcohol (70% v/v)	<b>Purpose</b> First aid antiseptic
<b>Uses</b> first aid to help prevent the risk of infection in minor cuts, scrapes, and burns.	
<b>WARNINGS</b> <b>FOR EXTERNAL USE ONLY</b> If taken internally, serious gastric disturbances will result. <b>FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.</b> <b>USE ONLY IN A WELL-VENTILATED AREA; FUMES MAY BE TOXIC.</b>	
<b>ASK A DOCTOR BEFORE USE IF YOU HAVE</b> Deep or puncture wounds, Animal bites, or serious burns.	
<b>WHEN USING THIS PRODUCT</b> ■ Do not get into eyes ■ Do not apply over large areas of the body ■ Do not use longer than one week unless directed by a doctor	
<b>STOP USE AND ASK A DOCTOR IF</b> condition persists or gets worse.	
<b>KEEP OUT OF REACH OF CHILDREN.</b> In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately (1-800-222-1222)	
<b>DIRECTIONS</b> ■ Clean affected area ■ Apply small amount of this product on the area 1-3 times daily ■ If bandaged, let dry first ■ May be covered with a sterile bandage	
<b>OTHER INFORMATION</b> ■ Store at controlled room temperature ■ Will produce serious gastric disturbances if taken internally	
<b>INACTIVE INGREDIENT</b> purified water, polihexanide hydrochloride	

**ETHYL ALCOHOL 70%****RUBBING ALCOHOL  
FIRST AID ANTISEPTIC  
AGENT ANTIBACTERIAL****55 gal (208 L)**

NDC: 75525-004-55

Lot: IWC101120  
Mfg.Dt. 11/2020  
Exp.Dt. 10/22

MADE IN COLOMBIA

Distributed by:  
**J&D INTERNATIONAL IMPORT &  
EXPORT**  
RIDGEFIELD, NJ 07657

IWSC033

TAMPER EVIDENT: DO NOT USE IF PRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**JD ANTISEPTIC ALCOHOL 70%**

alcohol liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:75525-004
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	72.9 mL in 100 mL

**Inactive Ingredients**

Ingredient Name	Strength
POLIHESANIDE HYDROCHLORIDE (UNII: 4XI6112496)	0.1 mL in 100 mL
WATER (UNII: 059QF0KO0R)	27 mL in 100 mL

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75525-004-50	500 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2020	
2	NDC:75525-004-10	1000 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2020	
3	NDC:75525-004-01	3800 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2020	
4	NDC:75525-004-05	18900 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2020	
5	NDC:75525-004-55	208000 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2020	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	10/01/2020	

**Labeler** - IWANNA CORP SAS (883223177)

**Registrant** - MASTER CUSTOMS BROKER (054631555)

## Establishment

Name	Address	ID/FEI	Business Operations
IWANNA CORP SAS		883223177	manufacture(75525-004)

Revised: 12/2020

IWANNA CORP SAS