SALINE LAXATIVE- saline laxative enema Major Pharmaceuticals

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

MAJOR® SALINE ENEMA LAXATIVE

Drug Facts

| Active ingredients (in each 118mL delivered dose) | Purpose |
|---|-----------------|
| Dibasic Sodium Phosphate 7 g | Saline laxative |
| Monobasic Sodium Phosphate | |
| 19 g | Saline laxative |

Use

- relieves occasional constipation
- this product generally produces bowel movement in 1 to 5 minutes

Warnings

Dosage warning

Using more than one enema in 24 hours can be harmful.

For rectal use only.

Do not use

• on children under 2 years of age

Ask a doctor before use if you

- have already used a laxative for more than 3 days
- have kidney disease, have heart problems, or are dehydrated
- are 55 years of age or older
- are on a sodium-restricted diet
- have abdominal pain, nausea, or vomiting
- have a sudden change in bowel habits lasting more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

When using this product

 do not use more than directed. Serious side effects may occur from excess dosage do not use for more than 3 days, without asking a doctor

Stop use and ask a doctor if

- you have rectal bleeding
- you have no bowel movement within 30 minutes of enema use
- you have symptoms of dehydration (thirstiness, dizziness, vomiting, urinating less often than normal)

These symptoms may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away: 800-222-1222.

Directions (or as directed by a doctor)

Single daily dosage (per 24 hours)

Do not use if taking another sodium phosphate product.

Do not use more unless directed by a doctor. See Warnings.

| Adults and children 12 years of age and older | 1 bottle once daily |
|---|---------------------|
| Children 2 to under 12 | Use Pediatric |
| years of age | Enema |
| Children under 2 years of age | DO NOT USE |

Other information

- each 118 ml contains sodium 4.3 g
- store at 15 to 30°C (59 to 86°F)
- additional liquids by mouth are recommended while using this product
- Tamper Evident: DO NOT USE IF TOP OR BOTTOM FLAP OF CARTON IS TORN OR MISSING.

Inactive ingredients

benzalkonium chloride, disodium EDTA, purified water

Questions or comments?

(800) 616-2471

Distributed by:

MAJOR® PHARMACEUTICALS

Livonia, MI 48152

PRINCIPAL DISPLAY PANEL - 133 mL Bottle Carton

 $\mathsf{MAJOR}^{\, \mathbb{R}}$

NDC 0904-6320-78

Compare to the active ingredients of Fleet® Saline Enema*

Ready-to-Use Enema Saline Laxative

For relief of occasional constipation

Not made with natural rubber latex

4.5 FL OZ (133 mL)

1 Bottle

Do not use it top or bottom flap lamper Resistant

MAJOR

CAUTION:
REMOVE
PROTECTIVE
SHIELD
BEFORE
INSERTING.

Ready-to-Use :nema Saline Laxative

Remove protective shield from enema tip before inserting.

PL4000B V17

Hold bottle upright, grasping bottle cap with fingers. Grasp protective shield with other hand and pull gently to remove.



Positions for using this enema

Left-side position: Lie on left side with knees bent and arms resting comfortably.



Knee-chest position: Kneel, then lower head and chest forward until left side of face is resting on surface with left arm folded comfortably.



How to use this enema

Administering enema:

- With steady pressure, gently insert enema tip into rectum with a slight sideenema up into recum with a signit sole-to-side movement, with tip pointing toward navel. Insertion may be easier if person receiving enema bears down, as if having a bowel movement. Doing so helps relax muscles around anus.
- . Do not force enema tip into rectum. Stop using if tip is hard to insert. Forcing the tip into the rectum can cause injury (especially if you have hemorrhoids). If enema tip causes rectal bleeding or pain, get immediate medical care.
- . Squeeze bottle until nearly all liquid is gone. It is not necessary to empty bottle completely, as it contains more liquid than needed.
- Remove tip from rectum and stay in position until urge to evacuate is strong (usually 1 to 5 minutes). If no urge is felt after 5 minutes of using, try to empty bowel. Call a doctor promptly if no liquid comes out of the rectum after 30 minutes because dehydration could occur.

Bottle, cap, and shield are NOT MADE WITH NATURAL RUBBER LATEX MADE IN



THE USA

Rev. 02/21 M-139 Re-order No. 700202



MAJOR[®]

NDC 0904-6320-78

Compare to the active ingredients of Fleet* Saline Enema*

Ready-to-Use Enema

Saline Laxative

For relief of occasional constipation

Not made with natural rubber latex

Drug Facts

Active ingredients (In each 118mL delivered Dibasic Sodium Purpose

CARTON SEALEI FOR SAFETY. DO NOT USE IF TOP OR BOTTOM FLAP OF CARTON IS TORN OR MISSING.

Saline laxative Phosphale 7 g Monobasic Sodium Phosphale 19 g. .Saline laxative

Use

relieves occasional constipation this product generally produces bowel movement in 1 to 5 minutes

Warnings

Dosage warning: Using more than one enema in 24 hours can be harmful.

For rectal use only.

- Do not use on children under 2 years of age
- on crimone interior 2 years or age
 Asik a doctor before use if you
 have already used a lexative for more than 3 days
 have kidney disease, have heart problems, or are dehydrated
 are 55 years of age or older
 are on a sodium-restricted diel
 have abdominal pain, nausea, or vemitting
 have a sudden change in bowel habits lasting more than 2 weeks

Ask a doctor or pharmacist before use if you are laking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

- When using this product
 do not use more than directed. Serious side effects may occur from excess dosage do not use for more than 3 days, without
- asking a doctor

- Stop use and ask a doctor if

 you have rectal bleeding

 you have no bowel movement within 30 minutes of enema use

 you have symptoms of dehydration (thirstiness, dizziness, wmitting, urtnating less often than normal)

These symptoms may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Polson Control Center right away: 800-222-1222.

Directions (or as directed by a doctor, Single daily dosage (per 24 hours) Do not use if taking another sodium phosphate product, Do not use more unless directed by a doctor. See Warnings.

| Adults and children | 1 bottle | |
|----------------------------------|---------------|--|
| 12 years of age and older | once daily | |
| Children 2 to | Use Pediatric | |
| under 12 years of age | Enema | |
| Children under 2 years of age | DO NOT USE | |

Drug Facts (continued)

Other information

- each 118 mi contains sodium 4.3 g
 store at 15 to 30°C (59 to 96°F)
 additional fiquids by mouth are recommended white using this product
 Tamper Evident: DO NOT USE IF TOP OR 90 TOM FLAP OF CARTON IS TORN OR MISSING.

Inactive ingredients benzalkonium chloride, disodium EDTA, purified water

Questions or comments? (800) 616-2471

This product is not manufactured or distributed by C.B. Fleet Company Inc., owner of the registered trademark Fleet®.

Distributed by: MAJOR® PHARMACEUTICALS Livonia, MI 48152



Image not actual size

4.5 FL OZ (133 mL)

SALINE LAXATIVE

saline laxative enema

| Product Information | | | | |
|-------------------------|----------------|--------------------|---------------|--|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:0904-6320 | |
| Route of Administration | RECTAL | | | |

| Active Ingredient/Active Moiety | | | | |
|--|--|-------------------|--|--|
| Ingredient Name | Basis of Strength | Strength | | |
| SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM (UNII: GR686LBA74) (PHOSPHATE ION - UNII:NK08V8K8HR) | SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM | 7 g in 133 mL | | |
| SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW) (PHOSPHATE ION - UNII:NK08V8K8HR) | SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM | 19 g in 133 mL | | |

| Inactive Ingredients | | | |
|--|----------|--|--|
| Ingredient Name | Strength | | |
| BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) | | | |
| EDETATE DISODIUM (UNII: 7FLD91C86K) | | | |
| WATER (UNII: 059QF0KO0R) | | | |

| l | Packaging | | | | |
|---|-----------|----------------------|---|-------------------------|-----------------------|
| | # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| | 1 | NDC:0904-6320- 78 | 133 mL in 1 BOTTLE; Type 0: Not a Combination Product | 07/11/2013 | |

| Marketing Information | | | |
|----------------------------|---|-------------------------|-----------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC MONOGRAPH NOT FINAL | part334 | 07/11/2013 | |
| | | | |

Labeler - Major Pharmaceuticals (191427277)

| Establishment | | | | |
|----------------|---------|-----------|----------------------------|--|
| Name | Address | ID/FEI | Business Operations | |
| NATUREPLEX LLC | | 062808196 | MANUFACTURE(0904-6320) | |

Revised: 3/2021 Major Pharmaceuticals