

PAIN RELIEF- acetaminophen tablet
Meijer Distribution Inc

Meijer 44-148 Delisted

Active ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - toothache
 - muscular aches
 - headache
 - the common cold
 - minor pain of arthritis
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over
 - take 2 tablets every 6 hours while symptoms last
 - do not take more than 6 tablets in 24 hours, unless directed by a doctor
 - do not take for more than 10 days unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, povidone, sodium starch glycolate, stearic acid

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday- Friday

Principal Display Panel

NDC 41250-148-03

meijer®

Compare to
Extra Strength Tylenol®
active ingredient*

EXTRA STRENGTH
pain relief
500 mg each
Acetaminophen | Pain Reliever | Fever Reducer

10 Tablets

actual size

TAMPER EVIDENT: Use Only if This Blister is Intact

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Extra Strength Tylenol®. 50844 REV0621D14803

DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com

meijer

NDC 41250-148-03
Compare to
Extra Strength Tylenol®
active ingredient*

EXTRA STRENGTH
pain relief
500 mg each
Acetaminophen | Pain Reliever | Fever Reducer

10 Tablets 

TAMPER EVIDENT: Use Only if This Blister is Intact

No Print / No Varnish Area
Lot & Exp. Area

Drug Facts	RETAIN VIAL CARD FOR COMPLETE PRODUCT INFORMATION	Purpose
Active ingredient (in each tablet) Acetaminophen 500 mg		Pain reliever/fever reducer
Uses temporarily relieves minor aches and pains due to: ■ backache ■ toothache ■ muscular aches ■ headache ■ the common cold ■ minor pain of arthritis ■ premenstrual and menstrual cramps ■ temporarily reduces fever		
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product Ask a doctor before use if you have liver disease. Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. Stop use and ask a doctor if ■ redness or swelling is present ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days		
Drug Facts (continued) ■ new symptoms occur These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.		
Directions ■ do not take more than directed ■ adults and children 12 years and over ■ take 2 tablets every 6 hours while symptoms last ■ do not take more than 6 tablets in 24 hours, unless directed by a doctor ■ do not take for more than 10 days unless directed by a doctor ■ children under 12 years: ask a doctor		
Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package		
Inactive ingredients corn starch, povidone, sodium starch glycolate, stearic acid		
Questions or comments? Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday		
* This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Extra Strength Tylenol®. 50844 REV0621D14803		
DIST. BY MEIJER DISTRIBUTION, INC. GRAND RAPIDS, MI 49544 www.meijer.com		
PID 3767074  7 13733 43292 4		

Meijer 44-148

PAIN RELIEF

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-148
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	12mm
Flavor		Imprint Code	44;148
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-148-03	1 in 1 PACKAGE	01/21/1993	09/30/2026
1		10 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/21/1993	09/30/2026

Labeler - Meijer Distribution Inc (006959555)

Establishment

Name	Address	ID/FEI	Business Operations
-------------	----------------	---------------	----------------------------

LNK International, Inc.		038154464	pack(41250-148)
-------------------------	--	-----------	-----------------

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(41250-148)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(41250-148)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(41250-148)

Revised: 2/2026

Meijer Distribution Inc