

**ALL DAY ALLERGY D- cetirizine hydrochloride, pseudoephedrine hydrochloride tablet, film coated, extended release**  
**L. Perrigo Company**

-----  
**Perrigo All Day Allergy-D Drug Facts**

**Active ingredients (in each extended release tablet)**

Cetirizine HCl 5 mg

Pseudoephedrine HCl 120 mg

**Purpose**

Antihistamine

Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- thyroid disease

- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are**

taking tranquilizers or sedatives.

**When using this product**

- **do not use more than directed**
- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not break or chew tablet; swallow tablet whole

adults and children 12 years and over	take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
adults 65 years and over	ask a doctor
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

## Other information

- store between 20° to 25°C (68° to 77°F)
- **do not use if blister unit is broken or torn**
- see side panel for lot number and expiration date
- meets USP *Dissolution Test 2*

## Inactive ingredients

colloidal silicon dioxide, hypromellose, lactose monohydrate, low-substituted hydroxypropyl cellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

## Questions or comments?

call **1-800-719-9260**

## Package/Label Principal Display Panel

Compare to Zyrtec-D® active ingredients

ORIGINAL PRESCRIPTION STRENGTH

ALL DAY allergy-D

Cetirizine Hydrochloride and Pseudoephedrine Hydrochloride Extended Release tablets,  
5 mg/120 mg

Antihistamine/Nasal Decongestant

ALLERGY + SINUS

INDOOR + OUTDOOR ALLERGIES

12 hour

12 HOUR RELIEF

Sneezing

Itchy, Watery Eyes

Runny Nose

Itchy Nose or Throat

Actual Size

NASAL CONGESTION + SINUS PRESSURE

24 EXTENDED RELEASE TABLETS

**Drug Facts (continued)**

**Other information** ■ store between 20° to 25°C (68° to 77°F)

■ do not use if blister unit is broken or torn

■ see side panel for lot number and expiration date

■ meets USP Dissolution Test 2

**Active ingredients** cetirizine hydrochloride, pseudoephedrine hydrochloride, low-substituted hydroxypropyl cellulose, magnesium stearate, microcrystalline cellulose, polyvinyl alcohol, talc, titanium dioxide

**Questions or comments?** call 1-800-719-9260



**Drug Facts**

**Warnings**

Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine, (H1) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores free breathing through the nose

**Directions**

adults and children 12 years and over ■ take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.

adults and children 6 to 11 years ■ ask a doctor

children 2 to 5 years ■ ask a doctor

12 years of age ■ ask a doctor

or kidney disease ■ ask a doctor

**Directions**

do not break or chew tablet; swallow tablet whole

help or contact a Poison Control Center right away, (1-800-222-1222)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Warnings**

If pregnant or breast-feeding:

- breast-feeding not recommended
- pregnant; ask a health professional before use.

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

**Active ingredients** (in each extended release tablet)

Pseudoephedrine HCl 120 mg ..... Nasal decongestant

Cetirizine HCl 5 mg ..... Antihistamine

**Purpose**

Ask a doctor or pharmacist before use if you are taking

benzodiazepines or sedatives

do not use more than directed

avoid alcoholic drinks

stomach upset may occur

alcohol, sedatives, and tranquilizers may increase drowsiness

be careful when driving a motor vehicle or operating machinery

an allergic reaction to this product occurs. Seek medical help right away.

symptoms do not improve within 7 days or are accompanied by fever

**Directions**

adults and children 12 years and over ■ take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.

adults and children 6 to 11 years ■ ask a doctor

children 2 to 5 years ■ ask a doctor

12 years of age ■ ask a doctor

or kidney disease ■ ask a doctor

Convenient Reclosing Tab

Distributed by Perrigo® Allegan, MI 49010 perigo.com

OPEN OTHER END

14762 55 C3

ORIGINAL PRESCRIPTION STRENGTH

# ALL DAY allergy-D

NDC 0113-2147-62

ORIGINAL PRESCRIPTION STRENGTH

# ALL DAY allergy-D

Cetirizine Hydrochloride and Pseudoephedrine Hydrochloride  
Extended Release Tablets, 5mg/120 mg  
Antihistamine/Nasal Decongestant

Compare to Zyrtec-D® active ingredients

## ALLERGY + SINUS

INDOOR + OUTDOOR ALLERGIES



12 HOUR RELIEF

- Sneezing
- Itchy, Watery Eyes
- Runny Nose
- Itchy Nose or Throat

NASAL CONGESTION + SINUS PRESSURE



Actual Size

24 EXTENDED RELEASE TABLETS

# ALL DAY ALLERGY D

cetirizine hydrochloride, pseudoephedrine hydrochloride tablet, film coated, extended release

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0113-2147
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CETIRIZINE HYDROCHLORIDE</b> (UNII: 64O047KTOA) (CETIRIZINE - UNII:YO7261ME24)	CETIRIZINE HYDROCHLORIDE	5 mg
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>LOW-SUBSTITUTED HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 2165RE0K14)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	L147
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-2147-62	24 in 1 CARTON	10/02/2025	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
ANDA	ANDA210719	10/02/2025	

**Labeler** - L. Perrigo Company (006013346)

Revised: 10/2025

L. Perrigo Company