

**MEDIQUE CHLORPHEN- chlorpheniramine maleate tablet**  
**Unifirst First Aid Corporation**

-----  
**Medique Chlorphen**

***Drug Facts***

***Active ingredient (in each tablet)***

Chlorpheniramine maleate 4 mg

***Purpose***

Antihistamine

***Uses***

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

***Warnings***

Do not use to make a child sleepy.

**Ask a doctor before use if you have**

- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before us if you are** taking sedatives or tranquilizers.

**When using this product**

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**If you are pregnant or breast-feeding,** ask a health professional before use.

**Keep out of the reach of children.** In case of overdose, get medical help or contact a Poison Control

Center (1-800-222-1222) right away.

## ***Directions***

**Adults and children (12 years and older):** 1 tablet every 4 to 6 hours. Do not take more than 6 tablets in 24 hours.

**Children under 12 years:** Do not give to children under 12 years of age

## ***Other information***

- tamper evident sealed packets
- do not use any opened or torn packets
- store at 77°F (25°C); excursions permitted between 59°-86°F (15°-30°C)
- protect from excessive moisture
- use by expiration date on packet

## ***Inactive ingredients***

corn starch, D&C yellow #10 aluminum lake, lactose anhydrous, magnesium stearate, microcrystalline cellulose

***Questions or comments?*** Call 1-800-634-7680

## **Medique Chlorphen Label**

Medique ®

Chlorphen

Allergy and Hay Fever Relief

Alivio a las alergias y la fiebre del heno

Antihistamine • Chlorpheniramine Maleate 4mg

Antihistminico • Maleato de Clorfeniramina 4mg

Pull To Open

Tire Para Abrir

250 Tablets

(250 x 1)

Tamper Evident Unit Dose Packets

Empaquetado con Sellado Evidente en Dosis Unitarias

**Drug Facts**

<b>Active ingredient (in each tablet)</b> Chlorpheniramine maleate, 4 mg	<b>Purpose</b> Antihistaminic				
<b>Uses</b> Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: <ul style="list-style-type: none"> <li>■ runny nose</li> <li>■ itchy, watery eyes</li> <li>■ sneezing</li> <li>■ itching of the nose or throat</li> </ul>					
<b>Warnings</b> Do not use to make a child sleepy.					
Ask a doctor before use if you have: <ul style="list-style-type: none"> <li>■ glaucoma</li> <li>■ a breathing problem such as emphysema or chronic bronchitis</li> <li>■ trouble urinating due to an enlarged prostate gland</li> </ul>					
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.					
<b>When using this product</b> <ul style="list-style-type: none"> <li>■ drowsiness may occur</li> <li>■ avoid alcoholic beverages</li> <li>■ alcohol, sedatives and tranquilizers may increase drowsiness</li> <li>■ use caution when driving a motor vehicle or operating machinery</li> <li>■ excitability may occur, especially in children</li> </ul>					
If you are pregnant or breast feeding, ask a health professional before use.					
Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.					
<b>Directions</b> <table border="0" style="width: 100%;"> <tr> <td style="width: 60%;">Adults and children 12 years and older</td> <td>1 tablet every 4 to 6 hours. Do not take more than 6 tablets in 24 hours.</td> </tr> <tr> <td>Children under 12 years</td> <td>Do not give to children under 12 years of age.</td> </tr> </table>		Adults and children 12 years and older	1 tablet every 4 to 6 hours. Do not take more than 6 tablets in 24 hours.	Children under 12 years	Do not give to children under 12 years of age.
Adults and children 12 years and older	1 tablet every 4 to 6 hours. Do not take more than 6 tablets in 24 hours.				
Children under 12 years	Do not give to children under 12 years of age.				
<b>Other information</b> <ul style="list-style-type: none"> <li>■ tamper-evident sealed packets</li> <li>■ do not use any opened or torn packets</li> <li>■ store at 20°C (68°F); excursions permitted between 15°-30°C (59°-86°F)</li> <li>■ protect from excessive moisture</li> <li>■ use for respiratory relief only</li> </ul>					
<b>Inactive ingredients</b> corn starch, D&C yellow #10 aluminum lake, lactose anhydrous, magnesium stearate, microcrystalline cellulose					
<b>Questions or comments?</b> 1-800-634-7680					

Retain carton for complete product information

**Medique**  
**Chlorpheniramine Maleate 4 mg**  
**Allergy & Hay Fever Relief**  
Alivio a las alergias y la fiebre del heno

✓ Antihistamine • Chlorpheniramine Maleate 4 mg  
Antihistaminico • Maleato de Clorfeniramina 4 mg

Pull to Open  
The Flip-Top

250  
Tablets  
(250 x 1)

Tamper Evident Unit Dose Packets  
Empaquetado con Sellado Evidente en Dosis Unitarias

**Información del Medicamento**

<b>Ingrediente activo (en cada tableta)</b> Maleato de clorfeniramina, 4 mg	<b>Propiedad</b> Antihistaminico				
<b>Usos</b> para el alivio temporal de los siguientes síntomas los cuales se asocian con fiebre del heno y otras alergias del sistema respiratorio superior: <ul style="list-style-type: none"> <li>■ goteo de la nariz</li> <li>■ ojos con comezón o llorosos</li> <li>■ estornudos</li> <li>■ congestión de la nariz o garganta</li> </ul>					
<b>Advertencias</b> No se debe usar para hacer a un niño con sueño.					
Consulte con el médico antes de consumirlo si usted tiene: <ul style="list-style-type: none"> <li>■ glaucoma</li> <li>■ problema al respirar como asma o bronquitis crónica</li> <li>■ dificultad al orinar debido al crecimiento de la glándula prostática</li> </ul>					
Consulte con su médico o farmacéutico antes de administrarlo, si usted está tomando sedativos o tranquilizantes.					
<b>Cuando usted consuma este producto</b> <ul style="list-style-type: none"> <li>■ puede ocurrir somnolencia</li> <li>■ evite el consumo de bebidas alcohólicas</li> <li>■ el alcohol, los sedativos y tranquilizantes pueden incrementar los efectos de somnolencia</li> <li>■ sea cuidadoso al manejar un vehículo motorizado o al operar maquinaria</li> <li>■ puede provocar excitación, en especial en menores de edad</li> </ul>					
Si usted está embarazada o en período de lactancia, consulte la opinión de un médico antes de consumir este producto.					
Manténgalo lejos y fuera del alcance de los niños. En caso de ocurrir una sobredosis, comuníquese de inmediato con un médico o llame a un centro de control de envenenamiento (1-800-222-1222).					
<b>Instrucciones</b> <table border="0" style="width: 100%;"> <tr> <td style="width: 60%;">Adultos y niños (de 12 años y mayores)</td> <td>Turne 1 tableta cada 4 a 6 horas. No consuma más de 6 tabletas en 24 horas.</td> </tr> <tr> <td>Niños menores de 12 años</td> <td>No se administre a niños menores de 12 años de edad.</td> </tr> </table>		Adultos y niños (de 12 años y mayores)	Turne 1 tableta cada 4 a 6 horas. No consuma más de 6 tabletas en 24 horas.	Niños menores de 12 años	No se administre a niños menores de 12 años de edad.
Adultos y niños (de 12 años y mayores)	Turne 1 tableta cada 4 a 6 horas. No consuma más de 6 tabletas en 24 horas.				
Niños menores de 12 años	No se administre a niños menores de 12 años de edad.				
<b>Otra información</b> <ul style="list-style-type: none"> <li>■ empaquetado con sellado evidente</li> <li>■ no use ningún paquete abierto o roto</li> <li>■ almacenar a 20°C (68°F); excursiones permitidas entre 15°-30°C (59°-86°F)</li> <li>■ evitar exposición a humedad excesiva</li> <li>■ usar antes de la fecha de expiración en el paquete</li> </ul>					
<b>Ingredientes inactivos</b> almidón de maíz, amarillo 10 y C #10 lacas de aluminio, lactosa anhidra, estearato de magnesio, celulosa microcristalina					
<b>¿Tiene alguna pregunta o comentario?</b> Llame al 1-800-634-7680					

Conserve la caja para obtener la información completa del producto

**MEDIQUE CHLORPHEN**

chlorpheniramine maleate tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-169
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9J9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3S5YLH9PMK)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6130)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	

**Product Characteristics**

<b>Color</b>	yellow	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	44;194
<b>Contains</b>			

**Packaging**

Marketing Start      Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-169-46	1 in 1 PACKET; Type 0: Not a Combination Product	11/25/2020	
2	NDC:47682-169-64	24 in 1 BOX	11/25/2020	
2		1 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-169-48	250 in 1 BOX	11/25/2020	
3		1 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/25/2020	

**Labeler** - Unifirst First Aid Corporation (832947092)

Revised: 9/2025

Unifirst First Aid Corporation