

MEDIQUE MEDI-MECLIZINE- meclizine hcl tablet
Unifirst First Aid Corporation

Medique Medi-Meclizine

Drug Facts

Active ingredient (in each tablet)

Meclizine Hydrochloride 25 mg

Purpose

Antiemetic

Uses

for the prevention and treatment of the nausea, vomiting, or dizziness associated with motion sickness.

Warnings

Do not use for children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- glaucoma
- a breathing problems such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **Do not take more than directed**
- take the first dose ½ hour to 1 hour before starting activity

Adults and children: (12 years and older) 1 to 2 tablets once daily, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information

- tamper-evident sealed packets
- do not use any opened or torn packets
- store at 77°F (25°C); excursions permitted between 59- 86° F (15-30°C)
- protect from heat and humidity
- use by expiration date on packet

Inactive ingredients

corn starch, D&C yellow # 10 aluminum lake, lactose monohydrate, magnesium stearate, silicon dioxide

Questions or comments? Call 1-800-634-7680

Medique Medi-Meclizine Label

Medique®

Medi-Meclizine

All Day Relief

Alivio que Perdura Todo el Dia

Pull to Open

Tire Para Abrir

For the Prevention and Treatment of Motion Sickness • Meclizine HCl 25mg

Para la Prevencion y el Tratamiento del Mareo "Mal del Viajero" • Hidrocloruro de Meclizina 25mg

100 Tablets

(50 x 2)

Tamper Evident Unit Dose Packets

Empaquetado con Sellado Evidente en Dosis Unitarias

<table border="1"><tr><td>Drug Facts</td><td>Drug Facts (continued)</td></tr><tr><td>Active ingredient (in each tablet) Purpose Meclizine Hydrochloride, 25 mg Antiemetic</td><td>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center at 1-800-222-1222, right away.</td></tr><tr><td>Uses for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.</td><td>Directions Do not take more than directed. ■ Do not take more than directed. ■ Take the first dose 1/2 hour to 1 hour before starting activity.</td></tr><tr><td>Warnings Do not use for children under 12 years of age unless directed by a doctor. ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis</td><td>Other information ■ tamper-evident sealed packets ■ do not use any opened or torn packets ■ store at 77°F (25°C); excursions permitted between 59-86°F (15-30°C) ■ protect from heat and humidity ■ use by expiration date on packet</td></tr><tr><td>Ask a doctor or pharmacist if you have ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis</td><td>Inactive ingredients corn starch, D&C yellow #10 aluminum lake, lactose monohydrate, magnesium stearate, silicon dioxide</td></tr><tr><td>When using this product ■ drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery</td><td>Questions or comments Call 1-800-634-7680</td></tr><tr><td>If pregnant or breast-feeding, ask a health professional before use.</td><td></td></tr></table>	Drug Facts	Drug Facts (continued)	Active ingredient (in each tablet) Purpose Meclizine Hydrochloride, 25 mg Antiemetic	Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center at 1-800-222-1222, right away.	Uses for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.	Directions Do not take more than directed. ■ Do not take more than directed. ■ Take the first dose 1/2 hour to 1 hour before starting activity.	Warnings Do not use for children under 12 years of age unless directed by a doctor. ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis	Other information ■ tamper-evident sealed packets ■ do not use any opened or torn packets ■ store at 77°F (25°C); excursions permitted between 59-86°F (15-30°C) ■ protect from heat and humidity ■ use by expiration date on packet	Ask a doctor or pharmacist if you have ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis	Inactive ingredients corn starch, D&C yellow #10 aluminum lake, lactose monohydrate, magnesium stearate, silicon dioxide	When using this product ■ drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery	Questions or comments Call 1-800-634-7680	If pregnant or breast-feeding , ask a health professional before use.		<p>Medique® Medi-Meclizine All Day Relief Alivio que Perdura Todo el Dia</p> <p>Pull to Open Tire Para Abrir</p> <p>For Prevention and Treatment of Motion Sickness • Meclizine HCl 25 mg Para la Prevencion y el Tratamiento del Mareo "Mal del Viajero" • Hidrocloruro de Meclizina 25 mg</p> <p>100 Tablets (50 x 2)</p>	<table border="1"><tr><td>Información del Medicamento</td><td>Propiedad</td></tr><tr><td>Ingrediente activo (en cada tableta) Meclizina Hidrocloruro 25 mg</td><td>Antiemético</td></tr><tr><td>Usos para la prevención y tratamiento de la náusea, vómito, laberintismo, o mareo asociados con el "mal del viajero"</td><td></td></tr><tr><td>Advertencias No se administre a niños menores de 12 años a menos que lo señale el médico. ■ dificultad para orinar debido al crecimiento de la glándula prostática ■ diabetes ■ problemas respiratorios como enfisema o bronquitis crónica</td><td></td></tr><tr><td>Consulte con el médico antes de tomar este producto al seguirlo. ■ dificultad para orinar debido al crecimiento de la glándula prostática ■ diabetes ■ problemas respiratorios como enfisema o bronquitis crónica</td><td></td></tr><tr><td>Cuando consuma este producto ■ evite el consumo de bebidas alcohólicas ■ el alcohol, los sedantes y tranquilizantes pueden aumentar los efectos de la somnolencia ■ tenga cuidado al manejar un vehículo motorizado o al operar maquinaria</td><td></td></tr><tr><td>Si está usted embarazada o en período de lactancia, consulte la opinión de un médico antes de consumir este producto.</td><td></td></tr></table>	Información del Medicamento	Propiedad	Ingrediente activo (en cada tableta) Meclizina Hidrocloruro 25 mg	Antiemético	Usos para la prevención y tratamiento de la náusea, vómito, laberintismo, o mareo asociados con el "mal del viajero"		Advertencias No se administre a niños menores de 12 años a menos que lo señale el médico. ■ dificultad para orinar debido al crecimiento de la glándula prostática ■ diabetes ■ problemas respiratorios como enfisema o bronquitis crónica		Consulte con el médico antes de tomar este producto al seguirlo. ■ dificultad para orinar debido al crecimiento de la glándula prostática ■ diabetes ■ problemas respiratorios como enfisema o bronquitis crónica		Cuando consuma este producto ■ evite el consumo de bebidas alcohólicas ■ el alcohol, los sedantes y tranquilizantes pueden aumentar los efectos de la somnolencia ■ tenga cuidado al manejar un vehículo motorizado o al operar maquinaria		Si está usted embarazada o en período de lactancia , consulte la opinión de un médico antes de consumir este producto.		<p>Medique® Medi-Meclizine</p> <p>Reorder #47933 Manufactured for Medique Products Fort Myers, FL 33967 1-800-634-7680 www.mediqueproducts.com</p> <p>4 7682 4 7933 6</p>
Drug Facts	Drug Facts (continued)																														
Active ingredient (in each tablet) Purpose Meclizine Hydrochloride, 25 mg Antiemetic	Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center at 1-800-222-1222, right away.																														
Uses for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.	Directions Do not take more than directed. ■ Do not take more than directed. ■ Take the first dose 1/2 hour to 1 hour before starting activity.																														
Warnings Do not use for children under 12 years of age unless directed by a doctor. ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis	Other information ■ tamper-evident sealed packets ■ do not use any opened or torn packets ■ store at 77°F (25°C); excursions permitted between 59-86°F (15-30°C) ■ protect from heat and humidity ■ use by expiration date on packet																														
Ask a doctor or pharmacist if you have ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis	Inactive ingredients corn starch, D&C yellow #10 aluminum lake, lactose monohydrate, magnesium stearate, silicon dioxide																														
When using this product ■ drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery	Questions or comments Call 1-800-634-7680																														
If pregnant or breast-feeding , ask a health professional before use.																															
Información del Medicamento	Propiedad																														
Ingrediente activo (en cada tableta) Meclizina Hidrocloruro 25 mg	Antiemético																														
Usos para la prevención y tratamiento de la náusea, vómito, laberintismo, o mareo asociados con el "mal del viajero"																															
Advertencias No se administre a niños menores de 12 años a menos que lo señale el médico. ■ dificultad para orinar debido al crecimiento de la glándula prostática ■ diabetes ■ problemas respiratorios como enfisema o bronquitis crónica																															
Consulte con el médico antes de tomar este producto al seguirlo. ■ dificultad para orinar debido al crecimiento de la glándula prostática ■ diabetes ■ problemas respiratorios como enfisema o bronquitis crónica																															
Cuando consuma este producto ■ evite el consumo de bebidas alcohólicas ■ el alcohol, los sedantes y tranquilizantes pueden aumentar los efectos de la somnolencia ■ tenga cuidado al manejar un vehículo motorizado o al operar maquinaria																															
Si está usted embarazada o en período de lactancia , consulte la opinión de un médico antes de consumir este producto.																															
Retain carton for complete product information	Tamper Evident Unit Dose Packets Empaquetado con Sellado Evidente en Dosis Unitarias	Conserve la caja para obtener la información completa del producto																													

MEDIQUE MEDI-MECLIZINE

meclizine hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-481
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;403
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-481-99	2 in 1 PACKET; Type 0: Not a Combination Product	12/14/2020	
2	NDC:47682-481-15	500 in 1 BOX	12/14/2020	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-481-33	50 in 1 BOX	12/14/2020	
3		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	12/14/2020	

Labeler - Unifirst First Aid Corporation (832947092)

Revised: 9/2025

Unifirst First Aid Corporation