

**MEDIQUE MEDI-MECLIZINE- meclizine hcl tablet**  
**Unifirst First Aid Corporation**

-----  
**Medique Medi-Meclizine**

***Drug Facts***

***Active ingredient (in each tablet)***

Meclizine Hydrochloride 25 mg

***Purpose***

Antiemetic

***Uses***

for the prevention and treatment of the nausea, vomiting, or dizziness associated with motion sickness.

***Warnings***

**Do not use** for children under 12 years of age unless directed by a doctor.

**Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- glaucoma
- a breathing problems such as emphysema or chronic bronchitis

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers.

**When using this product**

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- to prevent motion sickness, take the first dose ½ hour to 1 hour before starting activity
- to treat motion sickness, take at first signs of symptoms

**Adults and children: (12 years and older)** 1 to 2 tablets once daily, or as directed

by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

***Other information***

- tamper-evident sealed packets
- do not use any opened or torn packets
- store at 77°F (25°C); excursions permitted between 59- 86° F (15-30°C)
- protect from heat and humidity
- use by expiration date on packet

***Inactive ingredients***

corn starch, D&C yellow # 10 aluminum lake, lactose anhydrous, magnesium stearate, silicon dioxide

***Questions or comments?*** Call 1-800-634-7680

**Medique Medi-Meclizine Label**

Medique®

Medi-Meclizine

All Day Relief

Alivio que Perdura Todo el Dia

Pull to Open

Tire Para Abrir

For the Prevention and Treatment of Motion Sickness • Meclizine HCl 25mg


Para la Prevencion y el Tratamiento del Mareo “Mal del Viajero” • Hidrocloruro de Meclizina 25mg

100 Tablets

(50 x 2)

Tamper Evident Unit Dose Packets

Empaquetado con Sellado Evidente en Dosis Unitarias



**Medique**  
**Medi-Medicine**  
All Day Relief  
Anti-nausea Packets

✓ For Prevention and Treatment of Motion Sickness • **Medicine HC 25 mg**  
Para la Prevención y el Tratamiento del Mareo "Mal del Viaje" • **Fármaco de Acción HC 25 mg**

4793BOX  
Rev. 11/13/20A  
Mfg. 24

**Drug Facts**

**Active ingredient (in each tablet)** Purpose  
Meclizine Hydrochloride 25 mg Antiemetic

**Uses**  
for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.

**Warnings**  
Do not use for children under 12 years of age unless directed by a doctor.

**Ask a doctor before use if you have**  
■ difficulty in urination due to enlargement of the prostate gland.  
■ glaucoma  
■ breathing problem such as emphysema or chronic bronchitis

**When using this product**  
■ drowsiness may occur  
■ avoid alcoholic beverages  
■ alcohol, sedatives and tranquilizers may increase drowsiness  
■ use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

**Drug Facts (continued)**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center 1-800-222-1222 right away.

**Directions**  
■ To prevent motion sickness, take the first dose 1/2 hour to 1 hour before starting activity  
■ To treat motion sickness, take at first signs of symptoms.

**Adults and children (12 years and older)**  
Children under 12 years: Do not give to children under 12 years of age.

**Other information**  
■ tamper-evident sealed packets  
■ do not use any opened or torn packets  
■ store at 77°F (25°C), excursions permitted between 59°-86°F (15°-30°C)  
■ protect from heat and humidity  
■ use by expiration date on packet.

**Inactive ingredients**  
corn starch, D&C yellow #10 aluminum lake, lactose anhydrous, magnesium stearate, silicon dioxide

**Questions or comments** Call 1-800-634-7680

**Medique**  
**Medi-Meclizine**  
All Day Relief  
Alivio que Perdura Todo el Día

Pull to Open  
Tire Para Abrir

✓ For Prevention and Treatment of Motion Sickness • **Medicine HC 25 mg**  
Para la Prevención y el Tratamiento del Mareo "Mal del Viaje" • **Fármaco de Acción HC 25 mg**

100 Tablets (50 x 2)

**Información del Medicamento**

**Ingredientes activos (en cada tableta)** Propiedad  
Hidrocloruro de Meclizina 25 mg Antiemético

**Usos**  
para la prevención y tratamiento de la náusea, vómito, laberintismo, o mareo asociados con el "mal del viaje".

**Advertencias**  
No se administra a niños menores de 12 años a menos que lo señale el médico.  
Consulte con el médico antes de tomar este producto si usted tiene:  
■ dificultad para orinar debido al crecimiento de la glándula prostática  
■ glaucoma  
■ problemas respiratorios como asma o bronquitis crónica

Consulte con su médico o farmacéuta antes de administrarle, si usted está tomando sedantes o tranquilizantes.

**Cuando consuma este producto**  
■ puede causar somnolencia  
■ evite el consumo de bebidas alcohólicas  
■ el alcohol, los sedantes y tranquilizantes pueden aumentar los efectos de la somnolencia  
■ tenga cuidado al manejar un vehículo motorizado o al operar maquinaria

Si está usted embarazada o en período de lactancia, consulte la opinión de un médico antes de consumir este producto.

**Información del Medicamento (continuación)**

Mantenga este y todos los medicamentos fuera del alcance de los niños. En caso de ocurrir una sobredosis, comuníquese de inmediato con un médico o llame a un centro de control de envenenamiento (1-800-222-1222).

**Instrucciones**  
■ para prevenir el mareo, debe tomar la primera dosis 1/2 hora o 1 hora antes de comenzar la actividad (viaje)  
■ para tratar el mareo por movimiento, tome a la primera señal de síntomas.

**Adultos y niños (para mayores de 12 años)** 1 ó 2 tabletas una vez al día o como lo señale el médico.

**Niños menores de 12 años:** No se administra a niños menores de 12 años de edad.

**Otra información**  
■ empaquetado con sellado evidente  
■ no use ningún paquete abierto o roto  
■ almacenar a 77°F (25°C), excursiones permitidas entre 59°-86°F (15°-30°C)  
■ evite exponerlo al calor y humedad excesivo  
■ usar antes de la fecha de expiración en el paquete

**Ingredientes inactivos**  
división de maíz, amilante D y C #10 lake de aluminio, lactosa anhidra, estearato de magnesio, dióxido de silicio

¿Tiene alguna pregunta o comentario? Llame al 1-800-634-7680

Retain carton for complete product information

Tamper Evident Unit Dose Packets  
Empaquetado con Sellado Evidente en Paquetes Individuales

Reorder #47933  
Manufactured for  
Medique Products  
Fort Myers, FL 33967  
1-800-634-7680  
www.mediqueproducts.com

# MEDIQUE MEDI-MECLIZINE

meclizine hcl tablet

Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-481
<b>Route of Administration</b>	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
<b>MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)</b>	MECLIZINE HYDROCHLORIDE	25 mg	

Inactive Ingredients			
Ingredient Name	Strength		
<b>STARCH, CORN (UNII: O8232NY3SJ)</b>			
<b>SILICON DIOXIDE (UNII: ETJ7Z6XBU4)</b>			
<b>MAGNESIUM STEARATE (UNII: 70097M6I30)</b>			
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)</b>			
<b>ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)</b>			

Product Characteristics			
<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	44;403
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-481-99	2 in 1 PACKET; Type 0: Not a Combination Product	12/14/2020	
2	NDC:47682-481-15	500 in 1 BOX	12/14/2020	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-481-33	50 in 1 BOX	12/14/2020	
3		2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	12/14/2020	

**Labeler** - Unifirst First Aid Corporation (832947092)

Revised: 1/2024

Unifirst First Aid Corporation