

**EXTRA STRENGTH HEADACHE RELIEF GELTABS- acetaminophen, aspirin,
caffeine tablet**
Strive Pharmaceuticals Inc.

RIGHT REMEDIES EXTRA STRENGTH Headache Relief

Drug Facts

Active ingredients (in each geltab)

Acetaminophen 250 mg
Aspirin 250 mg(NSAID*)
Caffeine 65 mg
*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever
Pain reliever
Pain reliever aid

Uses

Temporarily relieves minor aches and pains due to:
• headache • a cold • arthritis • muscular aches • toothache • premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause skin reactions. Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
• more than 8 geltabs in 24 hours, which is the maximum daily amount
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause

severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness and occasional rapid heartbeat

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor before or pharmacist before use if you are taking

- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed (see overdose warning)**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 geltabs every 6 hours; not more than 8 geltabs in 24 hours
- children under 12 years: ask a doctor

Other information

- **each geltab contains:** sodium 4mg (very low sodium) and potassium 3mg
- store at room temperature between 20-25°C(68-77°F)
- avoid excessive heat, cold and humidity

Inactive ingredients

corn starch, D&C yellow # 10 aluminum lake, edible black ink, ethyl cellulose, FD&C blue #1, FD&C blue #1 aluminum lake, FD&C red # 40 aluminum lake, gelatin, glycerin, hypromellose, isopropyl alcohol, lactose monohydrate, light mineral oil, microcrystalline cellulose, povidone, pregelatinized potato starch, purified water, sorbitol sorbitan solution, stearic acid, talc, titanium dioxide, triacetin

Questions or comments?

1-888-577-8033 Monday-Friday 8am-4pm EST

RIGHT REMEDIES

Compare to the active ingredients of **Excedrin® Extra Strength**

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING FROM BOTTLE

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Excedrin® Extra Strength.


Distributed by:
 Strive Pharmaceuticals Inc.
 East Brunswick, NJ 08816

Made in UAE

Packaged and Quality Assured in the USA

REV.00-062021

Packaging



Headache Relief

ACETAMINOPHEN,
ASPIRIN (NSAID)
and CAFFEINE
Pain reliever,
Pain reliever aid

200
geltabs

Actual Size

AMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS

Drug Facts

Active ingredients (in each geltab)

Acetaminophen 250 mg
Aspirin 250 mg (NSAID*)
Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains due to
■ headache ■ a cold ■ arthritis ■ muscular aches ■ premenstrual and menstrual cramps


Warnings

Reye's syndrome: Children and teenagers who
from chicken pox or flu-like symptoms should not use this product.
When using this product, if changes in behavior with nausea
vomiting occur, consult a doctor because these symptoms could be an
early sign of Reye's syndrome, a rare but serious
Allergy alert: Acetaminophen may cause skin reactions. Symptoms may
include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may
include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock.
Liver warning: This product contains acetaminophen. Severe liver
damage may occur if you take
■ more than 8 geltabs in 24 hours, which is the maximum
daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Stomach bleeding warning: This product contains an NSAID, which
may cause severe stomach bleeding. The chance is higher if you
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs
(aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic
drinks every day while using this product ■ take more or for a longer
time than directed

Caffeine warning:

The recommended dose of this product contains
about as much caffeine as a cup of coffee. Limit the use of
caffeine-containing medications, foods, or beverages while taking this
product because too much caffeine may cause nervousness, irritability,
sleeplessness and occasional rapid heartbeat.

This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Excedrin® Extra Strength.



3 70692 00022 1

PEEL HERE

LOT: LB7069213220
EXP: REV/00-062021

AMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS MISSING FROM BOTTLE

Drug Facts

Active ingredients (in each geltab)

Acetaminophen 250 mg.....Pain reliever
Aspirin 250 mg(NSAID*).....Pain reliever
Caffeine 65 mg.....Pain reliever aid
*nonsteroidal anti-inflammatory drug

Purposes

Uses

Temporarily relieves minor aches and pains due to:
■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache
■ premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering
from chicken pox or flu-like symptoms should not use this product.
When using this product, if changes in behavior with nausea and
vomiting occur, consult a doctor because these symptoms could be an
early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause skin reactions. Symptoms may
include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may
include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock.
Liver warning: This product contains acetaminophen. Severe liver
damage may occur if you take
■ more than 8 geltabs in 24 hours, which is the maximum
daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Stomach bleeding warning: This product contains an NSAID, which
may cause severe stomach bleeding. The chance is higher if you
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs
(aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic
drinks every day while using this product ■ take more or for a longer
time than directed

Caffeine warning: The recommended dose of this product contains
about as much caffeine as a cup of coffee. Limit the use of
caffeine-containing medications, foods, or beverages while taking this
product because too much caffeine may cause nervousness, irritability,
sleeplessness and occasional rapid heartbeat.

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Excedrin® Extra Strength.



Distributed by:
Strive Pharmaceuticals Inc
East Brunswick, NJ 08816

Made in UAE
Packaged and Quality
Assured in the USA

PEEL HERE

LB7069213220
REV.00-062021

LOT: No Varnish
EXP:

Drug Facts (Continued)

Do not use ■ if you have ever had an allergic reaction to
acetaminophen, aspirin or any other pain reliever/fever reducer
■ with any other drug containing acetaminophen (prescription or
nonprescription). If you are not sure whether a drug contains
acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor before or pharmacist before use if you are taking
■ a prescription drug for diabetes, gout, or arthritis
■ any other drug, or are under a doctor's care for any serious
condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It
is especially important not to use aspirin during the last 3 months of
pregnancy unless definitely directed to do so by a doctor because it
may cause problems in the unborn child or complications during
delivery.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a
Poison Control Center right away. Prompt medical attention is critical
for adults as well as children even if you do not notice any signs
or symptoms.

Directions

- **do not use more than directed (see overdose warning)**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 geltabs every 6 hours;
not more than 8 geltabs in 24 hours
- children under 12 years: ask a doctor

Other information

- each geltab contains: sodium 4mg (very low sodium) and potassium 3mg
- store at room temperature between 20-25°C(68-77°F)
- avoid excessive heat, cold and humidity

Inactive ingredients

corn starch, D&C yellow # 10 aluminum lake, edible black ink, ethyl cellulose, FD&C blue #1, FD&C blue #1 aluminum lake, FD&C red # 40 aluminum lake, gelatin, glycerin, hypromellose, isopropyl alcohol, lactose monohydrate, light mineral oil, microcrystalline cellulose, povidone, pregelatinized potato starch, purified water, sorbitol sorbitan solution, stearic acid, talc, titanium dioxide, triacetin

Questions or comments?

1-888-577-8033 Monday-Friday 8am-4pm EST

EXTRA STRENGTH HEADACHE RELIEF GELTABS

acetaminophen, aspirin, caffeine tablet

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:70692-132 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 250 mg |
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | ASPIRIN | 250 mg |
| CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E) | CAFFEINE | 65 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| STARCH, CORN (UNII: O8232NY3SJ) | |
| D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6) | |
| ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| GELATIN, UNSPECIFIED (UNII: 2G86QN327L) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| ISOPROPYL ALCOHOL (UNII: ND2M416302) | |
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X) | |
| LIGHT MINERAL OIL (UNII: N6K5787QVP) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) | |
| STARCH, POTATO (UNII: 8I089SAH3T) | |
| WATER (UNII: 059QF0KO0R) | |
| SORBITOL (UNII: 506T60A25R) | |
| SORBITAN (UNII: 6O92ICV9RU) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| TRIACETIN (UNII: XHX3C3X673) | |

Product Characteristics

| | | | |
|-----------------|--------------------------------|---------------------|----------|
| Color | white, green (White and Green) | Score | no score |
| Shape | ROUND | Size | 13mm |
| Flavor | | Imprint Code | S77 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:70692-132-20 | 200 in 1 BOTTLE; Type 0: Not a Combination Product | 04/05/2023 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M013 | 04/05/2023 | |

Labeler - Strive Pharmaceuticals Inc. (080028013)

Revised: 12/2024

Strive Pharmaceuticals Inc.