

# DICYCLOMINE HYDROCHLORIDE- dicyclomine hydrochloride tablet

## Advagen Pharma Ltd.,

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DICYCLOMINE HYDROCHLORIDE TABLETS safely and effectively. See full prescribing information for DICYCLOMINE HYDROCHLORIDE TABLETS.

**DICYCLOMINE HYDROCHLORIDE tablets, for oral use**  
**Initial U.S. Approval: 1950**

#### RECENT MAJOR CHANGES

Warnings and Precautions, Peripheral and Central Nervous System ( 5.3)

07/2012

#### INDICATIONS AND USAGE

Dicyclomine hydrochloride is an antispasmodic and anticholinergic (antimuscarinic) agent indicated for the treatment of functional bowel/irritable bowel syndrome ( 1)

#### DOSAGE AND ADMINISTRATION

Dosage for dicyclomine hydrochloride tablets must be adjusted to individual patient needs ( 2).  
If a dose is missed, patients should continue the normal dosing schedule. ( 2).

##### Oral in adults ( 2.1):

- Starting dose: 20 mg four times a day. After a week treatment with the starting dose, the dose may be escalated to 40 mg four times a day, unless side effects limit dosage escalation
- Discontinue dicyclomine hydrochloride tablets if efficacy not achieved or side effects require doses less than 80 mg per day after two weeks of treatment

#### DOSAGE FORMS AND STRENGTHS

- Dicyclomine hydrochloride tablets 20 mg ( 3)
- Dicyclomine hydrochloride tablets 40 mg ( 3)

#### CONTRAINDICATIONS

- |                                                               |                                                      |
|---------------------------------------------------------------|------------------------------------------------------|
| • Infants less than 6 months of age ( 4)                      | • Glaucoma ( 4)                                      |
| • Nursing mothers ( 4)                                        | • Obstructive uropathy ( 4)                          |
| • Unstable cardiovascular status in the acute hemorrhage ( 4) | • Obstructive disease of gastrointestinal tract ( 4) |
| • Myasthenia gravis ( 4)                                      | • Severe ulcerative colitis ( 4)                     |
|                                                               | • Reflux esophagitis ( 4)                            |

#### WARNINGS AND PRECAUTIONS

- Cardiovascular conditions: worsening of conditions ( 5.2)
- Peripheral and central nervous system: heat prostration can occur with drug use (fever and heat stroke due to decreased sweating); drug should be discontinued and supportive measures instituted ( 5.3)
- Psychosis and delirium have been reported in patients sensitive to anticholinergic drugs (such as elderly patients and/or in patients with mental illness): signs and symptoms resolve within 12 to 24 hours after discontinuation of dicyclomine hydrochloride tablets ( 5.3)
- Myasthenia Gravis: overdose may lead to muscular weakness and paralysis. Dicyclomine hydrochloride tablets should be given to patients with myasthenia gravis only to reduce adverse muscarinic effects of an anticholinesterase ( 5.4)
- Incomplete intestinal obstruction: diarrhea may be an early symptom especially in patients with ileostomy or colostomy. Treatment with dicyclomine hydrochloride tablets would be inappropriate and possibly fatal ( 5.5)
- Salmonella dysenteric patients: due to risk of toxic megacolon ( 5.6)
- Ulcerative colitis: Dicyclomine hydrochloride tablets should be used with caution in these patients; large

- doses may suppress intestinal motility or aggravate the serious complications of toxic megacolon ( 5.7)
- Prostatic hypertrophy: Dicyclomine hydrochloride tablets should be used with caution in these patients; may lead to urinary retention ( 5.8)
  - Hepatic and renal disease: should be used with caution ( 5.9)
  - Geriatric: use with caution in elderly who may be more susceptible to dicyclomine hydrochloride's adverse events ( 5.10)

#### -----ADVERSE REACTIONS-----

The most serious adverse reactions include cardiovascular and central nervous system symptoms. The most common adverse reactions (> 5% of patients) are dizziness, dry mouth, vision blurred, nausea, somnolence, asthenia and nervousness ( 6)

**To report SUSPECTED ADVERSE REACTIONS, contact Advagen Pharma Ltd, at 866-488-0312 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

#### -----DRUG INTERACTIONS-----

- Antiglaucoma agents: anticholinergics antagonize antiglaucoma agents and may increase intraocular pressure ( 7)
- Anticholinergic agents: may affect the gastrointestinal absorption of various drugs; may also increase certain actions or side effects of other anticholinergic drugs ( 7)
- Antacids: interfere with the absorption of anticholinergic agents ( 7)

#### -----USE IN SPECIFIC POPULATIONS-----

- Pregnancy: use only if clearly needed ( 8.1)
- Pediatric Use: Safety and effectiveness not established ( 8.4)
- Hepatic and renal impairment: caution must be taken with patients with significantly impaired hepatic and renal function ( 8.6)

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 2/2026**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Oral Dosage and Administration in Adults**

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

#### **5.2 Cardiovascular Conditions**

#### **5.3 Peripheral and Central Nervous System**

#### **5.4 Myasthenia Gravis**

#### **5.5 Intestinal Obstruction**

#### **5.6 Toxic Dilatation of Intestinmegacolon**

#### **5.7 Ulcerative Colitis**

#### **5.8 Prostatic Hypertrophy**

#### **5.9 Hepatic and Renal Disease**

#### **5.10 Geriatric Population**

### **6 ADVERSE REACTIONS**

#### **6.1 Clinical Trials Experience**

#### **6.2 Postmarketing Experience**

#### **6.3 Adverse Reactions Reported with Similar Drugs with Anticholinergic/Antispasmodic Action**

## **7 DRUG INTERACTIONS**

### **7.1 Antiglaucoma Agents**

### **7.2 Other Drugs with Anticholinergic Activity**

### **7.3 Other Gastrointestinal Motility Drugs**

### **7.4 Effect of Antacids**

### **7.5 Effect on Absorption of Other Drugs**

### **7.6 Effect on Gastric Acid Secretion**

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

### **8.3 Nursing Mothers**

### **8.4 Pediatric Use**

### **8.5 Geriatric Use**

### **8.6 Renal Impairment**

### **8.7 Hepatic Impairment**

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

### **12.2 Pharmacodynamics**

### **12.3 Pharmacokinetics**

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

## **14 CLINICAL STUDIES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

### **17.2 Use in Infants**

### **17.3 Use in Nursing Mothers**

### **17.4 Peripheral and Central Nervous System**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Dicyclomine hydrochloride tablets are indicated for the treatment of patients with functional bowel/irritable bowel syndrome.

### **2 DOSAGE AND ADMINISTRATION**

Dosage must be adjusted to individual patient needs.

#### **2.1 Oral Dosage and Administration in Adults**

The recommended initial dose is 20 mg four times a day.

After one week treatment with the initial dose, the dose may be increased to 40 mg four times a day unless side effects limit dosage escalation.

If efficacy is not achieved within 2 weeks or side effects require doses below 80 mg per day, the drug should be discontinued.

Documented safety data are not available for doses above 80 mg daily for periods longer than 2 weeks.

### **3 DOSAGE FORMS AND STRENGTHS**

Dicyclomine hydrochloride tablets, USP 20 mg: Light blue to blue, speckled, round, unscored, flat-faced, beveled-edge tablets, debossed '^' and 'A2' on one side and plain on the other side.

Dicyclomine hydrochloride tablets, USP 40 mg: Light blue to blue, speckled, oval shaped, concave tablet with scoreline on both sides, debossed on one side with '^' and 'H6' on either side of scoreline.

### **4 CONTRAINDICATIONS**

Dicyclomine hydrochloride is contraindicated in infants less than 6 months of age [*see Use in Specific Populations ( 8.4)*] , nursing mothers [*see Use in Specific Populations ( 8.3)*], and in patients with:

- unstable cardiovascular status in acute hemorrhage
- myasthenia gravis [*see Warnings and Precautions ( 5.4)*]
- glaucoma [*see Adverse Reactions ( 6.3) and Drug Interactions ( 7.1)*]
- obstructive uropathy [*see Warnings and Precautions ( 5.8)*]
- obstructive disease of the gastrointestinal tract [*see Warnings and Precautions ( 5.5)*]
- severe ulcerative colitis [*see Warnings and Precautions ( 5.7)*]
- reflux esophagitis

### **5 WARNINGS AND PRECAUTIONS**

#### **5.2 Cardiovascular Conditions**

Dicyclomine hydrochloride needs to be used with caution in conditions characterized by tachyarrhythmia such as thyrotoxicosis, congestive heart failure and in cardiac surgery, where they may further accelerate the heart rate. Investigate any tachycardia before administration of dicyclomine hydrochloride. Care is required in patients with coronary heart disease, as ischemia and infarction may be worsened, and in patients with hypertension [*see Adverse Reactions ( 6.3)*] .

#### **5.3 Peripheral and Central Nervous System**

The peripheral effects of dicyclomine hydrochloride are a consequence of their inhibitory effect on muscarinic receptors of the autonomic nervous system. They include dryness of the mouth with difficulty in swallowing and talking, thirst, reduced bronchial secretions, dilatation of the pupils (mydriasis) with loss of accommodation (cycloplegia) and photophobia, flushing and dryness of the skin, transient bradycardia followed by tachycardia, with palpitations and arrhythmias, and difficulty in micturition, as well as reduction in the tone and motility of the gastrointestinal tract leading to constipation [*see*

*Adverse Reactions ( 6)] .*

In the presence of high environmental temperature heat prostration can occur with drug use (fever and heat stroke due to decreased sweating). It should also be used cautiously in patients with fever. If symptoms occur, the drug should be discontinued and supportive measures instituted. Because of the inhibitory effect on muscarinic receptors within the autonomic nervous system, caution should be taken in patients with autonomic neuropathy.

Central nervous system (CNS) signs and symptoms include confusional state, disorientation, amnesia, hallucinations, dysarthria, ataxia, coma, euphoria, fatigue, insomnia, agitation and mannerisms, and inappropriate affect.

Psychosis and delirium have been reported in sensitive individuals (such as elderly patients and/or in patients with mental illness) given anticholinergic drugs. These CNS signs and symptoms usually resolve within 12 to 24 hours after discontinuation of the drug.

Dicyclomine hydrochloride may produce drowsiness, dizziness or blurred vision. The patient should be warned not to engage in activities requiring mental alertness, such as operating a motor vehicle or other machinery or performing hazardous work while taking dicyclomine hydrochloride.

#### **5.4 Myasthenia Gravis**

With overdosage, a curare-like action may occur (i.e., neuromuscular blockade leading to muscular weakness and possible paralysis). It should not be given to patients with myasthenia gravis except to reduce adverse muscarinic effects of an anticholinesterase [*see Contraindications ( 4)] .*

#### **5.5 Intestinal Obstruction**

Diarrhea may be an early symptom of incomplete intestinal obstruction, especially in patients with ileostomy or colostomy. In this instance, treatment with this drug would be inappropriate and possibly harmful [*see Contraindications ( 4)] .*

Rarely development of Ogilvie's syndrome (colonic pseudo-obstruction) has been reported. Ogilvie's syndrome is a clinical disorder with signs, symptoms, and radiographic appearance of an acute large bowel obstruction but with no evidence of distal colonic obstruction.

#### **5.6 Toxic Dilatation of Intestinmegacolon**

Toxic dilatation of intestine and intestinal perforation is possible when anticholinergic agents are administered in patients with Salmonella dysentery.

#### **5.7 Ulcerative Colitis**

Caution should be taken in patients with ulcerative colitis. Large doses may suppress intestinal motility to the point of producing a paralytic ileus and the use of this drug may precipitate or aggravate the serious complication of toxic megacolon [*see Adverse Reactions ( 6.3)] .* Dicyclomine hydrochloride tablets are contraindicated in patients with severe ulcerative colitis [*see Contraindications ( 4)] .*

#### **5.8 Prostatic Hypertrophy**

Dicyclomine hydrochloride should be used with caution in patients with known or

suspected prostatic enlargement, in whom prostatic enlargement may lead to urinary retention [see *Adverse Reactions* ( 6.3)] .

## 5.9 Hepatic and Renal Disease

Dicyclomine hydrochloride should be used with caution in patients with known hepatic and renal impairment.

## 5.10 Geriatric Population

Dicyclomine hydrochloride should be used with caution in elderly who may be more susceptible to its adverse effects.

# 6 ADVERSE REACTIONS

The pattern of adverse effects seen with dicyclomine is mostly related to its pharmacological actions at muscarinic receptors [see *Clinical Pharmacology* ( 12)] . They are a consequence of the inhibitory effect on muscarinic receptors within the autonomic nervous system. These effects are dose-related and are usually reversible when treatment is discontinued.

The most serious adverse reactions reported with dicyclomine hydrochloride include cardiovascular and central nervous system symptoms [see *Warnings and Precautions* ( 5.2, 5.3)] .

## 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice. The data described below reflect exposure in controlled clinical trials involving over 100 patients treated for functional bowel/irritable bowel syndrome with dicyclomine hydrochloride at initial doses of 160 mg daily (40 mg four times a day) In these trials most of the side effects were typically anticholinergic in nature and were reported by 61% of the patients. Table 1 presents adverse reactions ( *MedDRA* 13.0 preferred terms) by decreasing order of frequency in a side-by-side comparison with placebo.

**Table 1: Adverse reactions experienced in controlled clinical trials with decreasing order of frequency**

| <b>MedDRA Preferred Term</b> | <b>Dicyclomine Hydrochloride (40 mg four times a day) %</b> | <b>Placebo %</b> |
|------------------------------|-------------------------------------------------------------|------------------|
| Dry Mouth                    | 33                                                          | 5                |
| Dizziness                    | 40                                                          | 5                |
| Vision blurred               | 27                                                          | 2                |
| Nausea                       | 14                                                          | 6                |

|             |   |   |
|-------------|---|---|
| Somnolence  | 9 | 1 |
| Asthenia    | 7 | 1 |
| Nervousness | 6 | 2 |

Nine percent (9%) of patients were discontinued from dicyclomine hydrochloride because of one or more of these side effects (compared with 2% in the placebo group). In 41% of the patients with side effects, side effects disappeared or were tolerated at the 160 mg daily dose without reduction. A dose reduction from 160 mg daily to an average daily dose of 90 mg was required in 46% of the patients with side effects who then continued to experience a favorable clinical response; their side effects either disappeared or were tolerated.

## 6.2 Postmarketing Experience

The following adverse reactions, presented by system organ class in alphabetical order, have been identified during post approval use of dicyclomine hydrochloride. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- *Cardiac disorders*: palpitations, tachyarrhythmias
- *Eye disorders*: cycloplegia, mydriasis, vision blurred
- *Gastrointestinal disorders*: abdominal distension, abdominal pain, constipation, dry mouth, dyspepsia, nausea, vomiting
- *General disorders and administration site conditions*: fatigue, malaise
- *Immune System Disorders*: drug hypersensitivity including face edema, angioedema, anaphylactic shock
- *Nervous system disorders*: dizziness, headache, somnolence, syncope
- *Psychiatric disorders*: As with the other anti-cholinergic drugs, cases of delirium or symptoms of delirium such as amnesia (or transient global amnesia), agitation, confusional state, delusion, disorientation, hallucination (including visual hallucination) as well as mania, mood altered and pseudodementia, have been reported with the use of dicyclomine. Nervousness and insomnia have also been reported.
- *Reproductive system and breast disorders*: suppressed lactation
- *Respiratory, thoracic and mediastinal disorders*: dyspnoea, nasal congestion
- *Skin and subcutaneous tissue disorder*: dermatitis allergic, erythema, rash

## 6.3 Adverse Reactions Reported with Similar Drugs with Anticholinergic/Antispasmodic Action

Gastrointestinal: anorexia

Central Nervous System: tingling, numbness, dyskinesia, speech disturbance, insomnia

Peripheral Nervous System: With overdosage, a curare-like action may occur (i.e., neuromuscular blockade leading to muscular weakness and possible paralysis)

Ophthalmologic: diplopia, increased ocular tension

Dermatologic/Allergic: urticaria, itching, and other dermal manifestations

Genitourinary: urinary hesitancy, urinary retention in patients with prostatic hypertrophy

Cardiovascular: hypertension

Respiratory: apnea

Other: decreased sweating, sneezing, throat congestion, impotence.

## **7 DRUG INTERACTIONS**

### **7.1 Antiglaucoma Agents**

Anticholinergics antagonize the effects of antiglaucoma agents. Anticholinergic drugs in the presence of increased intraocular pressure may be hazardous when taken concurrently with agents such as corticosteroids. Use of dicyclomine hydrochloride tablets in patients with glaucoma is not recommended [see *Contraindications ( 4)*].

### **7.2 Other Drugs with Anticholinergic Activity**

The following agents may increase certain actions or side effects of anticholinergic drugs including dicyclomine hydrochloride: amantadine, antiarrhythmic agents of Class I (e.g., quinidine), antihistamines, antipsychotic agents (e.g., phenothiazines), benzodiazepines, MAO inhibitors, narcotic analgesics (e.g., meperidine), nitrates and nitrites, sympathomimetic agents, tricyclic antidepressants, and other drugs having anticholinergic activity.

### **7.3 Other Gastrointestinal Motility Drugs**

Interaction with other gastrointestinal motility drugs may antagonize the effects of drugs that alter gastrointestinal motility, such as metoclopramide.

### **7.4 Effect of Antacids**

Because antacids may interfere with the absorption of anticholinergic agents including dicyclomine hydrochloride, simultaneous use of these drugs should be avoided.

### **7.5 Effect on Absorption of Other Drugs**

Anticholinergic agents may affect gastrointestinal absorption of various drugs by affecting on gastrointestinal motility, such as slowly dissolving dosage forms of digoxin; increased serum digoxin concentration may result.

### **7.6 Effect on Gastric Acid Secretion**

The inhibiting effects of anticholinergic drugs on gastric hydrochloric acid secretion are antagonized by agents used to treat achlorhydria and those used to test gastric secretion.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

Pregnancy Category B

Adequate and well-controlled studies have not been conducted with dicyclomine hydrochloride in pregnant women at the recommended doses of 80 to 160 mg/day. However, epidemiologic studies did not show an increased risk of structural malformations among babies born to women who took products containing dicyclomine hydrochloride at doses up to 40 mg/day during the first trimester of pregnancy. Reproduction studies have been performed in rats and rabbits at doses up

to 33 times the maximum recommended human dose based on 160 mg/day (3 mg/kg) and have revealed no evidence of harm to the fetus due to dicyclomine. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

### **8.3 Nursing Mothers**

Dicyclomine hydrochloride is contraindicated in women who are breastfeeding. Dicyclomine hydrochloride is excreted in human milk. Because of the potential for serious adverse reactions in breast-fed infants from dicyclomine hydrochloride tablets, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother [*see Use in Specific Populations ( 8.4)*].

### **8.4 Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

Dicyclomine hydrochloride is contraindicated in infants less than 6 months of age [*see Contraindications ( 4)*]. There are published cases reporting that the administration of dicyclomine hydrochloride to infants has been followed by serious respiratory symptoms (dyspnea, shortness of breath, breathlessness, respiratory collapse, apnea and asphyxia), seizures, syncope, pulse rate fluctuations, muscular hypotonia, and coma, and death, however; no causal relationship has been established.

### **8.5 Geriatric Use**

Clinical studies of dicyclomine hydrochloride tablets did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range in adults, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

### **8.6 Renal Impairment**

Effects of renal impairment on PK, safety and efficacy of dicyclomine hydrochloride have not been studied. Dicyclomine hydrochloride drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Dicyclomine hydrochloride tablets should be administered with caution in patients with renal impairment.

### **8.7 Hepatic Impairment**

Effects of renal impairment on PK, safety and efficacy of dicyclomine hydrochloride has not been studied. Dicyclomine hydrochloride tablets should be administered with caution in patients with hepatic impairment.

## 10 OVERDOSAGE

In case of an overdose, patients should contact a physician, poison control center (1-800-222-1222), or emergency room.

The signs and symptoms of overdose include: headache; nausea; vomiting; blurred vision; dilated pupils; hot, dry skin; dizziness; dryness of the mouth; difficulty in swallowing; and CNS stimulation including convulsion. A curare-like action may occur (i.e., neuromuscular blockade leading to muscular weakness and possible paralysis).

One reported event included a 37-year-old who reported numbness on the left side, cold fingertips, blurred vision, abdominal and flank pain, decreased appetite, dry mouth, and nervousness following ingestion of 320 mg daily (four 20 mg tablets four times daily). These events resolved after discontinuing the dicyclomine.

The acute oral LD50 of the drug is 625 mg/kg in mice.

The amount of drug in a single dose that is ordinarily associated with symptoms of overdose or that is likely to be life-threatening, has not been defined. The maximum human oral dose recorded was 600 mg by mouth in a 10-month-old child and approximately 1500 mg in an adult, each of whom survived. In three of the infants who died following administration of dicyclomine hydrochloride [see *Warnings and Precautions* ( 5.1)] , the blood concentrations of drug were 200, 220, and 505 ng/mL.


It is not known if dicyclomine hydrochloride is dialyzable.

Treatment should consist of gastric lavage, emetics, and activated charcoal. Sedatives (e.g., short-acting barbiturates, benzodiazepines) may be used for management of overt signs of excitement. If indicated, an appropriate parenteral cholinergic agent may be used as an antidote.

## 11 DESCRIPTION

Dicyclomine hydrochloride is an antispasmodic and anticholinergic (antimuscarinic) agent available as dicyclomine hydrochloride tablets, USP for oral use containing 20 mg and 40 mg dicyclomine hydrochloride USP. It also contain inactive ingredients: lactose monohydrate, hypromellose, pregelatinized maize starch, FD & C Blue 1 Alu.lake, colloidal silicon dioxide and magnesium stearate.

Dicyclomine hydrochloride is [bicyclohexyl]-1- carboxylic acid, 2-(diethylamino) ethyl ester, hydrochloride, with a molecular formula of  $C_{19}H_{35}NO_2 \cdot HCl$  and the following structural formula:



Molecular weight: 345.95

Dicyclomine hydrochloride occurs as a fine, white, crystalline, practically odorless powder with a bitter taste. It is soluble in water, freely soluble in alcohol and chloroform, and very slightly soluble in ether.

FDA approved dissolution test specifications differ from USP.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Dicyclomine relieves smooth muscle spasm of the gastrointestinal tract. Animal studies indicate that this action is achieved via a dual mechanism:

- a specific anticholinergic effect (antimuscarinic) at the acetylcholine-receptor sites with approximately 1/8 the milligram potency of atropine ( *in vitro*, guinea pig ileum); and
- a direct effect upon smooth muscle (musculotropic) as evidenced by dicyclomine's antagonism of bradykinin- and histamine-induced spasms of the isolated guinea pig ileum.

Atropine did not affect responses to these two agonists. *In vivo* studies in cats and dogs showed dicyclomine to be equally potent against acetylcholine (ACh)- or barium chloride (BaCl<sub>2</sub>)-induced intestinal spasm while atropine was at least 200 times more potent against effects of ACh than BaCl<sub>2</sub>. Tests for mydriatic effects in mice showed that dicyclomine was approximately 1/500 as potent as atropine; antisialagogue tests in rabbits showed dicyclomine to be 1/300 as potent as atropine.

### **12.2 Pharmacodynamics**

Dicyclomine hydrochloride can inhibit the secretion of saliva and sweat, decrease gastrointestinal secretions and motility, cause drowsiness, dilate the pupils, increase heart rate, and depress motor function

### **12.3 Pharmacokinetics**

#### Absorption and Distribution

In man, dicyclomine is rapidly absorbed after oral administration, reaching peak values within 60-90 minutes. Mean volume of distribution for a 20 mg oral dose is approximately 3.65 L/kg suggesting extensive distribution in tissues.

#### Elimination

The metabolism of dicyclomine was not studied. The principal route of excretion is via the urine (79.5% of the dose). Excretion also occurs in the feces, but to a lesser extent (8.4%). Mean half-life of plasma elimination in one study was determined to be approximately 1.8 hours when plasma concentrations were measured for 9 hours after a single dose. In subsequent studies, plasma concentrations were followed for up to 24 hours after a single dose, showing a secondary phase of elimination with a somewhat longer half-life.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term animal studies have not been conducted to evaluate the carcinogenic potential of dicyclomine. In studies in rats at doses of up to 100 mg/kg/day, dicyclomine produced no deleterious effects on breeding, conception, or parturition.

## **14 CLINICAL STUDIES**

In controlled clinical trials involving over 100 patients who received drug, 82% of patients treated for functional bowel/irritable bowel syndrome with dicyclomine hydrochloride at initial doses of 160 mg daily (40 mg four times daily) demonstrated a favorable clinical response compared with 55% treated with placebo ( $p < 0.05$ ).

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

Dicyclomine hydrochloride tablets USP, 20 mg and 40 mg are available as follows,

20 mg are available as light blue to blue, speckled, round, unscored, flat-faced, beveled-edge tablets, debossed '^' and 'A2' on one side and plain on the other side.

40 mg are available as light blue to blue, speckled, oval shaped, concave tablet with scoreline on both sides, debossed on one side with '^' and 'H6' on either side of scoreline.

They are supplied as follows:

20 mg:

Bottles of 30, NDC 72888-119-30

Bottles of 100, NDC 72888-119-01

Bottles of 500, NDC 72888-119-05

Bottles of 1,000, NDC 72888-119-00

40 mg:

Bottles of 30, NDC 72888-198-30

Bottles of 100, NDC 72888-198-01

Bottles of 120, NDC 72888-198-12

Bottles of 500, NDC 72888-198-05

Bottles of 1,000, NDC 72888-198-00

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light.

## **17 PATIENT COUNSELING INFORMATION**

### **17.2 Use in Infants**

Inform parents and caregivers not to administer dicyclomine hydrochloride tablets in

infants less than 6 months of age [see Use in Specific Populations ( 8.4)].

### 17.3 Use in Nursing Mothers

Advise lactating women that dicyclomine hydrochloride tablets should not be used while breastfeeding their infants [see Use in Specific Populations ( 8.3, 8.4)] .

### 17.4 Peripheral and Central Nervous System

In the presence of a high environmental temperature, heat prostration can occur with dicyclomine hydrochloride tablets use (fever and heat stroke due to decreased sweating). If symptoms occur, the drug should be discontinued and a physician contacted. Dicyclomine hydrochloride tablets may produce drowsiness or blurred vision. The patient should be warned not to engage in activities requiring mental alertness, such as operating a motor vehicle or other machinery or to perform hazardous work while taking dicyclomine hydrochloride tablets [see Warnings and Precautions ( 5.3)].

#### Distributed By:

Advagen Pharma Ltd.,  
East Windsor, NJ 08520, USA

#### Manufactured by:

Rubicon Research Ltd.,  
Thane 421506, India.

Rev.: 03, 02/26

### PRINCIPAL DISPLAY PANEL

#### Dicyclomine Hydrochloride Tablets, USP 20 mg - NDC 72888-119-30 - 30 Tablets Label

|                                                                                                                          |                                                                                                                                                                             |                                                                                             |                                                                             |                                                                                       |       |
|--------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|---------------------------------------------------------------------------------------|-------|
| <br>4<br>7 2 8 8 8 1 1 9 3 0 1 1<br>N | Each tablet contains:<br>20 mg of Dicyclomine Hydrochloride, USP.                                                                                                           | NDC 72888-119-30                                                                            | Distributed by: <b>Advagen Pharma Ltd.</b> ,<br>East Windsor, NJ 08520, USA |  |       |
|                                                                                                                          | <b>Dosage and Administration:</b><br>See package insert for dosage information.                                                                                             | <b>Dicyclomine Hydrochloride Tablets, USP</b>                                               | Manufactured by: <b>Rubicon Research Ltd.</b> ,<br>Thane 421506, India.     |                                                                                       |       |
|                                                                                                                          | Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light.                       |                                                                                             | Mfg Lic No. : KD-682                                                        | Rev.04, 11/24                                                                         | XXXXX |
|                                                                                                                          | <b>Pharmacist:</b> Dispense in a well-closed container with child-resistant closure.                                                                                        | <b>20 mg</b>                                                                                | <b>Overprinting Area</b><br>43 mm X 15 mm<br>↑                              |                                                                                       |       |
|                                                                                                                          | <b>WARNING:</b> Keep out of reach of children.                                                                                                                              |  ADVAGEN |                                                                             |                                                                                       |       |
|                                                                                                                          | Package Insert available at <a href="https://www.advagenpharma.com/pil/dicyclominehydrochloridetabets">https://www.advagenpharma.com/pil/dicyclominehydrochloridetabets</a> | 30 Tablets                                                                                  | Rx Only                                                                     |                                                                                       |       |

#### Dicyclomine Hydrochloride Tablets, USP 20 mg - NDC 72888-119-01 - 100 Tablets Label



Each tablet contains:  
20 mg of Dicyclomine Hydrochloride, USP.

**Dosage and Administration:**  
See package insert for dosage information.

**Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].** Protect from light.

**Pharmacist:** Dispense in a well-closed container with child-resistant closure.

**WARNING:** Keep out of reach of children.

**Package Insert available at** <https://www.advagenpharma.com/pil/dicyclominehydrochloridetabets>

NDC 72888-119-01

## Dicyclomine Hydrochloride Tablets, USP

**20 mg**



100 Tablets Rx Only

Distributed by: **Advagen Pharma Ltd.**, East Windsor, NJ 08520, USA

Manufactured by: **Rubicon Research Ltd.**, Thane 421506, India.

Mfg Lic No. : KD-682      Rev.04, 11/24      21934

**Overprinting Area**

43 mm X 15 mm

### Dicyclomine Hydrochloride Tablets, USP 20 mg - NDC 72888-119-05 - 500 Tablets Label



Each tablet contains: 20 mg of Dicyclomine Hydrochloride, USP.

**Dosage and Administration:**  
See package insert for dosage information.

**Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].** Protect from light.

**Pharmacist:** Dispense in a well-closed container with child-resistant closure.

**WARNING:** Keep out of reach of children.

**Package Insert available at** <https://www.advagenpharma.com/pil/dicyclominehydrochloridetabets>

NDC 72888-119-05

## Dicyclomine Hydrochloride Tablets, USP

**20 mg**



500 Tablets Rx Only

Distributed by: **Advagen Pharma Ltd.**, East Windsor, NJ 08520, USA


Manufactured by: **Rubicon Research Ltd.**, Thane 421506, India.

Mfg Lic No. : KD-682      Rev.04, 11/24      XXXXX

**Overprinting Area**

45 mm X 20 mm

### Dicyclomine Hydrochloride Tablets, USP 20 mg - NDC 72888-119-00 - 1000 Tablets Label



Each tablet contains: 20 mg of Dicyclomine Hydrochloride, USP.

**Dosage and Administration:**  
See package insert for dosage information.

**Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].** Protect from light.

**Pharmacist:** Dispense in a well-closed container with child-resistant closure.

**WARNING:** Keep out of reach of children.

**Package Insert available at** <https://www.advagenpharma.com/pil/dicyclominehydrochloridetabets>

NDC 72888-119-00

## Dicyclomine Hydrochloride Tablets, USP

**20 mg**



1000 Tablets Rx Only

Distributed by: **Advagen Pharma Ltd.**, East Windsor, NJ 08520, USA

Manufactured by: **Rubicon Research Ltd.**, Thane 421506, India.

Mfg Lic No. : KD-682      Rev.04, 11/24      21935

**Overprinting Area**

45 mm X 20 mm

## Dicyclomine Hydrochloride Tablets, USP 40 mg - NDC 72888-198-30 - 30 Tablets Label



N 3 72888 19830 9

Each tablet contains:  
40 mg of Dicyclomine Hydrochloride, USP.

**Dosage and Administration:**  
See package insert for dosage information.

**Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].** Protect from light.

**WARNING:** Keep out of reach of children.

**Pharmacist:** Dispense in a well-closed container with child-resistant closure.

**Package Insert available at** <https://www.advagenpharma.com/pil/dicyclominehydrochloridetables>

NDC 72888-198-30

**Dicyclomine Hydrochloride Tablets, USP**

**40 mg**

30 Tablets Rx Only

 ADVAGEN

Distributed by: **Advagen Pharma Ltd.**,  
East Windsor, NJ 08520, USA

Manufactured by: **Rubicon Research Ltd.**,  
Thane 421506, India.

Mfg Lic No. : KD-682      Rev.02, 03/25      21812



**Overprinting Area**  
45 mm X 15 mm



## Dicyclomine Hydrochloride Tablets, USP 40 mg - NDC 72888-198-01 - 100 Tablets Label



N 3 72888 19801 9

Each tablet contains: 40 mg of  
Dicyclomine Hydrochloride, USP.

**Dosage and Administration:**  
See package insert for dosage information.

**Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].** Protect from light.

**WARNING:** Keep out of reach of children.

**Pharmacist:** Dispense in a well-closed container with child-resistant closure.

**Package Insert available at** <https://www.advagenpharma.com/pil/dicyclominehydrochloridetables>

NDC 72888-198-01

**Dicyclomine Hydrochloride Tablets, USP**

**40 mg**

100 Tablets Rx Only

 ADVAGEN

Distributed by: **Advagen Pharma Ltd.**,  
East Windsor, NJ 08520, USA

Manufactured by: **Rubicon Research Ltd.**,  
Thane 421506, India.

Mfg Lic No. : KD-682

Rev.02, 03/25

21815



**Overprinting Area**  
43 mm X 20 mm



## Dicyclomine Hydrochloride Tablets, USP 40 mg - NDC 72888-198-12 - 120 Tablets Label



Each tablet contains: 40 mg of Dicyclomine Hydrochloride, USP.

**Dosage and Administration:**  
See package insert for dosage information.

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light.

**WARNING:** Keep out of reach of children.

**Pharmacist:** Dispense in a well-closed container with child-resistant closure.

Package Insert available at <https://www.advagenpharma.com/pil/dicyclominehydrochloridetables>

NDC 72888-198-12

**Dicyclomine Hydrochloride Tablets, USP**

**40 mg**

120 Tablets Rx Only

ADVAGEN

Distributed by: **Advagen Pharma Ltd.**, East Windsor, NJ 08520, USA

Manufactured by: **Rubicon Research Ltd.**, Thane 421506, India.

Mfg Lic No. : KD-682

Rev.01, 03/25

22391

**Overprinting Area**

43 mm X 20 mm

**Dicyclomine Hydrochloride Tablets,USP 40 mg - NDC 72888-198-05 - 500 Tablets Label**



Each tablet contains: 40 mg of Dicyclomine Hydrochloride, USP.

**Dosage and Administration:**  
See package insert for dosage information.

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light.

**WARNING:** Keep out of reach of children.

**Pharmacist:** Dispense in a well-closed container with child-resistant closure.

Package Insert available at <https://www.advagenpharma.com/pil/dicyclominehydrochloridetables>

NDC 72888-198-05

**Dicyclomine Hydrochloride Tablets, USP**

**40 mg**

500 Tablets Rx Only

ADVAGEN

Distributed by: **Advagen Pharma Ltd.**, East Windsor, NJ 08520, USA

Manufactured by: **Rubicon Research Ltd.**, Thane 421506, India.

Mfg Lic No. : KD-682

Rev.02, 03/25

21814

**Overprinting Area**

47 mm X 20 mm

**Dicyclomine Hydrochloride Tablets,USP 40 mg - NDC 72888-198-00 - 1000 Tablets Label**



Each tablet contains: 40 mg of  
Dicyclomine Hydrochloride, USP.

**Dosage and Administration:**  
See package insert for dosage information.

Store at 20° to 25°C (68° to 77°F);  
excursions permitted between  
15° to 30°C (59° to 86°F) [see USP  
Controlled Room Temperature].  
Protect from light.

**WARNING:** Keep out of reach of children.

**Pharmacist:** Dispense in a well-closed  
container with child-resistant closure.

**Package Insert available at**  
[https://www.advagenpharma.com  
/pil/dicyclominehydrochloridetables](https://www.advagenpharma.com/pil/dicyclominehydrochloridetables)

NDC 72888-198-00

# Dicyclomine Hydrochloride Tablets, USP

**40 mg**

1000 Tablets

Rx Only



Distributed by: **Advagen Pharma Ltd.**,  
East Windsor, NJ 08520, USA

Manufactured by: **Rubicon Research Ltd.**,  
Thane 421506, India.

Mfg Lic No. : KD-682

Rev.02, 03/25

21813



**Overprinting  
Area**  
47 mm X 20 mm



## DICYCLOMINE HYDROCHLORIDE

dicyclomine hydrochloride tablet

### Product Information

|                                |                         |                           |               |
|--------------------------------|-------------------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN PRESCRIPTION DRUG | <b>Item Code (Source)</b> | NDC:72888-198 |
| <b>Route of Administration</b> | ORAL                    |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name                                                                      | Basis of Strength         | Strength |
|--------------------------------------------------------------------------------------|---------------------------|----------|
| <b>DICYCLOMINE HYDROCHLORIDE</b> (UNII: CQ903KQA31) (DICYCLOMINE - UNII: 4KV4X8IF6V) | DICYCLOMINE HYDROCHLORIDE | 40 mg    |

### Inactive Ingredients

| Ingredient Name                                             | Strength |
|-------------------------------------------------------------|----------|
| <b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)               |          |
| <b>HYPROMELLOSE 2910 (15 MPA.S)</b> (UNII: 365FW2JZ0W)      |          |
| <b>STARCH, CORN</b> (UNII: O8232NY3SJ)                      |          |
| <b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM) |          |
| <b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)                   |          |
| <b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)                |          |

### Product Characteristics

|                 |                           |                     |                |
|-----------------|---------------------------|---------------------|----------------|
| <b>Color</b>    | blue (light blue to blue) | <b>Score</b>        | no score       |
| <b>Shape</b>    | OVAL                      | <b>Size</b>         | 13mm           |
| <b>Flavor</b>   |                           | <b>Imprint Code</b> | UpArrowhead;H6 |
| <b>Contains</b> |                           |                     |                |

### Packaging

| # | Item Code        | Package Description                                 | Marketing Start Date | Marketing End Date |
|---|------------------|-----------------------------------------------------|----------------------|--------------------|
| 1 | NDC:72888-198-30 | 30 in 1 BOTTLE; Type 0: Not a Combination Product   | 02/17/2026           |                    |
| 2 | NDC:72888-198-01 | 100 in 1 BOTTLE; Type 0: Not a Combination Product  | 03/25/2025           |                    |
| 3 | NDC:72888-198-12 | 120 in 1 BOTTLE; Type 0: Not a Combination Product  | 03/25/2025           |                    |
| 4 | NDC:72888-198-05 | 500 in 1 BOTTLE; Type 0: Not a Combination Product  | 03/25/2025           |                    |
| 5 | NDC:72888-198-00 | 1000 in 1 BOTTLE; Type 0: Not a Combination Product | 03/25/2025           |                    |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|------------------------------------------|----------------------|--------------------|
| ANDA               | ANDA216736                               | 03/25/2025           |                    |

DICYCLOMINE HYDROCHLORIDE

dicyclomine hydrochloride tablet

Product Information

|                         |                         |                    |               |
|-------------------------|-------------------------|--------------------|---------------|
| Product Type            | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:72888-119 |
| Route of Administration | ORAL                    |                    |               |

Active Ingredient/Active Moiety

| Ingredient Name                                                              | Basis of Strength         | Strength |
|------------------------------------------------------------------------------|---------------------------|----------|
| DICYCLOMINE HYDROCHLORIDE (UNII: CQ903KQA31) (DICYCLOMINE - UNII:4KV4X8IF6V) | DICYCLOMINE HYDROCHLORIDE | 20 mg    |

Inactive Ingredients

| Ingredient Name                                  | Strength |
|--------------------------------------------------|----------|
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)           |          |
| HYPROMELLOSE 2910 (15 MPA.S) (UNII: 36SFW2JZ0W)  |          |
| STARCH, CORN (UNII: O8232NY3SJ)                  |          |
| FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM) |          |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)               |          |
| MAGNESIUM STEARATE (UNII: 70097M6I3O)            |          |

Product Characteristics

|          |                           |              |                |
|----------|---------------------------|--------------|----------------|
| Color    | blue (light blue to blue) | Score        | no score       |
| Shape    | ROUND                     | Size         | 8mm            |
| Flavor   |                           | Imprint Code | UpArrowhead;A2 |
| Contains |                           |              |                |

## Packaging

| # | Item Code        | Package Description                                 | Marketing Start Date | Marketing End Date |
|---|------------------|-----------------------------------------------------|----------------------|--------------------|
| 1 | NDC:72888-119-30 | 30 in 1 BOTTLE; Type 0: Not a Combination Product   | 12/14/2022           |                    |
| 2 | NDC:72888-119-01 | 100 in 1 BOTTLE; Type 0: Not a Combination Product  | 12/14/2022           |                    |
| 3 | NDC:72888-119-05 | 500 in 1 BOTTLE; Type 0: Not a Combination Product  | 12/14/2022           |                    |
| 4 | NDC:72888-119-00 | 1000 in 1 BOTTLE; Type 0: Not a Combination Product | 12/14/2022           |                    |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|------------------------------------------|----------------------|--------------------|
| ANDA               | ANDA216736                               | 12/14/2022           |                    |

**Labeler** - Advagen Pharma Ltd., (051627256)

## Establishment

| Name                     | Address | ID/FEI    | Business Operations                                                                                                           |
|--------------------------|---------|-----------|-------------------------------------------------------------------------------------------------------------------------------|
| Rubicon Research Limited |         | 677604197 | manufacture(72888-119, 72888-198) , analysis(72888-119, 72888-198) , label(72888-119, 72888-198) , pack(72888-119, 72888-198) |

Revised: 2/2026

Advagen Pharma Ltd.,