

ACETAMINOPHEN PAIN RELIEVER FEVER REDUCER-
acetaminophen suppository
Padagis Israel Pharmaceuticals Ltd

Perrigo Acetaminophen Suppositories USP, 650 mg Drug Facts

Active ingredient (in each rectal suppository)

Acetaminophen 650 mg

Purposes

Pain reliever/fever reducer

Uses

temporarily

- reduces fever
- relieves minor aches, pains, and headache

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if

- an adult or child 12 years and older takes more than 6 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- an adult takes 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

For rectal use only

Do not use

- in children under 12 years
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

Ask a doctor before use if

- you have liver disease
- you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- fever lasts more than 3 days (72 hours), or recurs.
- pain gets worse or lasts more than 10 days.
- new symptoms occur.
- redness or swelling is present in the painful area. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- remove wrapper
- carefully insert suppository well up into the rectum
- adults and children 12 years and older:
- 1 suppository every 4 to 6 hours while symptoms last
- do not exceed 6 suppositories in any 24 hour period
- **children under 12 years: ask a doctor**

Other information

- store at 20-25°C (68-77°F) or in a cool place
- tamper-evident: suppositories are individually wrapped. Do not use if printed wrapper is opened or damaged.

Inactive ingredient

hydrogenated vegetable oil

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to FeverAll[®] active ingredient

For Adults

Acetaminophen Suppositories USP, 650 mg

Pain Reliever/Fever Reducer

Ages 12 years and older

12 RECTAL SUPPOSITORIES, 650 mg EACH

Padagis[®]



ACETAMINOPHEN PAIN RELIEVER FEVER REDUCER			
acetaminophen suppository			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:45802-730
Route of Administration	RECTAL		
Active Ingredient/Active Moiety			

Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	650 mg	
Inactive Ingredients				
Ingredient Name		Strength		
COTTONSEED OIL (UNII: H3E878020N)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:45802-730-32	50 in 1 CARTON	11/22/2010	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:45802-730-30	12 in 1 CARTON	10/28/2010	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:45802-730-33	100 in 1 CARTON	10/28/2010	11/30/2026
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:45802-730-00	1 in 1 BLISTER PACK; Type 0: Not a Combination Product	10/28/2010	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA070608	10/28/2010	11/30/2026	

Labeler - Padagis Israel Pharmaceuticals Ltd (600093611)

Revised: 2/2025

Padagis Israel Pharmaceuticals Ltd