

**ZEE-SELTZER ANTACID AND PAIN RELIEVER- aspirin, citric acid monohydrate, and sodium bicarbonate tablet, effervescent**  
**ZEE MEDICAL**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Zee-Seltzer™ Antacid and Pain Reliever**

**Drug Facts**

<b>Active ingredients (in each tablet)</b>	<b>Purposes</b>
Aspirin 325 mg (NSAID*)	Pain reliever
Citric acid 1000 mg	Antacid
Sodium bicarbonate (heat-treated) 1916 mg	Antacid

\* nonsteroidal anti-inflammatory drug

**Uses**

**For the temporary relief of:**

- heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches and pain alone

**Warnings**

**Reye's syndrome**

Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed

**Do not use:** if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have asthma, high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Ask a doctor or pharmacist before use if you are**

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- taking a prescription drug for diabetes, gout, or arthritis

**When using this product do not exceed the recommended dosage****Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any signs of stomach bleeding: (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better)
- pain lasts more than 10 days or gets worse
- ringing in the ears or loss of hearing occurs
- redness or swelling is present
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- completely dissolve tablets in 4 ounces of water before taking

Adults and children 12 years and over	2 tablets every 4 hours. Do not exceed 8 tablets in 24 hours
60 years and over	2 tablets every 4 hours. Do not exceed 4 tablets in 24 hours
Children under 12 years	Ask a doctor

**Other information**

- **each tablet contains:** sodium 540 mg
- store at 15-25°C (59-77°F)
- avoid excessive heat above 40°C (104°F)
- in water, this tablet contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate

**Inactive ingredients**

none

**Questions?**

1-800-841-8417 or zeemedical.com

Distributed by ZEE MEDICAL, INC., Irvine, CA 92606

**PRINCIPAL DISPLAY PANEL - 36 Tablet Carton**

**ORIGINAL FLAVOR • EFFERVESCENT**

**Zee-Seltzer™**

**ANTACID & PAIN RELIEVER**

**Aspirin (NSAID)** / Pain reliever

Citric acid / Antacid

Sodium bicarbonate / Antacid

**NBE**

**NATIONAL**

**BRAND**

**EQUIVALENT**

**COMPARE TO ACTIVE**

**INGREDIENTS OF ALKA-SELTZER®\***

**FAST RELIEF**

▼ **HEARTBURN**

▼ **ACID INDIGESTION**

▼ **SOUR STOMACH with HEADACHE**

▼ **BODY ACHES and PAINS**

**36 EFFERVESCENT TABLETS**

**ZEE**

**1408**

1408



ORIGINAL FLAVOR • EFFERVESCENT

# Zee-Seltzer™

ANTACID & PAIN RELIEVER

ORIGINAL FLAVOR • EFFERVESCENT

# Zee-Seltzer™

ANTACID & PAIN RELIEVER



ORIGINAL FLAVOR • EFFERVESCENT  
**Zee-Seltzer™**  
ANTACID & PAIN RELIEVER

**Aspirin (NSAID)** / Pain reliever  
Citric acid / Antacid  
Sodium bicarbonate / Antacid

**NBE** NATIONAL BRAND EQUIVALENT COMPARE TO ACTIVE INGREDIENTS OF ALKA-SELTZER®\*

## FAST RELIEF

- ▼ HEARTBURN
- ▼ ACID INDIGESTION
- ▼ SOUR STOMACH with HEADACHE
- ▼ BODY ACHES and PAINS

**36 EFFERVESCENT TABLETS**

1408



Place two (2) tablets  
in 4 oz. of water.  
Dissolve tablets  
completely.



You do not have to  
drink any of the  
residue that may be  
on the bottom of the  
glass. The medicines  
are in the water.



BOXD-014 REV-5



### Drug Facts

#### Active ingredients (in each tablet)

Aspirin 325 mg (NSAID)*	Pain reliever
Citric acid 1000 mg	Antacid
Sodium bicarbonate (heat-treated) 1916 mg	Antacid

\*nonsteroidal anti-inflammatory drug

#### Purposes

#### Uses

For the temporary relief of: ■ heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains ■ upset stomach with headache from overindulgence in food or drink ■ headache, body aches and pain alone

#### Warnings

**Reye's syndrome:** Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older

■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ have 3 or more alcoholic drinks every day while using this product

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ take more or for a longer time than directed

**Do not use:** if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if:** ■ stomach bleeding warning applies to you

■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic

■ you have asthma, high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Ask a doctor or pharmacist before use if you are:**

■ presently taking a prescription drug. Antacids may interact with certain prescription drugs.

■ taking a prescription drug for diabetes, gout, or arthritis

**When using this product do not exceed the recommended dosage**

**Stop use and ask a doctor if:** ■ an allergic reaction occurs. Seek medical help right away.

■ you experience any signs of stomach bleeding: (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better)

■ pain lasts more than 10 days or gets worse ■ ringing in the ears or loss of hearing occurs

■ redness or swelling is present ■ any new symptoms appear

**DO NOT USE IF PACKET IS TORN OR OPEN.**

### Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

■ completely dissolve tablets in 4 ounces of water before taking

Adults and children 12 years and over	2 tablets every 4 hours. Do not exceed 8 tablets in 24 hours
60 years and over	2 tablets every 4 hours. Do not exceed 4 tablets in 24 hours
Children under 12 years	Ask a doctor

#### Other information

■ each tablet contains: sodium 540 mg

■ store at 15-25°C (59-77°F)

■ avoid excessive heat above 40°C (104°F)

■ in water, this tablet contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate

**Inactive ingredients** none

**Questions?** 1-800-841-8417 or zeemedical.com

Distributed by ZEE MEDICAL, INC., Irvine, CA 92606

\* This product is not manufactured or distributed by Bayer HealthCare, distributor of Alka-Seltzer®.

Rev 0614

MADE IN THE USA

## ZEE-SELTZER ANTACID AND PAIN RELIEVER

aspirin, citric acid monohydrate, and sodium bicarbonate tablet, effervescent

### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:35418-233

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Aspirin (UNII: R16CO5Y76E) (Aspirin - UNII:R16CO5Y76E)	Aspirin	325 mg
Citric Acid Monohydrate (UNII: 2968PHW8QP) (Anhydrous Citric Acid - UNII:XF417D3PSL)	Anhydrous Citric Acid	1000 mg
Sodium Bicarbonate (UNII: 8MDF5V39QO) (Bicarbonate Ion - UNII:HN1ZRA3Q20)	Sodium Bicarbonate	1916 mg

### Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	25mm
Flavor		Imprint Code	C11
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:35418-233-67	18 in 1 CARTON		
1		2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part343	08/14/2006	

Labeler - ZEE MEDICAL (009645623)

Revised: 10/2014

ZEE MEDICAL