

**CALCIUM POLYCARBOPHIL- calcium polycarbophil tablet, film coated
LEADER/ Cardinal Health 110, Inc.**

0339-Cardinal

Drug Facts

Active ingredient (in each caplet)

Calcium polycarbophil 625 mg (equivalent to polycarbophil 500 mg)

Purpose

Bulk-forming laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 12 to 72 hours

Warnings

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product

- do not use for more than 7 days unless directed by a doctor
- do not take more than 8 caplets in a 24-hour period unless directed by a doctor

Stop use and ask a doctor if

you fail to have a bowel movement after use or have rectal bleeding. These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- take each dose of this product with a full glass of water (8oz.) or other fluid. Taking this product without enough liquid may cause choking. See choking warning.
- dosage will vary according to diet, exercise, previous laxative use or severity of constipation
- continued use for 1 to 3 days is normally required to provide full benefit

adults and children 12 years of age and over
children under 12 years of age

2 caplets, 1 to 4 times per day
ask a doctor

Other information

- **each caplet contains:** calcium 125 mg
- do not use if printed seal under cap is torn or missing □ store in a dry place at 15° - 30°C (59° - 86°F)
- protect contents from moisture

Inactive ingredients

caramel, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, silicon dioxide, sodium lauryl sulfate, stearic acid, talc.

Questions or comments?

1-800-231-4670

*This product is not manufactured or distributed by Foundation Consumer Brands, LLC, owner of the registered trademark Fibercon®.

DISTRIBUTED BY CARDINAL HEALTH, DUBLIN, OHIO 43017
www.myleader.com 1-800-200-6313

©2025 CARDINAL HEALTH, LEADER, and LOGOs are trademarks of Cardinal Health. All other marks are the property of their respective owners.

WARNING: This product can expose you to chemicals including lead, which is known to the State of California to cause cancer and birth defects or other reproductive harm. For more information go to www.P65Warnings.ca.gov.

LEADER

NDC 70000-0067-1

Fiber Laxative + Calcium

Calcium Polycarbophil, 625 mg

Bulk-Forming Laxative

Fiber Therapy for Regularity

COMPARE TO FIBERCON® active ingredient*

250 Caplets

LEADER²

NDC 70000-0067-1

Fiber Laxative + Calcium

Calcium Polycarbophil, 625 mg
Bulk-Forming Laxative

Fiber Therapy for Regularity

Actual Size

COMPARE TO FIBERCON® active ingredient*

100% Money Back Guarantee

250 CAPLETS

WARNING: This product can expose you to chemicals including lead, which is known to the State of California to cause cancer and birth defects or other reproductive harm. For more information go to www.PBSWarnings.ca.gov.

Drug Facts

Active Ingredient (in each caplet)
Calcium polycarbophil 625 mg (equivalent to polycarbophil 500 mg).....Bulk-forming laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 12 to 72 hours

Warnings

Children: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing, if you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product

- do not use for more than 7 days unless directed by a doctor
- do not take more than 6 caplets in a 24-hour period unless directed by a doctor

Stop use and ask a doctor if you fail to have a bowel movement after use or have rectal bleeding. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- Take each dose of this product with a full glass of water (8oz.) or other fluid. Taking this product without enough liquid may cause choking. See choking warning.
- Dosage will vary according to diet, exercise, previous laxative use or severity of constipation
- continued use for 1 to 3 days is normally required to provide full benefit.
- adults and children 12 years of age and over 2 caplets, 1 to 4 times per day
- children under 12 years of age ask a doctor

Other Information

- each caplet contains: calcium 125 mg
- do not use if printed seal under cap is torn or missing
- store in a dry place at 15° - 30°C (59° - 86°F)
- protect contents from moisture

Inactive Ingredients caramel, croscopidone, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, silicon dioxide, sodium lauryl sulfate, stearic acid, talc.

Questions or comments? 1-800-231-4670

CIN 5669397 REV. 09/25
DISTRIBUTED BY CARDINAL HEALTH, MURBIN, OHIO 43007
www.myLeader.com 1-800-200-5333

9629514027 9

©2025 CARDINAL HEALTH, LEADER, and LOGOS are trademarks of Cardinal Health. All other marks are the property of their respective owners.

This product is not manufactured or distributed by Foundation Consumer Brands, LLC. Consumer the registered trademark "FIBERCON".

653038
LOT & EXP. DATE:

CALCIUM POLYCARBOPHIL

calcium polycarbophil tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0067
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALCIUM POLYCARBOPHIL (UNII: 8F049NKY49) (POLYCARBOPHIL - UNII:W25LM17A4W)	CALCIUM POLYCARBOPHIL	625 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSPVIDONE (UNII: 2S7830E561)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CARAMEL (UNII: T9D99G2B1R)	

Product Characteristics

Color	white (off white)	Score	2 pieces
Shape	OVAL	Size	19mm
Flavor		Imprint Code	CPC;339
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0067-1	250 in 1 BOTTLE; Type 0: Not a Combination Product	11/18/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	11/18/2020	

Labeler - LEADER/ Cardinal Health 110, Inc. (063997360)

Revised: 11/2025

LEADER/ Cardinal Health 110, Inc.