ASPIRIN 81MG- aspirin 81mg tablet Strive Pharmaceuticals Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Right Remedies Aspirin Low Dose Aspirin 81mg

Aspirin 81mg (NSAID)

colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10, FD&C yellow #6, methacrylic acid copolymer, microcrytalline cellulose, silicon dioxide, stearic acid, talc, titanium dioxide, triethyl citrate

* drink a full glass of water with each dose **SEE OVERDOSE WARNING**.

adults and children 12	take 4 to 8 tablets every 4 hours not to exceed 48 tablets in
years of age and over	24 hours unless directed by a doctor
children under 12 years of age	ask a doctor

^{*} for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- Hives
- Facial swelling
- Asthma (wheezing)
- Shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- Are age 60 or older
- Have had stomach ulcers or bleeding problems
- Take a blood thinning (anticoagulant)or steroid drug
- Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- Have 3 or more alcoholic drinks everyday while using this product
- Take more or for a longer time than directed.

Do not use

^{*}ask your doctor about other uses.

- If you are allergic to aspirin or any other pain reliever/fever reducer.\
- If you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if

- Stomach bleeding warning applies to you
- You have a history of stomach problems such as heartburn
- You have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- You are taking a diuretic
- You have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- Gout
- Diabetes
- Arthritis

Stop use and ask a doctor if

- An allergic reaction occurs. Seek medical help right away.
- You experience any of the following signs of stomach bleeding:
- Feel faint
- Vomit blood
- Have bloody or black stools
- Have stomach pain that does not get better
- Pain gets worse or last more than 10 days
- Redness or swelling is present
- Fever gets worse or last more than 3 days
- New symptoms occur
- Ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

OVERDOSE WARNING: In case of overdose, get medical help or contact a Poison Control right away. Prompt medical attention is critical for adults as well as children eve if you do not notice any signs or symptoms.

Keep out of reach of children.

OVERDOSE WARNING: In case of overdose, get medical help or contact a Poison Control right away. Prompt medical attention is critical for adults as well as children eve if you do not notice any signs or symptoms.

Pain reliever

- * store at room temperature 20 25°C (68 77°F).
- * avoid excessive heat, cold and humidity.
- * close cap tightly after use.



ASPIRIN 81MG

aspirin 81mg tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-158
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients	
Ingredient Name	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)

METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A (UNII: NX76LV5T8J)

STEARIC ACID (UNII: 4ELV7Z65AP)

TALC (UNII: 7SEV7J4R1U)

CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)

CROSCARMELLOSE SODIUM (UNII: M280L1HH48)

Product Characteristics			
Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	I1
Contains			

Packaging				
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	1 NDC:70692-158-	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/18/2020	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	11/18/2020	
OTC monograph not			

Labeler - Strive Pharmaceuticals Inc. (080028013)

Revised: 4/2022 Strive Pharmaceuticals Inc.