

**RITE AID ANTACID AND PAIN RELIEF- aspirin, citric acid, sodium bicarbonate tablet, effervescent**  
**Rite Aid**

-----  
**Rite Aid Antacid & Pain Relief**

**Active ingredients**

**(in each effervescent tablet)**

Anhydrous citric acid 1000 mg

Aspirin 325 mg **(NSAID)\***

Sodium bicarbonate (heat-treated) 1916 mg

\*Nonsteroidal anti-inflammatory drug

**Purpose**

Anhydrous citric acid 1000 mg. . . . . Antacid

Aspirin 325 mg **(NSAID)\***. . . . . Pain reliever

Sodium bicarbonate (heat-treated) 1916 mg. . . . . Antacid

\*Nonsteroidal anti-inflammatory drug

**Uses**

**Uses** for the temporary relief of

- heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood

thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

### **Do not use**

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients.

### **Ask a doctor before use if**

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma ■ you have a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

■ presently taking a prescription drug. Antacids may interact with certain prescription drugs. ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

### **When using this product**

**When using this product** do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician.

### **Stop use and ask a doctor if**

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

### **If pregnant or breast-feeding**

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

**Keep out of reach of children.**

### **In case of overdose**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

■ do not exceed recommended dosage

■ to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking

adults and children 12 years and over (up to 60 years of age): 2 tablets every 4 hours - do not exceed 8 tablets in 24 hours

60 years and over: 2 tablets every 4 hours - do not exceed 4 tablets in 24 hours

Children under 12 years - consult a doctor

### **Other information**

■ **each tablet contains:** sodium 524 mg

■ store at room temperature (59°-86°F)

### **Inactive ingredients**

■ none

### **PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

PN: 42124 RAC3  
R: 04/23



**SATISFACTION GUARANTEE**  
If you're not satisfied, we'll happily refund your money.  
The maker of Rite Aid® is distributed or licensed by the maker of Rite Aid®.

DISTRIBUTED BY: RITE AID  
200 NEUMBERY COMMONS, ETTERS, PA 17319  
www.riteaid.com  
MANUFACTURED IN THE U.S.A. WITH DOMESTIC AND IMPORTED INGREDIENTS.

Compare to the active ingredients in Rite Aid's **Aspirin**®.

# ANTACID & PAIN RELIEF

**EFFERVESCENT TABLETS**

ANHYDROUS CITRIC ACID/ANTACID  
ASPIRIN (NSAID)/PAIN RELIEVER  
SODIUM BICARBONATE/ANTACID

Fast relief of acid indigestion, upset stomach, heartburn with headache, body aches & pains

ORIGINAL FLAVOR

24 EFFERVESCENT TABLETS

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not exceed recommended dosage
- to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking
- adults and children 17 years and over (up to 60 years of age) 2 tablets every 4 hours do not exceed 8 tablets in 24 hours
- 60 years and over 2 tablets every 4 hours do not exceed 4 tablets in 24 hours
- Children under 17 years consult a doctor

**Other Information**

- each tablet contains sodium 524 mg
- store at room temperature (59°-86°F)
- none

**Inactive Ingredients**



ANHYDROUS CITRIC ACID/ANTACID  
ASPIRIN (NSAID)/PAIN RELIEVER  
SODIUM BICARBONATE/ANTACID

# ANTACID & PAIN RELIEF

**EFFERVESCENT TABLETS**

**Drug Facts**

**Active ingredients (in each effervescent tablet)**

Active ingredient	Purpose
Anhydrous citric acid 1000 mg	Antacid
Aspirin 325 mg (NSAID)*	Pain Reliever
Sodium bicarbonate (heat-treated) 1916 mg	Antacid

\*Nonsteroidal anti-inflammatory drug

**Uses** for the temporary relief of

- heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

**Drug Facts (continued)**

- take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer. If you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

**When using this product do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician.**

**Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse
- new symptoms occur
- redness or swelling is present
- ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

42124 RAC3



**RITE AID ANTACID AND PAIN RELIEF**  
aspirin, citric acid, sodium bicarbonate tablet, effervescent

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-4212
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	ANHYDROUS CITRIC ACID	1000 mg
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1ZRA3Q20)	SODIUM BICARBONATE	1916 mg

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	25mm
<b>Flavor</b>		<b>Imprint Code</b>	L421
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:11822-4212-4	12 in 1 CARTON	03/16/2020	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:11822-4212-7	36 in 1 CARTON	12/09/2022	
2		2 in 1 POUCH; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	03/16/2020	

**Labeler** - Rite Aid (014578892)