

NICOTINE POLACRILEX COATED MINT- nicotine polacrilex gum, chewing
Chain Drug Consortium, LLC

Nicotine Polacrilex Gum 2 mg and 4 mg Coated Mint - Premier Value (Chain Drug Consortium, LLC)

Drug Facts

Active ingredient (in each chewing piece) - 2 mg

Nicotine Polacrilex 2 mg (nicotine)

Active ingredient (in each chewing piece) - 4 mg

Nicotine Polacrilex 4 mg (nicotine)

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug

- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions - 2 mg

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum according to the following 12 week schedule:

| Weeks 1 to 6 | Weeks 7 to 9 | Weeks 10 to 12 |
|----------------------------|----------------------------|----------------------------|
| 1 piece every 1 to 2 hours | 1 piece every 2 to 4 hours | 1 piece every 4 to 8 hours |

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information - 2 mg

- **each piece contains:** calcium 115 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Directions - 4 mg

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum according to the following 12 week schedule:

| Weeks 1 to 6 | Weeks 7 to 9 | Weeks 10 to 12 |
|-------------------------------|-------------------------------|-------------------------------|
| 1 piece every 1 to 2 hours | 1 piece every 2 to 4 hours | 1 piece every 4 to 8 hours |

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information - 4 mg

- **each piece contains:** calcium 120 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients - 2 mg

acacia, acesulfame potassium, carnauba wax, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide, xylitol.

Inactive ingredients - 4 mg

acacia, acesulfame potassium, carnauba wax, D&C yellow #10 lake, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide, xylitol.

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal display panel 2 mg

COMPARE TO THE ACTIVE INGREDIENT IN NICORETTE® GUM*

Nicotine Polacrilex Gum USP, 2 mg (nicotine)

STOP SMOKING AID

FOR THOSE WHO SMOKING THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg

COATED MINT 2 mg

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew **at least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a Support Program** as directed in the enclosed User's Guide.
 - not for sale to those under 18 years of age
 - proof of age required
 - not for sale in vending machines or from any source where proof of age cannot be verified

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

Distributed By:

Pharmacy Value Alliance, LLC

407 East Lancaster Avenue,

Wayne, PA 19087

Package label 2 mg 284

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** – Chew at least **9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** – Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a Support Program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit.



Peel off backing starting at corner with loose edge.



Push gum through foil.



COMPARE TO THE ACTIVE INGREDIENT IN NICORETTE® GUM*

2 mg

Nicotine Polacrilex Gum USP, 2 mg (nicotine)

STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.
If you smoke your first cigarette **WITHIN** 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg

COATED MINT

100 PIECES, 2 MG EACH



Nicotine Polacrilex Gum USP, 2 mg (nicotine)

STOP SMOKING AID

COATED MINT



PLD-017A PC007574

THEFT SURVEILLANCE TAG AREA

machines or from any source where proof of age cannot be verified

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.



Distributed By:
Pharmacy Value Alliance, LLC
407 East Lancaster Avenue,
Wayne, PA 19087

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

Call 1-877-753-9938 Monday-Friday 9AM-5PM EST

Questions or comments?

nicotine, sodium carbonate, sac, butylam device, yfth
scent, gum base, hydroxypropyl cellulose, mannitol, oza, sodium

Stop use and ask a doctor if
 ■ you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
 ■ you get symptoms of nicotine overdose occur
 ■ regular heartbeat or palpitations occur
 ■ mouth, teeth, or jaw problems occur
 ■ you have symptoms of an allergic reaction (such as difficulty breathing or rash)
Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. When used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.
Directions
 ■ If you are under 18 years of age, ask a doctor before use
 ■ Before using this product, read the enclosed User's Guide for complete directions and other important information.
 ■ If you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum.
 ■ If you smoke your first cigarette more than 30 minutes of waking up, use 2 mg nicotine gum.

■ not for sale to those under 18 years of age
 ■ proof of age required
 ■ not for sale in vending

Nicotine Gum USP, Polacrilex
2 mg (nicotine)
 STOP SMOKING AID
 COATED MINT



Drug Facts (continued)

Warnings

- If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement products. This medicine is intended to be used with other smoking cessation products. However, the risk to your child from this medicine is not fully known.
- Ask a doctor before use if you have a heart condition, recent heart attack, or regular heartbeat. Nicotine can increase your heart rate.
- High blood pressure not controlled with medication. Nicotine can increase blood pressure.
- History of seizures

Use

- Use as directed on the label, including nicotine strength, associated with quitting smoking.

Active ingredient (in each chewing piece) Nicotine prochlorate, 2 mg (nicotine)

Purpose Stop smoking aid

Other information

- Each piece contains nicotine 1.15 mg, sodium 6 mg
- Store at 20°-25°C (68°-77°F)
- Protect from light

Drug Facts (continued)

■ If you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

| | |
|----------------|-----------------------------|
| Weeks 1 to 6 | 1 piece every 2 to 4 hours |
| Weeks 7 to 9 | 1 piece every 4 to 8 hours |
| Weeks 10 to 12 | 1 piece every 8 to 12 hours |

■ Nicotine gum is a medication and must be used in a certain way to get the best results.

■ Chew the gum slowly until it begins to melt. Then park it between your cheek and gum. When the target is gone, begin chewing again, until the target returns. Repeat the process until most of the target is gone (about 30 minutes).

■ Do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece.

■ Do not repeat your chances of quitting, use at least 9 pieces per day for the first 6 weeks.

■ If you experience dizziness or frequent dizziness, you may use a second piece when the hour. However, do not continually use one piece after another since this may cause you nausea, headache, dizziness, or other side effects.

■ Do not use more than 24 pieces a day.

■ It is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

ALWAYS RECYCLE YOUR PACKAGING TO HELP PROTECT THE EARTH AND YOUR HEALTH.

Nicotine Gum USP, Polacrilex
2 mg (nicotine)
 STOP SMOKING AID
 COATED MINT



PREMIER VALUE Nicotine Gum Coated Mint

Package label 2 mg 280



PRIMIER VALUE Nicotine Gum Coated Mint

Package label 2 mg 559

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use With a Support Program** as directed in the enclosed User's Guide.

Drug Facts (continued)

Other information

- each piece contains: calcium 115 mg, sodium 8 mg
- store at 20-25°C (68 -77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium, carnauba wax, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide, xylitol

Questions or comments?

Call 1-877-753-3935
Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

Distributed By: Pharmacy Value Alliance, LLC
407 East Lancaster Avenue,
Wayne, PA 19087



If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

See inside for more details or visit www.nixthehabit.com.

Product of Ireland
PLD-E617A
FC008071



COMPARE TO THE ACTIVE INGREDIENT IN NICORETTE® GUM*

2 mg

Nicotine Polacrilex Gum USP, 2 mg (nicotine)

STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicotine Polacrilex Gum USP, 4 mg

COATED MINT

Sugar-free



20 PIECES, 2 MG EACH



THEFT SURVEILLANCE TAG AREA

■ not for sale to those under 18 years of age
■ proof of age required
■ not for sale in vending machines or from any source where proof of age cannot be verified



Drug Facts

Active ingredient (in each chewing piece) Purpose
Nicotine polacrilex.....2 mg (nicotine).....Stop smoking aid

Use ■ reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

| Weeks 1 to 6 | Weeks 7 to 9 | Weeks 10 to 12 |
|----------------------------|----------------------------|----------------------------|
| 1 piece every 1 to 2 hours | 1 piece every 2 to 4 hours | 1 piece every 4 to 8 hours |

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.



PREMIER VALUE Nicotine Polacrilex Gum Coated Mint Flavor

Principal display panel 4 mg

COMPARE TO THE ACTIVE INGREDIENT IN NICORETTE® GUM*

Nicotine Polacrilex Gum USP, 4 mg (nicotine)

STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicotine Polacrilex Gum USP, 2 mg

COATED MINT 4 MG

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew **at least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use With a Support Program** as directed in the enclosed User's Guide
 - not for sale to those under 18 years of age
 - proof of age required
 - not for sale in vending machines or from any source where proof of age cannot be required

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

Distributed By:

Pharmacy Value Alliance, LLC

407 East Lancaster Avenue,

Wayne, PA 19087

Package label 4 mg 560

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use With a Support Program** as directed in the enclosed User's Guide.

Drug Facts (continued)

Other information

- each piece contains: calcium 120 mg, sodium 8 mg
- store at 20-25°C (68 -77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium, carnauba wax, D&C yellow #10 aluminum lake, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide, xylitol

Questions or comments?

Call 1-877-753-3935
Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

Distributed By:
Pharmacy Value Alliance, LLC
407 East Lancaster Avenue,
Wayne, PA 19087



If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

See inside for more details or visit www.nixthehabit.com.

Product of Ireland
PLD-D623A
FC008083



COMPARE TO THE ACTIVE INGREDIENT IN NICORETTE® GUM*

4 mg

Nicotine Polacrilex Gum USP, 4 mg (nicotine)

STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicotine Polacrilex Gum USP, 2 mg

COATED MINT



actual size

20 PIECES, 4 MG EACH



THEFT SURVEILLANCE TAG AREA

not for sale to those under 18 years of age
proof of age required
not for sale in vending machines or from any source where proof of age cannot be verified



Push gum through foil.



Peel off backing starting at corner with loose edge.



To remove the gum, tear off single unit.

Drug Facts

Active ingredient (in each chewing piece) Purpose
Nicotine polacrilex.....4 mg (nicotine)..... Stop smoking aid

Use ■ reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum according to the following 12 week schedule:

| Weeks 1 to 6 | Weeks 7 to 9 | Weeks 10 to 12 |
|----------------------------|----------------------------|----------------------------|
| 1 piece every 1 to 2 hours | 1 piece every 2 to 4 hours | 1 piece every 4 to 8 hours |

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.



8 40986 03110 6

PREMIER VALUE Nicotine Polacrilex Gum Coated Mint

Package label 4 mg 285

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** – Chew at least **9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** – Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a Support Program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit.



Peel off backing starting at corner with loose edge.



Push gum through foil.



COMPARE TO THE ACTIVE INGREDIENT IN NICORETTE® GUM*

4 mg

Nicotine Polacrilex Gum USP, 4 mg (nicotine)

STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.
If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use Nicotine Polacrilex Gum USP, 2 mg

COATED MINT

100 PIECES, 4 MG EACH



actual size



Nicotine Polacrilex Gum USP, 4 mg (nicotine)


STOP SMOKING AID

COATED MINT



PRODUCT OF IRELAND
PLD-823A F000366

Thrift surveillance
age cannot be verified.
source where proof of
machines or from any
NOT FOR SALE IN VERMONT



If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

Distributed By:
Pharmacy Value Alliance, LLC
407 East Lancaster Avenue,
Wayne, PA 19087

*The product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, LP, distributor of Nicorette® Gum.

Call 1-877-750-9505 Monday-Friday 9AM-5PM EST

Questions or comments?

■ You get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
■ You have symptoms of an allergic reaction (such as difficulty breathing or rash)
■ Keep out of reach of children and pets. Pieces of nicotine gum may have paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

■ If you are under 18 years of age, ask a doctor before use
■ Before using this product, read the enclosed User's Guide for complete directions and other important information
■ Begin using the gum on your quit day
■ If you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum



PREMIER VALUE Nicotine Gum Coated Mint

NICOTINE POLACRILEX COATED MINT

nicotine polacrilex gum, chewing

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:68016-280 |
| Route of Administration | BUCCAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|----------------------------------------------------------|-------------------|----------|
| NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R) | NICOTINE | 2 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---------------------------------------------------------|----------|
| ACACIA (UNII: 5C5403N26O) | |
| ACESULFAME POTASSIUM (UNII: 23OV73Q5G9) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH) | |
| MAGNESIUM OXIDE (UNII: 3A3U0GI71G) | |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) | |
| SODIUM CARBONATE (UNII: 45P3261C7T) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| XYLITOL (UNII: VCQ006KQ1E) | |

Product Characteristics

| | | | |
|----------|--------------------|--------------|----------|
| Color | white | Score | no score |
| Shape | SQUARE | Size | 14mm |
| Flavor | MINT (coated mint) | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:68016-280-49 | 4 in 1 CARTON | 07/01/2013 | |
| 1 | | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|------------------------------------------|----------------------|--------------------|
| ANDA | ANDA078699 | 07/01/2013 | 12/31/2026 |

NICOTINE POLACRILEX COATED MINT

nicotine polacrilex gum, chewing

Product Information

| | | | |
|--------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:68016-284 |
|--------------|----------------|--------------------|---------------|

Route of Administration BUCCAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|----------------------------------------------------------|-------------------|----------|
| NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R) | NICOTINE | 2 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---------------------------------------------------------|----------|
| ACACIA (UNII: 5C5403N26O) | |
| ACESULFAME POTASSIUM (UNII: 23OV73Q5G9) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH) | |
| MAGNESIUM OXIDE (UNII: 3A3U0GI71G) | |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) | |
| SODIUM CARBONATE (UNII: 45P3261C7T) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| XYLITOL (UNII: VCQ006KQ1E) | |

Product Characteristics

| | | | |
|----------|--------------------|--------------|----------|
| Color | white | Score | no score |
| Shape | SQUARE | Size | 14mm |
| Flavor | MINT (coated mint) | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:68016-284-44 | 10 in 1 CARTON | 06/01/2013 | |
| 1 | | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|------------------------------------------|----------------------|--------------------|
| ANDA | ANDA078699 | 06/01/2013 | 12/31/2026 |

NICOTINE POLACRILEX COATED MINT

nicotine polacrilex gum, chewing

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:68016-281 |
| Route of Administration | BUCCAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|-----------------------------------------------------------------|-------------------|----------|
| NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R) | NICOTINE | 4 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|----------------------------------------------------------------|----------|
| ACACIA (UNII: 5C5403N26O) | |
| ACESULFAME POTASSIUM (UNII: 230V73Q5G9) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH) | |
| MAGNESIUM OXIDE (UNII: 3A3U0GI71G) | |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) | |
| SODIUM CARBONATE (UNII: 45P3261C7T) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| XYLITOL (UNII: VCQ006KQ1E) | |

Product Characteristics

| | | | |
|-----------------|--------------------|---------------------|----------|
| Color | yellow | Score | no score |
| Shape | SQUARE | Size | 14mm |
| Flavor | MINT (coated mint) | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:68016-281-49 | 4 in 1 CARTON | 06/01/2013 | |
| 1 | | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|------------------------------------------|----------------------|--------------------|
| ANDA | ANDA078697 | 06/01/2013 | 12/31/2026 |

NICOTINE POLACRILEX COATED MINT

nicotine polacrilex gum, chewing

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:68016-285 |
| Route of Administration | BUCCAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|-----------------------------------------------------------------|--------------------------|-----------------|
| NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R) | NICOTINE | 4 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|----------------------------------------------------------------|-----------------|
| ACACIA (UNII: 5C5403N26O) | |
| ACESULFAME POTASSIUM (UNII: 23OV73Q5G9) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH) | |
| MAGNESIUM OXIDE (UNII: 3A3U0GI71G) | |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) | |
| SODIUM CARBONATE (UNII: 45P3261C7T) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| XYLITOL (UNII: VCQ006KQ1E) | |

Product Characteristics

| | | | |
|-----------------|--------------------|---------------------|----------|
| Color | yellow | Score | no score |
| Shape | SQUARE | Size | 14mm |
| Flavor | MINT (coated mint) | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:68016-285-44 | 10 in 1 CARTON | 09/01/2013 | |
| 1 | | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------------|-------------------------------------------------|-----------------------------|---------------------------|
| ANDA | ANDA078697 | 09/01/2013 | 12/31/2026 |

NICOTINE POLACRILEX COATED MINT

nicotine polacrilex gum, chewing

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:68016-560 |
| Route of Administration | BUCCAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|-----------------------------------------------------------------|-------------------|----------|
| NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R) | NICOTINE | 4 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|----------------------------------------------------------------|----------|
| ACACIA (UNII: 5C5403N26O) | |
| ACESULFAME POTASSIUM (UNII: 230V73Q5G9) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH) | |
| MAGNESIUM OXIDE (UNII: 3A3U0GI71G) | |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) | |
| SODIUM CARBONATE (UNII: 45P3261C7T) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| XYLITOL (UNII: VCQ006KQ1E) | |

Product Characteristics

| | | | |
|-----------------|--------------------|---------------------|----------|
| Color | yellow | Score | no score |
| Shape | SQUARE | Size | 14mm |
| Flavor | MINT (coated mint) | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:68016-560-00 | 2 in 1 CARTON | 10/01/2014 | |
| 1 | | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|------------------------------------------|----------------------|--------------------|
| ANDA | ANDA078697 | 10/01/2014 | 12/31/2026 |

NICOTINE POLACRILEX COATED MINT

nicotine polacrilex gum, chewing

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:68016-559 |
| Route of Administration | BUCCAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|-----------------------------------------------------------------|-------------------|----------|
| NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R) | NICOTINE | 2 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|----------------------------------------------------------------|----------|
| ACACIA (UNII: 5C5403N26O) | |
| ACESULFAME POTASSIUM (UNII: 23OV73Q5G9) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH) | |
| MAGNESIUM OXIDE (UNII: 3A3U0GI71G) | |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) | |
| SODIUM CARBONATE (UNII: 45P3261C7T) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| XYLITOL (UNII: VCQ006KQ1E) | |

Product Characteristics

| | | | |
|-----------------|--------------------|---------------------|----------|
| Color | white | Score | no score |
| Shape | SQUARE | Size | 14mm |
| Flavor | MINT (coated mint) | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:68016-559-00 | 2 in 1 CARTON | 10/01/2014 | |
| 1 | | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing | Application Number or Monograph | Marketing Start | Marketing End |
|-----------|---------------------------------|-----------------|---------------|
|-----------|---------------------------------|-----------------|---------------|

| Category | Citation | Date | Date |
|----------|------------|------------|------------|
| ANDA | ANDA078699 | 10/01/2014 | 12/31/2026 |

Labeler - Chain Drug Consortium, LLC (101668460)

Revised: 6/2025

Chain Drug Consortium, LLC