

GERICARE SORBITOL SOLUTION- sorbitol liquid
GERI-CARE PHARMACEUTICALS, CORP

GeriCare- Sorbitol 758

Active Ingredient

Sorbitol 70% USP

Purpose

Laxative

Uses

- relieves occasional constipation and irregularity
- generally produces bowel movement in 1/4 to 1 hour when used rectally
- as a pharmaceutical aide (sweetner)
- for other uses, ask your doctor

Warnings

For rectal use only

Do not use

- when abdominal pain, nausea, or vomiting are present
- for more than one week unless directed by a doctor

Ask a doctor before use if

- are taking mineral oil
- have noticed a sudden change in bowel habits that lasts over 2 weeks

Stop use and ask a doctor if you have a rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.

If pregnant or breast feeding ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions Do not exceed recommended dose

| AGE | DOSE |
|---------------------------------------|--|
| Adults and children 12 years and over | For rectal use only. Enema dosage is 120 mL of a 25 to 30% w/v solution (1 part of this product with 2.3 parts water) in a single daily dose as needed, or as directed by your doctor. |
| Children under 12 years of age | Ask a doctor |

Inactive ingredients water

Questions or comments? 1-800-540-3765

| Drug Facts | |
|---|----------------------------|
| Active ingredient Sorbitol 70%, USP | Purpose Laxative |
| Uses | |
| <ul style="list-style-type: none"> relieves occasional constipation and irregularity generally produces bowel movement in ¼ to 1 hour when used rectally as a pharmaceutical aide (sweetener) for other uses, ask your doctor | |
| Warnings | |
| For rectal use only | |
| Do not use - when abdominal pain, nausea, or vomiting are present - for more than one week unless directed by a doctor. | |
| Ask a doctor before use if you - are taking mineral oil - have noticed a sudden change in bowel habits that lasts over 2 weeks | |
| Stop use and ask a doctor if you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition. | |
| If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. | |

Code#: L-126 Rev. 12/20
 Lot.#
 Exp. Date:



SORBITOL SOLUTION USP
70% W/W
LAXATIVE

TAMPER EVIDENT: Do not use this product if inner foil seal over the mouth of bottle is missing or broken.

16 FL OZ (1 PT) (473 mL)

| Drug Facts (continued) | |
|--|--|
| Directions - do not exceed recommended dose | |
| Age | Dose |
| Adults and children 12 years and over | For rectal use only. Enema dosage is 120 mL of a 25 to 30% w/v solution (1 part of this product with 2.3 parts water) in a single daily dose as needed, or as directed by your doctor. |
| Children under 12 years of age | Ask a doctor. |
| Other information | |
| <ul style="list-style-type: none"> store at room temperature 15°- 30° C (59°-86°F) below 59°F cloudiness and thickening may occur; warming will restore clarity and fluidity without affecting product quality do not freeze Pharmacist-Preserve and dispense in tight, light-resistant container with a child resistant cap as defined in the USP | |
| Inactive ingredient: water | |
| Questions or comments? 1-800-540-3765 | |



DIST. BY: GERI-CARE PHARMACEUTICALS CORP.
 1850 63rd Street
 Brooklyn, NY 11204

GERICARE SORBITOL SOLUTION

sorbitol liquid

| Product Information | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:57896-758 |
| Route of Administration | RECTAL | | |

| Active Ingredient/Active Moiety | | |
|---|-------------------|-----------------|
| Ingredient Name | Basis of Strength | Strength |
| SORBITOL (UNII: 506T60A25R) (SORBITOL - UNII:506T60A25R) | SORBITOL | 70 mg in 100 mL |

| Inactive Ingredients | |
|---------------------------------|----------|
| Ingredient Name | Strength |
| WATER (UNII: 059QF0KO0R) | |

| Packaging | | | | |
|-----------|------------------|---|----------------------|--------------------|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:57896-758-16 | 473 mL in 1 BOTTLE; Type 0: Not a Combination Product | 01/01/2020 | |

| Marketing Information | | | |
|-----------------------|--|----------------------|--------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| | | | |

OTC Monograph Drug M007

01/01/2020

Labeler - GERI-CARE PHARMACEUTICALS, CORP (611196254)

Revised: 12/2024

GERI-CARE PHARMACEUTICALS, CORP