

**RIGHT REMEDIES ALLERGY RELIEF- diphenhydramine hydrochloride capsule**  
**Strive Pharmaceuticals Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**RIGHT REMEDIES Allergy Relief**

**Drug Facts**

**Active ingredient (in each capsule)**

Diphenhydramine HCl 25 mg

**Purpose**

Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: • runny nose • sneezing • itchy, watery eyes • itching of the nose or throat  
temporarily relieves these symptoms due to the common cold: • runny nose • sneezing

**Warnings**

**Do not use** • to make a child sleepy  
• with any other product containing diphenhydramine, even one used on skin

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers.

**When using this product**

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control

Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

### **Directions**

- take every 4 to 6 hours, or as directed by a doctor
- do not take more than 6 times in 24 hours

adults and children 12 years and older	1 to 2 capsules
children 6 to under 12 years	1 capsule
children under 6 years old	do not use

### **Other information**

- store between 20-25°C (68-77°F) • protect from light.

### **Inactive ingredients**

butylparaben, D&C red #28, edible ink, FD&C blue #1, FD&C red #40, gelatin, lactose monohydrate, magnesium stearate, methylparaben, polysorbate-80, propylparaben, purified water, sodium lauryl sulphate, starch

### **Questions or comments?**

**1-888-577-8033** Monday-Friday 8am-4pm EST

Compare to the active ingredient of **Benadryl® Allergy\***

- Sneezing
- Runny nose
- Itchy, watery eyes
- Itchy throat


**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Benadryl® Allergy

Distributed by:  
Strive Pharmaceuticals Inc.  
East Brunswick, NJ 08816

Product of India  
Packaged and Quality Assured in the USA

# Packaging



Compare to the active ingredient of Benadryl® Allergy\*


NDC 70692-832-11


# Allergy Relief

Diphenhydramine HCl 25 mg

**Antihistamine**

- Sneezing
- Runny nose
- Itchy, watery eyes
- Itchy throat





Actual Size

**Drug Facts**

**Active ingredient (in each capsule)**  
Diphenhydramine HCl 25 mg.....Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat temporarily relieves these symptoms due to the common cold: ■ runny nose ■ sneezing

**Warnings**  
Do not use ■ to make a child sleepy ■ with any other product containing diphenhydramine, even one used on skin

**Ask a doctor before use if you have**  
■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.**

**When using this product**  
■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**  
■ take every 4 to 6 hours, or as directed by a doctor ■ do not take more than 6 times in 24 hours


adults and children 12 years and older	1 to 2 capsules
children 6 to under 12 years	1 capsule
children under 6 years old	do not use

**Other information**  
■ store between 20°-25°C (68°-77°F) ■ protect from light.

**Inactive ingredients** butylparaben, D&G red #28, ethyl red #1, FD&C blue #1, FD&C red #40, gelatin, lactose monohydrate, magnesium stearate, polyethylene glycol, polyethylene glycol, purified water, sodium lauryl sulfate, starch

**Questions or comments?** 1-888-577-4033 Monday-Friday 8am-4pm EST

\*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Benadryl® Allergy.  
Distributed by:  
Stive Pharmaceuticals Inc.  
East Brunswick, NJ 08816  
Product of India,  
Packaged and Quality Assured in the USA



Product of India,  
Packaged and Quality Assured in the USA

Lot: LB7069283211  
Exp: REV:00-06:2021

## DRUG FACTS LABEL

### Drug Facts

<b>Active ingredient (in each capsule)</b> Diphenhydramine HCl 25 mg.....	<b>Purpose</b> Antihistamine
--	---------------------------------

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat temporarily relieves these symptoms due to the common cold: ■ runny nose ■ sneezing

**Warnings**  
Do not use ■ to make a child sleepy ■ with any other product containing diphenhydramine, even one used on skin

**Ask a doctor before use if you have**  
■ a breathing problem such as emphysema or chronic bronchitis  
■ glaucoma  
■ trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.**

**When using this product**  
■ marked drowsiness may occur  
■ avoid alcoholic drinks  
■ alcohol, sedatives, and tranquilizers may increase drowsiness  
■ be careful when driving a motor vehicle or operating machinery  
■ excitability may occur, especially in children

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**  
■ take every 4 to 6 hours, or as directed by a doctor  
■ do not take more than 6 times in 24 hours

adults and children 12 years and older	1 to 2 capsules
children 6 to under 12 years	1 capsule
children under 6 years old	do not use

**Other information**

■ store between 20-25°C (68-77°F) ■ protect from light.

**Inactive ingredients** butylparaben, D&C red #28, edible ink, FD&C blue #1, FD&C red #40, gelatin, lactose monohydrate, magnesium stearate, methylparaben, polysorbate-80, propylparaben, purified water, sodium lauryl sulphate, starch

**Questions or comments?** 1-888-577-8033 Monday-Friday 8am-4pm EST

**RIGHT REMEDIES ALLERGY RELIEF**

diphenhydramine hydrochloride capsule

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70692-832
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII: 8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>BUTYLPARABEN</b> (UNII: 3QPI1U3FV8)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	

**Product Characteristics**

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	capsule (Cylindrical)	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	P25
<b>Contains</b>			

**Packaging**

Marketing Start      Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-832-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	12/20/2021	
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	12/20/2021		

**Labeler** - Strive Pharmaceuticals Inc (080028013)

Revised: 12/2021

Strive Pharmaceuticals Inc