

**SMOOTH TEXTURE ORANGE FLAVOR - psyllium husk powder, for suspension
McKesson**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Sunmark Fiber Smooth Texture Orange Flavor

Active ingredient (per each rounded tablespoon=12 g)

Psyllium husk 3.4 g

Purpose

Bulk-forming laxative

Uses

- effective in treating occasional constipation and restoring regularity

Warnings

Allergy alert: May cause allergic reaction in people sensitive to ingested or inhaled psyllium

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have

- abdominal pain, nausea or vomiting
- a sudden change in bowel habits persisting for 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Laxatives may affect how other drugs work.

When using this product

- do not exceed the maximum recommended daily dosage in a 24-hour period
- you may experience changes in bowel habits or minor bloating as your body adjusts to increased fiber intake

Stop use and ask a doctor if

- constipation lasts more than 7 days
- rectal bleeding occurs

These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take at the first sign of irregularity up to 3 times a day
- put one dose into an empty glass
- fill glass with 8oz of water or favorite beverage
- stir briskly and drink promptly
- if mixture thickens add more water and stir
- take this product (child or adult dose) with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning.
- if you are taking a prescription medicine by mouth, take this product at least 2 hours before or 2 hours after the prescribed medicine.

New Users: Start with 1 dose per day; gradually increase to 3 doses per day as necessary

adults and children 12 years and over	1 rounded tablespoon
6 to under 12 years	1/2 tablespoon
under 6 years	ask a doctor

Other information

- each tablespoon contains: potassium 30 mg and sodium 5 mg
- store at (20°-25°C) 68°-77°F

Inactive ingredients

citric acid, FD and C yellow 6, natural and artificial flavor, silicon dioxide, sucrose

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR BROKEN

Fiber Supplement information

Uses Diets low in saturated fat and cholesterol that include 7 grams of soluble fiber per day from psyllium seed husk, may reduce the risk of heart disease by lowering cholesterol levels. One ADULT serving provides 2.4 grams of this soluble fiber. Consult a doctor if you are considering use of the product as part of a cholesterol-lowering program.

Warnings Read entire label, including warnings in Drug Facts

Directions Adults 12 years and older: 1 rounded TABLESPOON (tbsp) in 8 oz of liquid 3 times daily.

Children under 12 years: Consult a doctor.

NOTICE: Mix this product with at least 8 ounces (a full glass) of liquid. Taking this product without enough liquid may cause choking. Do not take if you have difficulty swallowing. **See choking warning.**

PDP


Sunmark

compare to the active ingredient in Metamucil®

fiber

Smooth Texture
 Bulk-forming laxative
 Dietary fiber supplement
 100% Natural Psyllium Seed Husk Fiber
 Orange Flavor
 Naturally and Artificially Flavored

48 Tablespoon doses
 Net wt 20.3 oz (575 g) (1 LB 4.3 OZ)
 FILL CONTROLLED BY WEIGHT, NOT VOLUME



FIBER LAXATIVE

Drug Facts

Active ingredient (per each rounded tablespoon = 12 g)	Purpose
Psyllium husk 3.4 g	Bulk-forming laxative

Uses

- effective in treating occasional constipation and restoring regularity

Warnings

Allergy alert: May cause allergic reaction in people sensitive to ingested or inhaled psyllium

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have

- abdominal pain, nausea or vomiting
- a sudden change in bowel habits persisting for 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Laxatives may affect how other drugs work.

When using this product

- do not exceed the maximum recommended daily dosage in a 24-hour period
- you may experience changes in bowel habits or minor bloating as your body adjusts to increased fiber intake

Stop use and ask a doctor if

- constipation lasts more than 7 days
- rectal bleeding occurs

These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

READ ENTIRE LABEL INCLUDING WARNINGS IN DRUG FACTS.

◀ NOTE SEPARATE USES ▶

Drug Facts (continued)

Directions

- take at the first sign of irregularity up to 3 times a day
 - put one dose into an empty glass
 - fill glass with 8oz of water or favorite beverage
 - stir briskly and drink promptly
 - if mixture thickens add more water and stir
 - take this product (child or adult dose) with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning.
 - if you are taking a prescription medicine by mouth, take this product at least 2 hours before or 2 hours after the prescribed medicine.
 - New Users:** Start with 1 dose per day; gradually increase to 3 doses per day as necessary
- | adults and children 12 years and over | 1 rounded tablespoon |
|---------------------------------------|----------------------|
| 6 to under 12 years | 1/2 tablespoon |
| under 6 years | ask a doctor |

Other information

- each tablespoon contains: potassium 30 mg and sodium 5 mg
 - store at (20°-25°C) 68°-77°F
- Inactive ingredients** citric acid, FD&C yellow #6, natural and artificial orange flavor, silicon dioxide, sucrose.

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR BROKEN

**This product is not manufactured or distributed by Procter & Gamble, the distributor of METAMUCIL®.



Another Quality Product
 Distributed By McKesson
 One Post Street, San Francisco, CA 94104
 Money Back Guarantee
 Please visit us at www.sunmarkbrand.com

DIETARY FIBER SUPPLEMENT

Fiber Supplement Information

Uses Diets low in saturated fat and cholesterol that include 7 grams of soluble fiber per day from psyllium seed husk, may reduce the risk of heart disease by lowering cholesterol levels. One ADULT serving provides 2.4 grams of this soluble fiber. Consult a doctor if you are considering use of the product as part of a cholesterol-lowering program.

Warnings Read entire label, including warnings in Drug Facts.

Directions Adults 12 years & older: 1 rounded TABLESPOON (Tbsp) in 8 oz of liquid 3 times daily. Children under 12 years: Consult a doctor.

NOTICE: Mix this product with at least 8 ounces (a full glass) of liquid. Taking this product without enough liquid may cause choking. Do not take if you have difficulty swallowing. See choking warning.

Supplement Facts

Serving Size: 1 rounded Tbsp. (12 g)
 Servings Per Container: 48

Amount Per Serving	% DV*
Calories 45	
Total Carb. 12 g	4%*
Dietary Fiber 3 g	12%*
Soluble Fiber 2 g	†
Sugars 9 g	†
Iron 1 mg	4%
Sodium 5 mg	<1%
Potassium 30 mg	<1%

*Percent Daily Values (DV) are based on a 2,000 calorie diet.
 † Daily Value not established.

Ingredients: Sucrose, Psyllium Husk, Citric Acid, Natural and Artificial Orange Flavor, FD&C Yellow #6, Silicon Dioxide.

LOT:

EXP:

SMOOTH TEXTURE ORANGE FLAVOR

psyllium husk powder, for suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49348-091
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PSYLLIUM HUSK (UNII: 0SHO53407G) (PSYLLIUM HUSK - UNII:0SHO53407G)	PSYLLIUM HUSK	3.4 g in 12 g

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SUCROSE (UNII: C151H8M554)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics

Color	brown	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49348-091-68	575 g in 1 CANISTER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	04/05/2007	

Labeler - McKesson (177667227)

Revised: 9/2012

McKesson