

DG HEALTH IBUPROFEN- ibuprofen tablet, film coated
Dolgencorp Inc

Dolgencorp, LLC Ibuprofen Drug Facts

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

Adults and children 12 years and older:

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

Children under 12 years: ask a doctor

Other information

- read all warnings and directions before use
- store between 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- see end panel for lot number and expiration date

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, iron oxides, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, stearic acid, talc, titanium dioxide

Questions or comments?

1-888-309-9030

Principal Display Panel

DG™ | health

SEE REVISED WARNING

Compare to the active ingredient of Motrin® IB

Ibuprofen Tablets, 200 mg

Pain Reliever / Fever Reducer (NSAID)


50 Caplets**

**Capsule-Shaped Tablets

200 mg

Actual Caplet Size

51774 VT C3



A1491

Drug Facts

Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)..... Pain reliever/fever reducer
Nonsteroidal anti-inflammatory drug

Purposes

Uses ■ Temporarily relieves minor aches and pains due to:
■ headache ■ muscle aches ■ minor pain of arthritis ■ toothache ■ backache
■ the common cold ■ menstrual cramps ■ Temporarily reduces fever

Warnings

Warning: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or aspirin drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ If you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ Right before or after heart surgery
Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ The stomach bleeding warning applies to you ■ you have a history of stomach problems, such as

Drug Facts (continued)

asthma ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or a bad stroke ■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug
When using this product ■ Take with food or milk if stomach upset occurs.

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not go away ■ you have symptoms of heart problems or stroke ■ chest pain ■ trouble breathing ■ weakness in one part of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor or because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ Do not take more than directed ■ The smallest effective dose should be used. Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ If pain or fever does not respond to 1 caplet, 2 caplets may be used ■ Do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ask a doctor

100% Satisfaction Guaranteed!
(888) 309-9030

Important: Read all product information before using.
Keep this box for important information.
Questions or comments? 1-888-309-9030
Inactive ingredients include: silicon dioxide, corn starch, croscarmellose sodium, FD&C red no. 40 aluminum lake, FD&C yellow no. 5 aluminum lake, iron oxide, microcrystalline cellulose, polyethylene glycol, diphenyl dicarbonyl, stearic acid, talc, titanium dioxide, and glycerin.
Other information ■ Store between 20° to 25°C (68° to 77°F) ■ Avoid high humidity and moisture ■ See end panel for directions before use and expiration date



DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.
***This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Motrin® B.

SEE REVISED WARNING
Compare to the active ingredient of Motrin® IB***

DG™ health
Ibuprofen
Tablets, 200 mg
Pain Reliever/
Fever Reducer
(NSAID)



50 Caplets**

200 mg

Actual Caplet Size

**Capsule-Shaped Tablets

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-758
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	I2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-758-71	1 in 1 CARTON	05/03/2022	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55910-758-78	1 in 1 CARTON	05/03/2022	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:55910-758-85	250 in 1 BOTTLE; Type 0: Not a Combination Product	05/03/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077349	05/03/2022	

Labeler - Dolgencorp Inc (068331990)

Revised: 4/2024

Dolgencorp Inc