

MEDIQUE AT HOME IPRIN- ibuprofen tablet, film coated
Unifirst First Aid Corporation

Medique @ Home Iprin Tablets

Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains associated with:

- headache ■ toothache ■ backache ■ menstrual cramps
- common cold ■ muscular aches ■ minor arthritis pain

temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ skin reddening ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning : This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever

reducer

- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not use more than directed**

■ **the smallest effective dose should be used**

■ do not take longer than 10 days, unless directed by a doctor (see Warnings)

Adults and children: (12 years and older)

■ take 1 tablet every 4 to 6 hours while symptoms persist

■ if pain or fever does not respond to 1 tablet, 2 tablets may be used

■ do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years:

■ ask a doctor

Other information

■ store at 68-77°F (20-25°C)

■ avoid excessive heat 104°F (above 40°C)

■ **TAMPER EVIDENT** child resistant packets

■ do not use any opened or torn packets

Inactive ingredients

carnauba wax*, corn starch, hypromellose*, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose*, polydextrose*, polyethylene glycol, polyvinyl alcohol*, povidone k-30*, silicon dioxide, sodium starch glycolate, stearic acid, talc*, titanium dioxide

*may contain

Questions or comments? 1-800-634-7680

Medique® @ Home

I-Prin Tablets

Ibuprofen 200 mg

Pain Reliever/Fever Reducer

(NSAID)

Pull to Open

Child-Resistant Packets

*not actual size

50 Packets

(2 Tablets per Packet)

Compare to the active ingredient in Advil® Tablets

Medique® @ Home
I-Prin Tablets
Ibuprofen 200 mg
Pain Reliever/Fever Reducer
(NSAID)

Child-Resistant Packets
*not actual size
Compare to the active ingredient in Advil® Tablets*

Medique® @ Home
I-Prin Tablets
Ibuprofen 200 mg
Pain Reliever/Fever Reducer
(NSAID)

Child-Resistant Packets
*not actual size
Compare to the active ingredient in Advil® Tablets*

Retain carton for complete product information

The Brands That Work™ Now Work At Home

100 Tablets 50 Packets (2 Tablets Per Packet) 50 Packets (2 Tablets Per Packet)

Compare to the active ingredient in Advil® Tablets* Compare to the active ingredient in Advil® Tablets*

1Advil® is a registered trademark of Pfizer Consumer and is not affiliated with Medique Products or this product.

Reorder 70833
Manufactured for
Medique Products
Fort Myers, FL 33967
1-800-634-7680
www.mediqueproducts.com

Medique® @ Home

I-Prin Tablets

Ibuprofen 200 mg

Pain Reliever/Fever Reducer

(NSAID)

Pull to Open

Child-Resistant Packets

*not actual size

50 Packets

(2 Tablets per Packet)

Compare to the active ingredient in Advil® Tablets



Medique Home
I-Prin Tablets
Ibuprofen 200 mg
Pain Reliever/Fever Reducer
(NSAID)

Child-Resistant Packets
*not actual size

Compare to the active ingredient in Advil® tablets



Medique Home
I-Prin Tablets
Ibuprofen 200 mg
Pain Reliever/Fever Reducer
(NSAID)

Child-Resistant Packets
*not actual size

The Brands That Work™ Now Work At Home

100 Tablets 50 Packets (2 Tablets Per Packet)

Compare to the active ingredient in Advil® Tablets*



Medique Home
I-Prin Tablets
Ibuprofen 200 mg
Pain Reliever/Fever Reducer
(NSAID)

Child-Resistant Packets
*not actual size

50 Packets (2 Tablets Per Packet)

Compare to the active ingredient in Advil® tablets*

Drug Facts

Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses
Temporarily relieves minor aches and pains associated with:

- headache
- toothache
- backache
- menstrual cramps
- common cold
- rheumatic aches
- minor arthritis pain

 Sore throats reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- skin redness
- swelling (swelling)
- facial swelling
- rash
- shock
- stomach

 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant or blood) drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Drug Facts (continued)

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- stay up and ask a doctor if
- you experience any of the following signs of stomach bleeding:
 - red, dark, or vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - numbness or tingling
 - swelling
 - sting
 - pain gets worse or lasts for more than 10 days
 - fever gets worse or lasts for more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 30 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222) (continued on back panel)

Drug Facts (continued)

Directions

- do not use more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

Adults and children (12 years and older)

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

Children under 12 years

- ask a doctor

Other information

- store at 68°F to 77°F (20° to 25°C)
- avoid excessive heat (104°F above 40°C)
- IMPROPER EXCESSIVE CHILD-RESISTANT PACKETS
- do not use any opened or torn packets

Inactive ingredients
 carnauba wax*, corn starch, hydroxypropylcellulose, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose, polydextrose, polyethylene glycol, polyvinyl alcohol*, potassium hydroxide, silicon dioxide, sodium starch glycolate, stearic acid, titanium dioxide
 *may contain

Questions or comments? 1-800-634-7680

1Advil® is a registered trademark of Pfizer Consumer and is not affiliated with Medique Products or this product.

Reader 70833
 Manufactured for
 Medique Products
 Fort Myers, FL 33967
 1-800-634-7680
 www.mediqueproducts.com

MEDIQUE AT HOME IPRIN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-683
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z65AP)	
ANHYDROUS LACTOSE (UNII: 3S5Y5LH9PMK)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDWL1A)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

STARCH, CORN (UNII: O8232NY3SJ)

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;291
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-683-33	50 in 1 BOX	02/01/2021	
1	NDC:47682-683-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	02/01/2021	

MEDIQUE AT HOME IPRIN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-783
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE K30 (UNII: U725QWY32X)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	

SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	brown (Reddish Brown)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	G;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-783-33	50 in 1 BOX	02/01/2021	
1	NDC:47682-783-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	02/01/2021	

Labeler - Unifirst First Aid Corporation (832947092)

Revised: 9/2025

Unifirst First Aid Corporation