

MEDIQUE AT HOME MEDIPROXEN- naproxen sodium tablet, film coated
Unifirst First Aid Corporation

Medique @ Home Mediproxen

Drug Facts

Active ingredient (in each tablet)

Naproxen Sodium USP, 220 mg

(naproxen 200 mg) (NSAID*)

* nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

■ temporarily relieves minor aches and pains due to: ■ backache ■ muscular aches ■ minor pain of arthritis ■ menstrual cramps

■ headache ■ toothache ■ the common cold

temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin,

ibuprofen,
naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAID, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems such, as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children: (12 years and older)

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8 to 12 hour period
- do not exceed 3 tablets in a 24-hour period

Children under 12 years: ■ ask a doctor

Other information

- each tablet contains: sodium 20 mg
- store at 68-77°F (20-25°C)
- avoid high humidity and excessive heat above 104°F (40°C)
- **Tamper Evident** child resistant packets
- do not use any open or torn packets

Inactive ingredients

croscarmellose sodium, FD&C blue #2, macrogol, magnesium stearate, polyvinyl alcohol, povidone, pregelatinized starch, talc, titanium dioxide

Questions or comments? 1-800-634-7680

Medique® @ Home

Mediproxen

Naproxen Sodium 220 mg

Pain Reliever/Fever Reducer

(NSAID)

Pull to Open

Child-Resistant Packets

*not actual size

50 Packets

(1 Tablet Per Packet)

Compare to the active ingredient in Aleve® Tablets

Medique Home Mediproxen
Naproxen Sodium 220 mg Pain Reliever/Fever Reducer (NSAID)

Medique Home Mediproxen
Naproxen Sodium 220 mg Pain Reliever/Fever Reducer (NSAID)

Drug Facts
Active ingredient (in each tablet) Purpose
Naproxen Sodium USP 220 mg... Pain reliever/fever reducer (Naproxen 220 mg NSAID)*
Nonsteroidal anti-inflammatory drug

Directions
Do not take more than directed
The smallest effective dose should be used
Drink a full glass of water with each dose

Other Information
Each tablet contains: sodium 20mg
Store at 68-77°F (20-25°C)
Keep away from moisture and excessive heat above 104°F (40°C)
TAMPER EVIDENT child resistant packets
Do not use any opened or torn packets

Inactive ingredients
croscarmellose sodium, FD&C blue #2, magnesium, magnesium stearate, polyvinyl alcohol, povidone, polyethylene glycol, titanium dioxide

Questions or comments? 1-800-634-7680

3 47682 73750 4

Reorder 73750
Manufactured for
Medique Products
Fort Myers, FL 33967
1-800-634-7680
www.mediqueproducts.com

MEDIQUE AT HOME MEDIPROXEN			
naproxen sodium tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-737
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)		NAPROXEN SODIUM	220 mg
Inactive Ingredients			
Ingredient Name		Strength	
STARCH, CORN (UNII: O8232NY3SJ)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDWL4)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)			
TALC (UNII: 7SEV7J4R1U)			
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)			

POVIDONE (UNII: FZ989GH94E)

CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)

Product Characteristics

Color	blue	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	I;3
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-737-50	50 in 1 BOX	02/01/2021	
1	NDC:47682-737-46	1 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079096	02/01/2021	

Labeler - Unifirst First Aid Corporation (832947092)

Establishment

Name	Address	ID/FEI	Business Operations
Medique Products		086911794	pack(47682-737)

Revised: 9/2025

Unifirst First Aid Corporation