

**DOCUSATE SODIUM - STOOL SOFTENER LAXATIVE- docusate sodium capsule,  
liquid filled  
Cardinal Health 107, LLC**

-----  
**Docusate Sodium - Stool Softener Laxative**

***Drug Facts***

***Active ingredient (in each softgel)***

Docusate Sodium 100 mg

***Purpose***

Stool softener laxative

***Uses***

- for relief of occasional constipation (irregularity)
- This generally produces a bowel movement within 12 to 72 hours.

***Warnings***

**Ask a doctor before use if you have**

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

**Ask a doctor or pharmacist before use if you are** presently taking mineral oil

**Stop use and ask a doctor if**

- you have rectal bleeding or no bowel movement after using this product.

These can be signs of a serious condition

- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,** ask a health care professional before use.

**Keep out of reach of children**

**Overdose Warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

***Directions***

- take softgels with a full glass (8 oz) of water

---

adults and children 12 years of age and	take 1 to 3 softgels daily or as directed
--	--

older	by a doctor. This dose may be taken as a single daily dose or in divided doses.
children 6 to under 12 years of age	take 1 softgel daily or as directed by a doctor
children under 6 years of age	ask a doctor

### ***Other information***

- **each softgel contains:** sodium 10 mg (very low sodium)
- store at 25°C (77°F) in a dry place. Avoid excessive heat 40°C (140°F).

Overbagged with 10 liquid-filled capsules per bag, NDC 55154-3369-0

WARNING: This Unit Dose package is not child resistant and is Intended for Institutional Use Only. Keep this and all drugs out of the reach of children.

### ***Inactive ingredients***

edible white ink, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, isopropyl alcohol, light mineral oil, medium chain triglycerides, polyethylene glycol, propylene glycol, purified water, sorbitol sorbitan solution

### ***Questions or comments?***

**(800)616-2471**

**MAJOR®**

**Unit Dose**

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

**READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

Distributed by:

**MAJOR® PHARMACEUTICALS**

Indianapolis, IN 46268

**Distributed By:**

**Cardinal Health**

Dublin, OH 43017

L57466800124

Product of UAE

Packaged and Quality Assured in the USA

CT0904718361

REV.01-032022

## Principal Display Panel

NDC 55154-3369-0

### DOCUSATE SODIUM 100 mg

Stool Softener Laxative

10 SOFTGELS



NDC 55154-3369-0

# R122

## DOCUSATE SODIUM 100 mg Stool Softener Laxative

### 10 SOFTGELS

Each softgel contains: sodium 10 mg (very low sodium)

#### Drug Facts

Active ingredient (in each softgel)

Docosate Sodium 100 mg

Purpose

Stool softener laxative

Inactive ingredients edible white ink, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, isopropyl alcohol, light mineral oil, medium chain triglycerides, polyethylene glycol, propylene glycol, purified water, sorbitol sorbitan solution

Uses -for relief of occasional constipation (irregularity)

-This generally produces a bowel movement within 12 to 72 hours

Directions - take softgels with a full glass (8 oz) of water

adults and children 12 years of age and older	take 1 to 3 softgels daily or as directed by a doctor. This dose may be taken as a single daily dose or in divided doses.
---	---

children 6 to under 12 years of age	take 1 softgel daily or as directed by a doctor
-------------------------------------	---

children under 6 years of age	ask a doctor
-------------------------------	--------------

**STORAGE:** Store at 25° C (77° F); in a dry place.

Avoid excessive heat 40° C (140° F).

#### Warnings

Ask a doctor before use if you have -abdominal pain, nausea or vomiting

-a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are presently taking mineral oil

Stop use and ask a doctor if -you have rectal bleeding or no bowel

movement after using this product .These can be signs of a serious condition.

-you need to use a laxative for more than 1 week

**WARNING:** This Unit Dose package is not child resistant and is Intended for

Institutional Use Only. If pregnant or breast-feeding, ask a health care

professional before use. Keep out of reach of children.

**Overdose Warning:** In case of overdose, get medical help or contact a

Poison Control Center right away (1-800-222-1222). Prompt medical

attention is critical for adults as well as children even if you do not

notice any signs or symptoms.

**TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR  
SHOWS ANY SIGNS OF TAMPERING**

**READ FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

Questions or comments? (800) 616-2471

Distributed by MAJOR® PHARMACEUTICALS

Indianapolis, IN 46268

Product of UAE

Packaged and Quality Assured in the USA

Distributed by Cardinal Health

Dublin, OH 43017

L57466800124

Lot:           Exp:

# DOCUSATE SODIUM - STOOL SOFTENER LAXATIVE

docusate sodium capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55154-3369(NDC:0904-7183)
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)	DOCUSATE SODIUM	100 mg

## Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O921CV9RU)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	777
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-3369-0	10 in 1 BAG	09/04/2021	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M007	09/04/2021	

**Labeler** - Cardinal Health 107, LLC (118546603)

Revised: 10/2025

Cardinal Health 107, LLC