

WALGREENS ANTACID PAIN RELIEF EXTRA STRENGTH- aspirin, citric acid, sodium bicarbonate tablet, effervescent
Walgreens

Walgreens Antacid and Pain Relief Extra Strength Effervescent Tablets

Active ingredients

(in each effervescent tablet)

Anhydrous citric acid 1000mg

Aspirin 500 mg **(NSAID)***

Sodium bicarbonate (heat-treated) 1985 mg

*Nonsteroidal anti-inflammatory drug

Purpose

Anhydrous citric acid 1000mg.....Antacid

Aspirin 500 mg **(NSAID)***.....Pain Reliever

Sodium bicarbonate (heat-treated) 1985 mg.....Antacid

*Nonsteroidal anti-inflammatory drug

Uses

Uses for the temporary relief of

■ heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains

■ upset stomach with headache from overindulgence in food or drink

■ headache, body aches and pain alone

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin,

ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer
■ if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma ■ you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

■ presently taking a prescription drug. Antacids may interact with certain prescription drugs.
■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product

When using this product do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician.

Stop use and ask a doctor if

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding

- feel faint ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use. **It is especially**

important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Directions

■ do not exceed recommended dosage

■ to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking
adults and children 12 years and over (up to 60 years of age): 2 tablets every 4 hours
do not exceed 8 tablets in 24 hours

60 years and over: 2 tablets every 4 hours do not exceed 4 tablets in 24 hours

Children under 12 years: consult a doctor

Take Walgreens Antacid + Pain Relief any time - morning, noon, or night - when you need relief from heartburn, acid indigestion, sour stomach with headache or body aches and pains.

Place two (2) tablets in 4 oz. of water. Dissolve tablets completely.

You do not have to drink any of the residue that may be at the bottom of the glass. The medicines are in the water.

Other information

■ **each tablet contains:** sodium 575 mg

■ store at room temperature (59°-86°F)

Inactive ingredients:

flavor

Questions or comments?

1-888-228-6937

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Antacid + Pain Relief



Walgreens

Drug Facts (continued)

such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma ■ you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are ■ presently taking a prescription drug, Antacids may interact with certain prescription drugs. ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician.

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ an allergic reaction occurs. Seek medical help right away. ■ pain lasts for more than 10 days or gets worse ■ new symptoms occur ■ redness or swelling is present ■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact the Poison Control Center right away at 1-800-222-1222.

Directions

■ do not exceed recommended dosage ■ to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking

adults and children 12 years and over 2 tablets every 4 hours do not exceed 8 tablets (up to 60 years of age)

60 years and over 2 tablets every 4 hours do not exceed 4 tablets in 24 hours

Children under 12 years consult a doctor

Other information

■ each tablet contains: sodium 576 mg ■ store at room temperature (59°-86°F)

Inactive ingredients: flavor

Questions or comments? 1-888-228-6937

Take Walgreens Antacid + Pain Relief any time – morning, noon or night – when you need relief from heartburn, acid indigestion or sour stomach with headache or body aches and pains.

Mornings



Midday



Evenings



Directions for use:

Place two (2) tablets in 4 oz. of water. Dissolve tablets completely.



You do not have to drink any of the residue that may be on the bottom of the glass. The medicines are in the water.

Walgreens

Compare to the active ingredient in Alka-Seltzer® Extra Strength®



NDC 0363-0541-36

Antacid + Pain Relief

ANHYDROUS CITRIC ACID / ANTACID
ASPIRIN (NSAID) / PAIN RELIEVER
SODIUM BICARBONATE / ANTACID

Extra Strength

Relieves heartburn, acid indigestion, upset stomach with headache, body aches & pains

36 EFFERVESCENT TABLETS



ACTUAL SIZE

Drug Facts

Active ingredients (in each effervescent tablet)
Anhydrous citric acid 1000 mg Antacid
Aspirin 500 mg (NSAID) Pain Reliever
Sodium bicarbonate (heat-treated) 1985 mg Antacid
*Nonsteroidal anti-inflammatory drug

Uses for the temporary relief of ■ heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains ■ upset stomach with headache from overindulgence in food or drink ■ headache, body aches, and pain alone

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems.

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Alka-Seltzer®.



DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.

ITEM 319567 W10108-0322-L
3 11917 00273 6

51068 WA C2

PN: 51068 WA C2
R: 07/21
W30R1021-F

WALGREENS ANTACID PAIN RELIEF EXTRA STRENGTH

aspirin, citric acid, sodium bicarbonate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0541
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	ANHYDROUS CITRIC ACID	1000 mg
SODIUM BICARBONATE (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1Z RA3Q20)	SODIUM BICARBONATE	1985 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg

Inactive Ingredients

Ingredient Name	Strength
LEMN (UNII: 24RS0A988O)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor		Imprint Code	PXS
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0541-36	18 in 1 CARTON	02/05/2021	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/05/2021	12/31/2027

Labeler - Walgreens (008965063)

Revised: 10/2025

Walgreens