

PAIN RELIEVER EXTRA STRENGTH- acetaminophen liquid
Rite Aid Corporation

Drug Facts

Active ingredient (in each 15 mL)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- do not take more than 4 doses (120 mL) in any 24 hours period
- measure only with dosing cup provided. Do not use any other dosing device
- mL=milliliter
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL every 6 hours while symptoms last
 - do not take more than 10 days unless directed by a doctor
- children under 12 years: do not use

Other information

- **each 15 mL contains:**sodium 6 mg
- store between 20-25°C (68-77°F) do not refrigerate
- protect from light

Inactive ingredients

citric acid, D&C red 33, FD&C red 40, flavors, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol

Questions or comments?

Call 1-877753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in Tylenol® Extra Strength*

ADULT • EXTRA STRENGTH

PAIN RELIEVER

ACETAMINOPHEN 500 mg per 15 mL

PAIN RELIEVER/FEVER REDUCER

Rapid burst liquid

For ages 12 years & over

Alcohol free

CHERRY FLAVOR

FL OZ (mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Extra Strength.

MANUFACTURED BY: RITE AID, 30 HUNTER LANE,

CAMP HILL, PA 17011

www.riteaid.com

Product Label



TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011 www.riteaid.com

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.



PEEL CORNER FOR DRUG FACTS

Drug Facts

Active ingredient (in each 15 mL) Acetaminophen 500 mg.....Pain reliever/fever reducer

Purpose

Uses

- temporarily relieves minor aches and pains due to
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
 - temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen.

Drug Facts (continued)

Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Drug Facts (continued)

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses (120 mL) in any 24-hour period

Drug Facts (continued)

- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL every 6 hours while symptoms last
 - do not take more than 10 days unless directed by a doctor
- children under 12 years: do not use

Other information

- each 15 mL contains: sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.
- protect from light

Drug Facts (continued)

Inactive ingredients citric acid, D&C red #33, FD&C red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Extra Strength.

RITE AID PHARMACY Adult Extra Strength Pain Reliever Cherry Flavor

PAIN RELIEVER EXTRA STRENGTH

acetaminophen liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-6440
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDWLA)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-6440-1	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/31/2020	05/29/2026

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/31/2020	05/29/2026

Labeler - Rite Aid Corporation (014578892)

Revised: 8/2025

Rite Aid Corporation