

**PAIN RELIEVER EXTRA STRENGTH- acetaminophen liquid**  
**Rite Aid Corporation**

-----  
**Drug Facts**

**Active ingredient (in each 15 mL)**

**Acetaminophen 500 mg**

**Purpose**

**Pain reliever/fever reducer**

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks ever day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

## **Ask a doctor before use if you have**

liver disease.

## **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

## **Stop use and ask a doctor if**

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

- do not take more than directed (see overdose warning)
- do not take more than 4 doses (120 mL) in any 24 hours period
- measure only with dosing cup provided. Do not use any other dosing device
- mL=milliliter
- keep dosing cup with product
- adults and children 12 years and over
  - 30 mL every 6 hours while symptoms last
  - do not take more than 10 days unless directed by a doctor
- children under 12 years: do not use

## **Other information**

- **each 15 mL contains:**sodium 6 mg
- store between 20-25°C (68-77°F) do not refrigerate
- protect from light

## **Inactive ingredients**

citric acid, D&C red 33, FD&C red 40, flavors, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol

## Questions or comments?

Call 1-877753-3935 Monday-Friday 9AM-5PM EST

## Principal Display Panel

Compare to the active ingredient in Tylenol® Extra Strength\*

ADULT • EXTRA STRENGTH

PAIN RELIEVER

ACETAMINOPHEN 500 mg per 15 mL

PAIN RELIEVER/FEVER REDUCER

Rapid burst liquid

For ages 12 years & over

Alcohol free

CHERRY FLAVOR

FL OZ (mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Extra Strength.

MANUFACTURED BY: RITE AID, 30 HUNTER LANE,

CAMP HILL, PA 17011

[www.riteaid.com](http://www.riteaid.com)

## Product Label



**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

**DISTRIBUTED BY: RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011 [www.riteaid.com](http://www.riteaid.com)**

**SATISFACTION GUARANTEE:** If you're not satisfied, we'll happily refund your money.



PEEL CORNER FOR DRUG FACTS

**Drug Facts**

**Active ingredient (in each 15 mL)**  
Acetaminophen 500 mg.....Pain reliever/fever reducer

**Purpose**  
Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
  - temporarily reduces fever

**Warnings**  
Liver warning: This product contains acetaminophen.

**Drug Facts (continued)**

Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Drug Facts (continued)**

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease.**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Drug Facts (continued)**

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses (120 mL) in any 24-hour period

**Drug Facts (continued)**

- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over
  - 30 mL every 6 hours while symptoms last
  - do not take more than 10 days unless directed by a doctor
- children under 12 years: do not use

**Other information**

- each 15 mL contains: sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.
- protect from light

**Drug Facts (continued)**

**Inactive ingredients** citric acid, D&C red #33, FD&C red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol

**Questions or comments?**  
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Extra Strength.

**RITE AID PHARMACY Adult Extra Strength Pain Reliever Cherry Flavor**

PAIN RELIEVER EXTRA STRENGTH			
acetaminophen liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-6440
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	500 mg in 15 mL
Inactive Ingredients			
Ingredient Name			Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
D&C RED NO. 33 (UNII: 9DBA0SBB0L)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SORBITOL (UNII: 506T60A25R)			

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-6440-1	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/31/2020	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/31/2020	

**Labeler** - Rite Aid Corporation (014578892)

Revised: 5/2024

Rite Aid Corporation