COPPERTONE GLOW SUNCREEN SPF 30- avobenzone, homosalate, octisalate, octocrylene aerosol, spray Beiersdord Inc
Coppertone Glow Sunscreen Spray SPF 30
Drug Facts
Drug Facts
Active Ingredients
Active ingredients
(To Deliver) Avobenzone 3%, Homosalate 9%, Octisalate 4.5%, Octocrylene 8%
Purpose
Purpose
Sunscreen
Use

■ use only as directed. Intentional misuse by deliberately concentrating and inhaling the

do not puncture or incinerate. Contents under pressure. Do not store at

Use

Warnings

Warnings

helps prevent sunburn

For external use only

When using this product

contents can be harmful or fatal.

temperatures above 120°F.

Flammable: Do not use near heat, flame, or while smoking

Do not use on damaged or broken skin

■ keep away from face to avoid breathing it

Stop use and ask a doctor if rash occurs

■ keep out of eyes. Rinse with water to remove.

Keep out of reach of children.If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

Directions

- shake well before each use
- spray liberally and spread evenly by hand 15 minutes before sun exposure
- hold container 4 to 6 inches from the skin to apply
- do not spray directly into face. Spray on hands then apply to face.
- do not apply in windy conditions
- use in a well-ventilated area
- reapply:
- after 80 minutes of swimming or sweating
- immediately after towel drying
- at least every 2 hours
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
- limit time in the sun, especially from 10 a.m. 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor

Other Information

Other information

- protect this product from excessive heat and direct sun
- store between 20° to 25°C (68° to 77°F)
- may stain or damage some fabrics or surfaces

Inactive Ingredients

Inactive ingredients water, isobutane, SD alcohol 40-B (12.6% v/v), diethylhexyl carbonate, butyloctyl salicylate, diisopropyl adipate, glycereth-26, butylene glycol, dimethicone, polyamide-8, isododecane, styrene/acrylates copolymer, 1,2-hexanediol, ammonium acryloyldimethyltaurate/beheneth-25 methacrylate crosspolymer, glycerin, hydroxyacetophenone, mica, synthetic fluorphlogopite, tocopherol, ethylhexyl methoxycrylene, arachidyl alcohol, iron oxides, bismuth oxychloride, fragrance, xanthan gum, behenyl alcohol, silica, arachidyl glucoside, disodium EDTA, sodium ascorbyl

phosphate

Questions

Questions?1-866-288-3330

Principal Display Panel

Coppertone®

Sunscreen Lotion Spray

Glow

Lightweight Spray

With Shimmer

SPF 30

Free of oxybenzone, octinoxate, parabens, PABA & phthalates

Water Resistant (80 minutes)

Broad Spectrum SPF 30

NET WT 5 OZ (142 g)





Coppertone® Glow Sunscreen Spray helps protect skin against the sun's most damaging UVA/UVB rays. This nourishing spray is ultra-hydrating, dries in seconds, and helps leave skin feeling soft & smooth with a beautiful, glowing shimmer.

Drug Facts Active ingredients (To Deliver) Avobenzone 3%, Homosalate 9%, Octisalate 4.5%, Octocrylene 8% Purpose **Use** ■ helps prevent sunburn

Warnings

For external use only

Flammable: Do not use near heat, flame, or while smoking

Do not use on damaged or broken skin

When using this product


- keep away from face to avoid breathing itkeep out of eyes. Rinse with water to remove
- use only as directed. Intentional misuse by deliberately concentrating and inhaling the contents can be
- do not puncture or incinerate. Contents under pressure. Do not store at temperatures above 120°F.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

- shake well before each use
- spray liberally and spread evenly by hand 15 minutes before sun exposure
- hold container 4 to 6 inches from the skin to apply
- do not spray directly into face. Spray on hands then apply to face.
 do not apply in windy conditions
- use in a well-ventilated area
- reapply:
- after 80 minutes of swimming or sweating
- immediately after towel drying at least every 2 hours
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
- limit time in the sun, especially from 10 a.m. 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor



Beiersdorf

Beiersdorf Inc. Wilton, CT 06897 ®=reg. TM. of Beiersdorf AG, Germany

COPPERTONE GLOW SUNCREEN SPF 30

avobenzone, homosalate, octisalate, octocrylene aerosol, spray

Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:66800-0003 **Route of Administration TOPICAL**

7

Active Ingredient/Active Moiety			
Ingredient Name Basis o		Strength	
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	80 mg in 1 g	
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	30 mg in 1 g	

l	HOMOSALATE (UNII: V06SV4M95S) (HOMOSALATE - UNII: V06SV4M95S)	HOMOSALATE	90 mg in 1 g
l	OCTISALATE (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	45 mg in 1 g

Inactive Ingredients		
Ingredient Name	Strength	
WATER (UNII: 059QF0KO0R)		
SODIUM ASCORBYL PHOSPHATE (UNII: 836SJG51DR)		
ISOBUTANE (UNII: BXR49TP611)		
GLYCERETH-26 (UNII: NNE56F2N14)		
DOCOSANOL (UNII: 9G10E216XY)		
FRAGRANCE FRESH CITRUS FLORAL ORC1501495 (UNII: OU4GI2R2WB)		
STYRENE/ACRYLAMIDE COPOLYMER (MW 500000) (UNII: 5Z4DPO246A)		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
AMMONIUM ACRYLOYLDIMETHYLTAURATE/BEHENETH-25 METHACRYLATE CROSSPOLYMER (52000 MPA.S) (UNII: LZ291VH90H)		
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)		
POLYAMIDE-8 (4500 MW) (UNII: 77723GV81A)		
BUTYLOCTYL SALICYLATE (UNII: 2EH13UN8D3)		
FERRIC OXIDE YELLOW (UNII: EX43802MRT)		
DIETHYLHEXYL CARBONATE (UNII: YCD5000Z6L)		
MAGNESIUM POTASSIUM ALUMINOSILICATE FLUORIDE (UNII: YK3DC63Y5M)		
TOCOPHEROL (UNII: R0ZB2556P8)		
ETHYLHEXYL METHOXYCRYLENE (UNII: S3KFG6Q5X8)		
DIISOPROPYL ADIPATE (UNII: P7E6YFV72X)		
ISODODECANE (UNII: A8289P68Y2)		
GLYCERIN (UNII: PDC6A3C0OX)		
DIMETHICONE (UNII: 92RU3N3Y1O)		
COBALT DISODIUM EDETATE (UNII: 3EY1Y2QRLI)		
HYDROXYACETOPHENONE (UNII: G1L3HT4CMH)		
ARACHIDYL GLUCOSIDE (UNII: 6JVW35JOOJ)		
ARACHIDYL ALCOHOL (UNII: 1QR1QRA9BU)		
MICA (UNII: V8A1AW0880)		
XANTHAN GUM (UNII: TTV12P4NEE)		
ALCOHOL (UNII: 3K9958V90M)		
BISMUTH OXYCHLORIDE (UNII: 4ZR792I587)		
1,2-HEXANEDIOL (UNII: TR046Y3K1G)		

Product Characteristics			
Color	white (White to Pale Yellow)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66800-	142 g in 1 CAN; Type 0: Not a Combination	11/02/2020	

0003-5	Product	1/02/2020	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	11/02/2020	

Labeler - Beiersdord Inc (001177906)

Revised: 12/2025 Beiersdord Inc