

NUMQUICK ANALGESIC- epinephrine hydrochloride, lidocaine hydrochloride spray
Unit Dose, Ltd.

Numquick Analgesic

Drug Facts

Active ingredient

Epinephrine HCl 0.1 mg

Lidocaine HCl 50 mg

Purpose

Vasoconstrictor

Anesthetic

Uses:

Temporarily relieves local discomfort or pain and swelling or burning associated with anorectal disorders.

Warnings

External use only

certain persons can develop allergic reactions to the ingredients in this product **Allergy alert:**

Allergy alert:

certain persons can develop allergic reactions to the ingredients in this product

Do Not Use

this product in the rectum by using fingers or any mechanical device or applicator

When Using This Product

do not use more than directed.

Stop use and ask a doctor

if pain worsens or does not improve in 7 days

- redness, irritation, swelling, pain or other symptoms develop or increase
- if bleeding occurs

Keep out of the reach of children.

If accidentally swallowed get medical help or contact a Poison Control Center right away.

Directions:

Adults and Children over age 12 Spray 1-2ml to clean, dry affected area as needed up to 6 times daily. ---

Children under age 12: ask a doctor

Store in a cool, dark place or refrigerate.

Inactive Ingredients:

Aqua (DI Water), Chlorobutanol, Glycerin, Sodium chloride, Sodium metabisulfite


Questions?

Call Toll Free 888/664-9990

Package Labeling:



Drug Facts

Active ingredient (in each ml).....Purpose
 Epinephrine HCl 0.1 mgVasoconstrictor
 Lidocaine HCl 50 mgAnesthetic
Uses: Temporarily relieves local discomfort or pain and swelling or burning associated with anorectal disorders.
Warnings External Use Only
 Allergy alert: certain persons can develop allergic reactions to the ingredients in this product
 Do Not Use this product in the rectum by using fingers or any mechanical device or applicator
When Using This Product do not use more than directed.
 Stop use and ask a doctor if pain worsens or does not improve in 7 days • redness, irritation, swelling, pain or other symptoms develop or increase • if bleeding occurs
 Keep out of the reach of children. If accidentally swallowed get medical help or contact a Poison Control Center right away.
Directions: Adults and Children over age 12--- Spray 1-2 ml to clean, dry affected area as needed up to 6 times daily.
 Children under age 12: ask a doctor
 Store in a cool, dark place or refrigerate.
 Inactive Ingredients: Aqua (DI Water), Chlorobutanol, Glycerin, Sodium chloride, Sodium metabisulfite
 Questions? Call Toll Free 888/664-9990  6/02

NUMQUICK ANALGESIC

epinephrine hydrochloride, lidocaine hydrochloride spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67194-008
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

EPINEPHRINE HYDROCHLORIDE (UNII: WBB0470038) (EPINEPHRINE - UNII:YKH834O4BH)	EPINEPHRINE HYDROCHLORIDE	0.1 mg in 1 mL
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	50 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
CHLOROBUTANOL (UNII: HM4YQM8WRC)	
GLYCERIN (UNII: PDC6A3C0OX)	
EDETATE SODIUM (UNII: MP1J8420LU)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67194-008-01	59 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/20/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M015	01/20/2016	

Labeler - Unit Dose, Ltd. (119080393)

Revised: 2/2024

Unit Dose, Ltd.