

IBUPROFEN CHILDRENS- ibuprofen suspension

Walgreens

Ibuprofen Oral Suspension, USP

Active ingredient (in each 5 mL)

Ibuprofen, USP 100 mg (NSAID)**

**nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless

directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL =milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- **each 5 mL contains:** sodium 2 mg
- store between 20° to 25°C (68° to 77°F)

Inactive ingredients

acesulfame potassium, citric acid anhydrous, glycerin, polysorbate 80, pregelatinized corn starch, purified water, sodium benzoate, strawberry flavor, sucrose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday – Friday 9AM-5PM EST

Principal display panel

Compare to Children's Motrin® Dye-Free Berry active ingredient††

FOR AGES 2 TO 11 YEARS

Children's

Dye-Free

Ibuprofen 100

Ibuprofen Oral Suspension, USP **(NSAID)**

100 mg per 5 mL

Pain Reliever/Fever Reducer

Lasts up to 8 hours

Alcohol Free

Non-staining

Berry Flavor

Shake Well Before Using

FL OZ (mL)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

††This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin®.

Walgreens Pharmacist Recommended Walgreens Pharmacist Survey

DISTRIBUTED BY : WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015

Package label

Drug Facts (continued)

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Drug Facts (continued)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20° to 25°C (68° to 77°F)

Inactive ingredients acesulfame potassium, citric acid anhydrous, glycerin, polysorbate 80, pregelatinized corn starch, purified water, sodium benzoate, strawberry flavor, sucrose, xanthan gum

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

DISTRIBUTED BY: WALGREEN CO. 200 WILMOT RD. DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2019 Walgreen Co.

ITEM 982192 W10141-1119-O



3 11917 19116 4

17610317 GW710263 R0717 PLD-A630A FC006553

Walgreens

Compare to Children's Motrin®
Dye-Free Berry active ingredient!

NDC 0363-9370-04

FOR AGES 2 TO 11 YEARS
children's
Dye-Free
Ibuprofen 100

IBUPROFEN
ORAL SUSPENSION, USP
(NSAID)
100 mg PER 5 mL
PAIN RELIEVER /
FEVER REDUCER



DYE FREE
LASTS UP TO 8 HOURS
ALCOHOL FREE

• Non-staining

AGES
2-11
YEARS



BERRY FLAVOR
Shake well before using

4 FL OZ (118 mL)

IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING. KEEP THIS BOX FOR IMPORTANT INFORMATION. THIS PRODUCT IS INTENDED FOR USE IN CHILDREN.

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Ibuprofen, USP 100 mg (NSAID)**	Pain reliever/fever reducer
**nonsteroidal anti-inflammatory drug	

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

hives	facial swelling
asthma (wheezing)	shock
skin redness	rash
blisters	

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

TAMPER EVIDENT: DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

Walgreen Children's Ibuprofen 100mg/5mL Berry flavor

IBUPROFEN CHILDRENS

ibuprofen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9370
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
STARCH, CORN (UNII: O8232NY3SJ)	
WATER (UNII: 059QF0KOOR)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	

SUCROSE (UNII: C151H8M554)
XANTHAN GUM (UNII: TTV12P4NEE)

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9370-04	1 in 1 CARTON	10/24/2019	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074916	10/24/2019	

Labeler - Walgreens (008965063)

Revised: 11/2022

Walgreens