

COLD AND FLU- acetaminophen, dextromethorphan hbr, doxylamine succinate solution
Walgreen Company

Walgreen Co. Cold & Flu Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children. Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

| | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 6 hrs |
| children 4 to under 12 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- **each 30 mL contains:** sodium 39 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions

1-800-719-9260

Package/Label Principal Display Panel

Walgreens

WALGREENS PHARMACIST RECOMMENDED

Compare to the active ingredients in Vicks® NyQuil®

NIGHTTIME

Cold & Flu

ACETAMINOPHEN / ACHES / FEVER / SORE THROAT

DEXTROMETHORPHAN HBr / COUGH

DOXYLAMINE SUCCINATE / SNEEZING / RUNNY NOSE

Multi-Symptom

- Pain reliever, fever reducer, cough suppressant, & antihistamine
- ALCOHOL 10%

Cherry Flavor

12 FL OZ (355 mL)

NDC 0363-0568-40

Walgreens



Compare to the active ingredients in Vicks[®] NyQuil[®]††

NIGHTTIME

Cold & Flu

ACETAMINOPHEN /
ACHES / FEVER / SORE THROAT
DEXTROMETHORPHAN HBr / COUGH
DOXYLAMINE SUCCINATE /
SNEEZING / RUNNY NOSE

Multi-Symptom

- Pain reliever, fever reducer, cough suppressant & antihistamine
- ALCOHOL 10%



Cherry flavor

12 FL OZ (355 mL)

: 45940 94 F9

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.

ITEM 163057 W10107-0522-F

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

Gluten Free
W3ORG0222-F

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



Drug Facts

| Active ingredients (in each 30 mL) | Purpose |
|------------------------------------|-----------------------------|
| Acetaminophen 650 mg | Pain reliever/fever reducer |
| Dextromethorphan HBr 30 mg | Cough suppressant |
| Doxylamine succinate 12.5 mg | Antihistamine |

Uses temporarily relieves common cold/flu symptoms:
 cough due to minor throat and bronchial irritation
 sore throat headache minor aches and pains
 fever runny nose and sneezing

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening blisters rash
 If a skin reaction occurs, stop use and seek medical help right away.

PEEL BACK AT CORNER FOR MORE INFORMATION

: 45940 94 B1

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 if you have ever had an allergic reaction to this product or any of its ingredients

ADHESIVE AREA
NO VARNISH • NO TYPE

Drug Facts (continued)


When using this product excitability may occur, especially in children marked drowsiness may occur avoid alcoholic drinks
 be careful when driving a motor vehicle or operating machinery alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if pain or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions take only as directed - see Overdose warning only use the dose cup provided do not exceed 4 doses per 24 hrs

ADHESIVE AREA
NO VARNISH • NO TYPE



COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:0363-0566 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|---------------------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 650 mg in 30 mL |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 30 mg in 30 mL |
| DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL) | DOXYLAMINE SUCCINATE | 12.5 mg in 30 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ALCOHOL (UNII: 3K9958V90M) | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) | |

Product Characteristics

| | | | |
|-----------------|------------------------|---------------------|--|
| Color | RED (Clear/Dark Red) | Score | |
| Shape | | Size | |
| Flavor | CHERRY (Menthol Aroma) | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:0363-0566-50 | 2 in 1 PACKAGE | 08/16/2011 | |
| 1 | | 355 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 2 | NDC:0363-0566-34 | 237 mL in 1 BOTTLE; Type 0: Not a Combination Product | 08/10/2012 | 09/30/2025 |
| 3 | NDC:0363-0566-43 | 473 mL in 1 BOTTLE; Type 0: Not a Combination Product | 09/09/2011 | 04/16/2016 |
| 4 | NDC:0363-0566-40 | 355 mL in 1 BOTTLE; Type 0: Not a Combination Product | 08/16/2011 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 08/16/2011 | |

Labeler - Walgreen Company (008965063)

Revised: 10/2024

Walgreen Company