

**IBUPROFEN- ibuprofen capsule, liquid filled
Preferred Plus (Kinray)**

DRUG FACTS

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*

(present as the free acid and potassium salt)

***nonsteroidal anti-inflammatory drug**

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
 - take 1 capsule every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 capsule, 2 capsules may be used
 - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

Inactive ingredients

FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

†Compare to the active ingredient in Advil® Liqui-Gels®

Ibuprofen Capsules, 200 mg

Pain Reliever/Fever Reducer (**NSAID**)

Softgels**

(**Liquid filled capsules)

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil® Liqui-Gels®.

Distributed by: Kinray, Inc.

152-35 Tenth Avenue

Whitestone, NY 11357

©2013 Kinray Inc., All Rights Reserved, PREFERRED PLUS, PREFERRED PLUS++++ and the PREFERRED PLUS logo are trademarks and/or registered trademarks of Kinray Inc. All other marks are property of their respective owners.

Product Label

NO INK
NO VARNISH



Ibuprofen

Capsules, 200 mg

Pain Reliever/Fever Reducer (NSAID)

80 Softgels**
(*Liqud filled capsules)



3222Q7A

NDC 61715-054-38

*Compare to the active ingredient in Advil® Liqui-Gels®

NO INK
NO VARNISH

Ibuprofen

Capsules, 200 mg

Pain Reliever/Fever Reducer (NSAID)



80 Softgels**
(*Liqud filled capsules)

PLD-D F0001133
Rev# 03/12
Lot No.:
Exp. Date:



3 597226 27180 6

KIN# 713-834

Distributed by: Kinay, Inc.
152-35 10th Avenue
Whitestone, NY 11357

All Preferred Plus Pharmacy products are satisfaction guaranteed or your money back from retailer where purchased.

Drug Facts

Active ingredient (in each capsule) Purpose
Sulfonamide free ibuprofen is equal to 200 mg ibuprofen (NSAID). Pain reliever/fever reducer (NSAID). Present as the free acid and potassium salt nonsteroidal anti-inflammatory drug.

Uses
temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- joint pain of arthritis
- backache
- menstrual cramps
- temporarily reduces fever

Warnings
Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- swollen face
- skin redness
- facial swelling
- asthma (wheezing)
- blistering
- itching

 If these symptoms occur, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- take aspirin, other NSAIDs, blood thinners, or blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, etc.)
- use alcohol while taking this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Drug Facts (continued)

Ask a doctor before if

- the stomach is upset
- you are taking other pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver disease, or asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- do not take more than directed or for longer than directed
- you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - stools are black and sticky
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important during pregnancy because the use of NSAIDs in the 3rd trimester may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
- do not take more than 12 capsules in 24 hours, unless directed by a doctor
- take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not take more than 12 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- contains potassium 20 mg
- keep all warnings and directions before use.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

Inactive ingredients FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil® Liqui-Gels®.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IS IMPRINTED OR MISSING.

©2013 Kinay, Inc. All Rights Reserved. PREFERRED PLUS, PREFERRED PLUS+ and the PREFERRED PLUS logo are trademarks and/or registered trademarks of Kinay Inc. All other marks are property of their respective owners.

Ibuprofen Softgels

IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:61715-054

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0K00R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	BLUE	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	IB200
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61715-054-38	1 in 1 BOX	08/30/2010	12/31/2020
1		80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078682	08/30/2010	12/31/2020

Labeler - Preferred Plus (Kinray) (012574513)