

**MEDI-FIRST TOPICAL ANALGESIC- lidocaine hci spray**  
**Ultra Distributors Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**MEDI-FIRST®**  
**Burn spray**

***Active ingredient***

Lidocaine HCl 2.0%

***Purpose***

Topical pain relief

***Uses***

Temporary pain relief associated with minor burns

***Warnings***

**For external use only.**

**Do not use**

- in large quantities, particularly over raw or blistered areas
- near eyes, if this happens rinse thoroughly with water

**Stop use and ask a doctor**

if condition worsens or persists for more than 7 days or clears up and returns.

**Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away.

***Directions***

- spray an even layer of burn spray over cleaned affected area not more than 3-4 times daily
- not to be used on Children under 12 years of age

***Inactive ingredients***

glycerin, hydroxypropyl methylcellulose, melaleuca alternifolia (tea tree) leaf oil, octoxynol 9, PEG-40 hydrogenated castor oil, phenoxyethanol, propylene glycol, triethanolamine, water

***Questions or comments?***

1-800-634-7680

Topical Analgesic

Relieves pain in minor burns  
 Store at 68°-77°F (20°-25°C)

# MEDI-FIRST® Burn Spray

**Pump Spray**  
**Lidocaine HCl 2.0%**  
**Topical Analgesic**

Relieves pain in  
 minor burns

Store at 68°-77°F (20°-25°C)

**2 FL OZ (59.1 ML)**

**Drug Facts**

Active Ingredient	Purpose
Lidocaine HCl 2.0%	Topical pain relief

**Uses** Temporary pain relief associated with minor burns

**Warnings**  
 For external use only.  
 Do not use in large quantities, particularly over raw or blistered areas near eyes, if this happens rinse thoroughly with water  
 Stop use and ask a doctor if condition worsens or persists for more than 7 days or clears up and returns.  
 Keep out of reach of children. If swallowed get medical help or contact a Poison Control Center right away.

**Directions** ■ spray an even layer of burn spray over cleaned affected area not more than 3-4 times daily ■ not to be used on children under 12 years of age

**Inactive ingredients** glycerin, hydroxypropyl methylcellulose, melaleuca alternifolia (tea tree) leaf oil, octoxynol 9, PEG-40 hydrogenated castor oil, phenoxyethanol, propylene glycol, triethanolamine, water

**Questions or comments?** 1-800-634-7680



3 47682 22502 5

Rev. 03/20  
 Reorder #22502  
 Mfg. for Medique Products  
 Fort Myers, FL 33967 USA  
 www.mediqueproducts.com  
 MADE IN CHINA

## MEDI-FIRST TOPICAL ANALGESIC

lidocaine hci spray

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:78495-142
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	20 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
GLYCERIN (UNII: PDC6A3C0OX)	
OCTOXYNOL-9 (UNII: 7JPC6Y25QS)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

<b>TROLAMINE</b> (UNII: 9O3K93S3TK)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>TEA TREE OIL</b> (UNII: VIF565UC2G)	
<b>POLYOXYL 40 HYDROGENATED CASTOR OIL</b> (UNII: 7YC686GQ8F)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:78495-142-01	59.1 mL in 1 BOTTLE, PUMP; Type 0: Not a Combination Product	12/19/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	12/19/2020	

**Labeler** - Ultra Distributors Inc (007160073)

Revised: 12/2020

Ultra Distributors Inc