

**PENTREXCILINA DAYTIME- acetaminophen, chlorpheniramine maleate,
phenylephrine hcl tablet
OPMX LLC**

Pentrexilina - Daytime

ACTIVE INGREDIENTS (in each tablet)

Acetaminophen 500 mg

Chlorpheniramine Maleate 4 mg

Phenylephrine HCl 10 mg

PURPOSE

Pain Reliever/Fever Reducer

Antihistamine

Nasal decongestant

Uses:

- temporarily relieves these symptoms of hay fever or other respiratory allergies:
- headache
- sinus congestion & pressure
- nasal congestion
- runny nose and sneezing
- minor aches & pain
- temporarily relieves these additional symptoms of hay fever:
- nose or throat itching
- itchy, watery eyes
- helps clear nasal passages
- helps decongest sinus opening and passages

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take :

- more than 6 tablets in 24 hours, which is the maximum daily amount for this product
- with other drugs contains acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult doctor

promptly

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- to make a child sleepy

Ask a doctor before use if you have:

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- glaucoma
- diabetes
- trouble urinating due to enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are:

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product:

- do not use more than directed
- excitability may occur, especially in children
- drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if:

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or last more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding: ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not

notice any signs or symptoms.

Directions:

- Do not take more than directed - see overdose warning
- Swallow whole; do not crush, chew or dissolve
- Do not exceed 6 tablets per 24 hours

Adults and children 12 years of age & over	1 tablet with water every 4-6 hours as needed
Children 4 to 12 years of age	Ask a doctor
Children under 4 years of age	Do not use

Other information:

- Store at room temperature between 20-25°C (68-77°F): excursions permitted between 15°C-30°C (59°F-86°F)
- Avoid excessive heat and humidity
- Tamper Evident: do not use if package or blister unit is open or shows any sing of tampering

Inactive ingredients: Microcrystalline cellulose, Polyvinylpyrrolidone, Crospovidone, Magnesium stearate, Colloidal silicon dioxide, Purified talc

Questions or comments?

Call: (619) 600-5632 (Mon-Fri 9am-5pm EST) or <https://www.opmx.us>

TAMPER EVIDENT: DO NOT USE IF POUCH IS TORN, BROKEN OR SHOWS ANY SIGN OF TAMPERING

Exclusively distributed by:

OPMX

San Diego, CA 92154

Phone: 619-600-5632

PRINCIPAL DISPLAY PANEL - 2 TABLETS

NDC 69729-223-02

PENTREXCILINA

Actaminophen 500 mg.....Pain reliever/Fever reducer

Chlorpheniramine Maleate 4 mg.....Antihistamine

Phenylephrine HCl 10 mg.....Nasal decongestant


Effective and Quick Relief from cold & flu symptoms

Headache

Nasal Congestion

Itchy, Watery Eyes

Runny Nose & Sneezing



PRINCIPAL DISPLAY PANEL - 144 TABLETS

COLD & FLU

NDC 69729-223-38

DAYTIME

Actaminophen 500 mg • Chlorpheniramine Maleate 4 mg • Phenylephrine HCl 10 mg

PENTREXCILINA

Effective and Quick Relief from cold & flu symptoms

Headache

Nasal Congestion

Itchy, Watery Eyes

Runny Nose & Sneezing

144 Tablets (72 Pouches of 2 tablets)



PRINCIPAL DISPLAY PANEL - 12 TABLETS

COLD & FLU

NDC 69729-223-37

DAYTIME

Actaminophen 500 mg • Chlorpheniramine Maleate 4 mg • Phenylephrine HCl 10 mg

PENTREXCILINA

Effective and Quick Relief from cold & flu symptoms


Headache

Nasal Congestion

Itchy, Watery Eyes

Runny Nose & Sneezing

12 Tablets (6 Pouches of 2 tablets)



PRINCIPAL DISPLAY PANEL - 6 TABLETS

COLD & FLU

NDC 69729-223-06

DAYTIME

Actaminophen 500 mg • Chlorpheniramine Maleate 4 mg • Phenylephrine HCl 10 mg

PENTREXCILINA

Effective and Quick Relief from cold & flu symptoms

Headache

Nasal Congestion

Itchy, Watery Eyes
Runny Nose & Sneezing

6 Tablets (3 Pouches of 2 tablets)

**Eficaz y rápido
alivio de los
síntomas de
gripe y resfriado**

- Dolor de Cabeza
- Congestión Nasal
- Ojos llorosos
- Escurrimiento nasal y estornudos

INSTRUCCIONES EN ESPAÑOL EN EL INTERIOR DE LA CAJA.

PENTREXCILINA

 **DAYTIME**

COLD & FLU

NDC 69729-223-06

 **DAYTIME**

Acetaminophen 500 mg - Chlorpheniramine Maleate 4 mg - Phenylephrine HCl 10 mg

PENTREXCILINA

**Effective and
quick relief
from Cold & Flu
Symptoms**

- Headache
- Nasal Congestion
- Itchy, Watery Eyes
- Runny Nose & Sneezing

6 Tablets

**3 Packs
of 2 Tablets**



READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT INFORMATION

Exclusively distributed by:


Chula Vista, CA 91910
Phone: 619-600-5632



Made for:


San Diego, CA

PENTREXCILINA  **DAYTIME**

2A07DB01-00



REV. 01/2024 ITEM # RRX131
Code
LOT
EXP. 1

Directions
Do not take more than directed
- See Overdose warning
Swallow whole; do not crush, chew or dissolve
Do not exceed 6 tablets per 24 hours

Other information
Store at room temperature 25°C (77°F); excursions permitted between 15°C - 30°C (59°F - 86°F)
Avoid excessive heat and humidity
Tamper Evident: do not use if package or blister unit is open or shows any sign of tampering

Inactive ingredients
Colloidal silicon dioxide, Purified talc

Questions or comments? Call: (619) 600-5632 (Mon-Fri 9am-5pm EST) or <https://www.opmx.us>

Adults and children	1 tablet with water every 4-6 hours as needed
Children 4 to 12 years of age	Ask a doctor
Children under 4 years of age	Do not use

Keep out of reach of children
Overdose warning: taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If pregnant or breast-feeding: ask a health professional before use.

new symptoms occur. These could be signs of a serious condition.

Use or reuse more than 7 days - see your doctor or health professional if you have any of the following symptoms:

Drug Facts

Active Ingredients (in each tablet)
 Acetaminophen 500 mg
 Chlorpheniramine Maleate 4 mg
 Phenylephrine HCl 10 mg
 Nasal decongestant

Uses ■ temporarily relieves these symptoms of hay fever or other upper respiratory allergies ■ headache ■ sinus congestion & pressure ■ nasal congestion ■ runny nose and sneezing ■ minor aches and pains ■ temporarily relieves these additional symptoms of hay fever: ■ itching of the nose or throat ■ itchy, watery eyes ■ helps clear nasal passages ■ helps decongest sinus openings and passages

Warnings **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 6 tablets in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ Skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription or over-the-counter (OTC) drug for depression, psychiatric or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients ■ to make a child sleepy

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ glaucoma ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ drowsiness may occur ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain or nasal congestion gets worse or lasts more than 7 days ■ fever lasts more than 3 days ■ redness or swelling is present

No. : GUL/DRUGS/G/25/2644

DATE

PENTREXCILINA DAYTIME

acetaminophen, chlorpheniramine maleate, phenylephrine hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69729-223
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1W5297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white	Score	no score
--------------	-------	--------------	----------

Shape	ROUND	Size	13mm
Flavor		Imprint Code	A15
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69729-223-06	3 in 1 CARTON	01/10/2024	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:69729-223-38	72 in 1 CARTON	12/30/2020	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:69729-223-37	6 in 1 CARTON	01/10/2024	
3		2 in 1 POUCH; Type 0: Not a Combination Product		
4	NDC:69729-223-02	2 in 1 POUCH; Type 0: Not a Combination Product	12/30/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/30/2020	

Labeler - OPMX LLC (029918743)

Revised: 1/2024

OPMX LLC