

## **4X MEDICATED GUM CREAM- benzocaine cream Albertsons**

-----  
**5820621 Albertsons 4x Cream**

Benzalkonium chloride 0.13%

Benzocaine 20%

Menthol 0.26%

Zinc chloride 0.15%

Oral antiseptic

Oral pain reliever

Oral pain reliever

Oral astringent

- for the temporary relief of pain due to minor irritation or injury of the mouth and gums
- first aid to help prevent infection in minor oral irritation
  
- cut open tip of tube on score mark
- **adults and children 2 years of age and older:** apply to affected area up to 4 times daily or as directed by a dentist or doctor
- **children between 2 and 12 years of age:** ask a doctor before use. Should be supervised in the use of this product
- **children under 2 years of age:** do not use

**Methemoglobinemia warning:** use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy alert:** do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics

**Do not use** \* more than directed \* for more than 7 days unless told to do so by a dentist or doctor \* for teething \* in children under 2 years of age

**Stop use and ask a doctor if** \* swelling, rash or fever develops \* irritation, pain or redness persists or worsens \* symptoms do not improve in 7 days \* allergic reaction occurs

Keep out of reach of children. In case of overdose or allergic reaction, get medical help

or contact a Poison Control Center right away.

- do not use if tip is cut prior to opening
- avoid using toothpaste or drinking soft drinks or fruit juices for at least one hour after applying

ammonium glycyrrhizate, flavor, PEG-8, PEG-75, sodium saccharin, sorbic acid, titanium dioxide



Quality Guaranteed

Compare to the active ingredients in  
**Orajel™ 4X Medicated  
For Toothache & Gum Cream\***

NDC 21130-621-19

**CREAM**

# 4X MEDICATED TOOTHACHE & GUM PAIN RELIEF

**IMMEDIATE RELIEF OF SEVERE PAIN**

- 20% Benzocaine to relieve oral pain
- Kills harmful bacteria
- Astringent to help prevent infection
- Cools gum & relieves irritation

Oral antiseptic / Pain Reliever / Astringent

- 20% Benzocaine to relieve oral pain
- Kills harmful bacteria
- Astringent to help prevent infection
- Cools gum & relieves irritation

Oral antiseptic / Pain Reliever / Astringent

NET WT 0.33 OZ (9.4 g)

**TAMPER - EVIDENT: Do not use if tube tip is cut prior to use  
Retain carton for complete product information**

\*Orajel is a registered trademark of Church & Dwight Co.,  
Inc. and is not affiliated with this product.

DISTRIBUTED BY  
**BETTER LIVING BRANDS LLC**  
P.O. BOX 99, PLEASANTON, CA 94566-0009  
1-888-723-3929  
www.betterlivingbrandsLLC.com



Scan here  
for more  
information



RD 21082 S5538

2821621R2  
5820621

<b>Drug Facts</b>	
<b>Active ingredients</b>	Benzalkonium chloride 0.13% Benzocaine 20% Menthol 0.26% Zinc chloride 0.15%
<b>Purpose</b>	Oral antiseptic Oral pain reliever Oral pain reliever Oral astringent
<b>Uses</b>	• for the temporary relief of pain due to minor irritation or injury of the mouth and gums • first aid to help prevent infection in minor oral irritation
<b>Warnings</b>	<b>Methemoglobinemia warning:</b> use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: • pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy <b>Allergy alert:</b> do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics
<b>Do not use and ask a doctor if</b>	• more than directed • for more than 7 days unless told to do so by a dentist or doctor • for teething in children under 2 years of age • symptoms do not improve in 7 days • swelling, rash or fever develops • irritation, pain or redness persists or worsens
<b>Keep out of reach of children.</b>	In case of overdose or allergic reaction, get medical help or contact a Poison Control Center right away.
<b>Directions</b>	• cut open tip of tube on score mark • adults and children 2 years of age and older: apply to affected area up to 4 times daily or as directed by a dentist or doctor • children between 2 and 12 years of age: ask a doctor before use. Should be supervised in the use of this product • children under 2 years of age: do not use
<b>Other information</b>	• do not use if tip is cut prior to opening • avoid using toothpaste or drinking soft drinks or fruit juices for at least one hour after applying
<b>Inactive ingredients</b>	ammonium glycyrrhizate, flavor, PEG-8, PEG-75, PEG-8, PEG-75, sodium saccharin, sorbic acid, titanium dioxide

## 4X MEDICATED GUM CREAM

benzocaine cream

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:21130-621
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 g
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A) (MENTHOL, UNSPECIFIED FORM - UNII:L7T10EIP3A)	MENTHOL, UNSPECIFIED FORM	0.25 g in 100 g
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	20 g in 100 g
<b>ZINC CHLORIDE</b> (UNII: 86Q357L16B) (ZINC CATION - UNII:13S1S8SF37)	ZINC CATION	0.15 g in 100 g

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>AMMONIUM GLYCYRRHIZATE</b> (UNII: 3VRD35U26C)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SORBIC ACID</b> (UNII: X045WJ989B)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	

## Product Characteristics

<b>Color</b>	white (Cloudy white to yellow)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	WMTERGREEN	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:21130-621-19	9.4 g in 1 TUBE; Type 0: Not a Combination Product	05/07/2021	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M017	05/07/2021	

**Registrant** - Lornamead Products (119559733)

**Establishment**

Name	Address	ID/FEI	Business Operations
Lornamead Products		119559733	manufacture(21130-621) , pack(21130-621)

Revised: 11/2025

Albertsons