BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE- liquid bandage liquid Topco Associates LLC

TopCare Liquid Bandage

Active ingredients

Benzethonium chloride 0.2% Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic

Topical analgesic

Uses

• first aid to prevent bacteria contamination in minor cuts and scrapes

Warnings

For external use only.

Flammable:

do not use while smoking or near heat or flame.

Do not use

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor
- on infected areas or wounds that are draining
- with other first aid products such as lotions and creams
- over sutures
- on mucous membranes

Ask a doctor before use if you have

- deep puncture wounds
- deep cuts
- animal bites
- serious bleeding
- diabetes
- poor circulation
- serious burns

When using this product

do not sotre at temperaure above 120 °F

Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

Stop use and ask a doctor if

- conditions persists or gets worse
- infection occurs

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- Clean affected area
- apply a small amount on the area 1-3 times daily
- let dry
- a second coating may be applied for extra protection
- to remove, apply more Liquid Bandage and quickly wipe off
- finger nail polish remover may dissolve Liquid Bandage

Other information

- cap bottle tightly
- store at room temperature away from heat
- may temporarily sting upon application
- do not allow to come in contact with floors, countertops, or other finished surfaces will stain

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD slcohol 40

Questions

Call 1-866-964-0939

Principal Display Panel

TopCare

Liquid

Bandage

Topical Analgesic & Antiseptic to Help Prevent Infection

FOR SMALL CUTS & WOUNDS

• Dries Quickly

• Forms a Flexible, Invisible Protective Cover

USE LIQUID BANDAGE FOR:

- Blisters
- Chapped & Cracked Finger Tips
- Hangnails
- Paper Cuts
- Shaving Nicks
- Help in Preventing the Formulation of Calluses

GREAT FOR:

Athletes, Active Lifestyles, Outdoor Sportmen & Musicians

NET 0.3 FL OZ (9 mL)



ontrol Center right away.	
sti .nen bild of children. It s	, get medical help or call a Poison
to use and ask a doctor if a continue of task of any standard and the continue of task of the continue of t	
hen using this product avoid	jith eyes
avimal bites	or puncture wounds 🔳 deep cuts
orger than 1 meek unless ∎ over the sevent in a meek unless directed	
/arnings or external use only. Flammable:	.email to etit mort ya
to teiler yusroqmet ertt tot =	anud bna seqans, stus rorim ri noitar
%70.0 ebilothorbord eninology sets at the transmission of the transmission of the transmission of t	iseglana labiqoT
%2.0 ebrolrho muinorthezne %7.0 ebrolrhoorbyrt eninolw eiretsed meveng of bis fant = 252 to feiler ynsrogmer ent rot =	idqeatme bis tanii. Jiseglene leoiqoT
cTive ingredication scalably of the state 0.2% section by drochondo 0.75% since and of the same state and of the same of the same section of the state state of the same section of the same section of the section of the same section of the same section of the section of the same section of the same section of the section of the same section of the same section of the section of the same section of the same section of the section of the same section of the same section of the section of the same section of the same section of the section of the same section of t	iseglana labiqoT
%2.0 ebrolrho muinorthezne %7.0 ebrolrhoorbyrt eninolw eiretsed meveng of bis fant = 252 to feiler ynsrogmer ent rot =	idqeatme bis tanii. Jiseglene leoiqoT
Prug Facts crive ingredients crive ingredients crive indroching of 2% crives cristad neveration cristad neveration cristad	ອຂວຕາມໆ ອັດອອນເຫຼັດເອັດອອນເອີ້ອງ ອີເອອອີເລາຍ ໂອວິເຊວາ
cTive ingredication scalabolic muinofiscre scalabolic provent of the static section of the static section of the section of the section of the section of the section of the section of the section of the section of the section of the section of the section of the section of the section of the section of the section of the section of th	idqeatme bis tanii. Jiseglene leoiqoT
Prug Facts crive ingredients crive ingredients crive indroductor of 2% crives criments cri	ອຂວຕາມໆ ອັດອອນເຫຼັດເອັດອອນເອີ້ອງ ອີເອອອີເລາຍ ໂອວິເຊວາ

BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE

liquid bandage liquid

_							
Ρ	roduct Info	ormation					
P	roduct Type		HUMAN OTC DRUG	Item Code	(Source)	NDC:	36800-382
R	oute of Admi	nistration	TOPICAL				
A	ctive Ingre	dient/Active	Moiety				
		Ingred	lient Name		Basis of S	trength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - BENZ					BENZ ETHONIL CHLORIDE	JM	0.2 mg in 9 mL
	YCLONINE HYE		INII: ZEC193879Q) (DYCLON	NINE -	DYCLONINE HYDROCHLORI	IDE	0.75 mg in 9 mL
In	nactive Ing	redients					
		I	ngredient Name			St	trength
AC	CETONE (UNII:	1364PS73AF)					
AN	MYL ACETATE	(UNII: 92Q24NH7A	S)				
CASTOR OIL (UNII: D5340Y2I9G)							
ET	HYL ACETATE	(UNII: 7684508N	MZ)				
PYROXYLIN (UNII: KYR8BR2X6O)							
AL	COHOL (UNII:	3K9958V90M)					
	ackaging						
P					Marketing S	Start M	arketing End
	Item Code	Р	ackage Description		Date		Date

Marketing Information								
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date					
OTC Monograph Drug	M003	07/10/2014						

Labeler - Topco Associates LLC (006935977)

Revised: 12/2024

Topco Associates LLC