

ALLEROFF - cetirizine hydrochloride syrup
Corporacion Infarmasa

Drug Facts

(in each 5 ml teaspoonful) Cetirizine HCL 5 mg.....Antihistamine

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product:

- Drowsiness may occur
- Avoid alcoholic drinks
- Alcohol, sedatives and tranquilizers may increase drowsiness
- Be careful when driving a motor vehicle or operating machinery

Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

If pregnant or breast feeding:

- if breast feeding: not recommended
- if pregnant: ask a health professional before use

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

Uses

Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- Runny Nose
- Itchy, water eyes
- Itching of the nose or throat
- Sneezing

If you have questions of a medical nature, please contact your pharmacist, doctor or health care professional

Questions or comments? 561 338 5221

Store between 20°C to 25°C (68°F to 77°F)

See bottom panel for lot number and expiration date

Use only with enclosed dosing cup

Adults and Children 6 years and over 1 teaspoonful (5ml) or 2 teaspoonful (10ml) once daily depending upon severity of symptoms; do not take more than 2 teaspoonful (10ml) in 24 hours

Adults 65 years and over 1 teaspoonful (5ml) once daily; do not take more than 1 teaspoonful (5ml) in 24 hours

Children 2 to under 6 years of age 1/2 teaspoonful (2.5ml) once daily. If needed, dose can be increased to a maximum of 1 teaspoon (5ml) once daily or 1/2 teaspoonful (2.5ml) every 12 hours. Do not give more than 1 teaspoonful (5ml) in 24 hours

Children under 2 years of age Ask a Doctor

Consumer with liver or kidney disease Ask a Doctor

Citric and monohydrate, Sodium Benzoate, Sodium Citrate Tribasic Dihydrate, FDC Red N° 40, Disodium Edetate, Strawberry Essence Liquid, Glycerol, Saccharin Sodium, Sorbitol, Purified Water

Expiration Date:
Lot #:
oral solution
1 mg/mL
Antihistamine /
Cetirizine HCl
alleroff
allergy

Drug facts

Active Ingredient	Purpose
Cetirizine HCl (5 mg in each teaspoonful)	Antihistamine

Uses
Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- Sneezing
- Itchy, watery eyes
- Runny nose
- Itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
If you are taking tranquilizers or sedatives, ask a doctor or a health professional before using this product.
When using this product:
- Drowsiness may occur
- Avoid alcoholic drinks
- Alcohol, sedatives and tranquilizers may increase drowsiness.
- Be careful when driving a motor vehicle or operating machinery.
Stop using and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
- If you are breast-feeding, not recommended.
- If you are pregnant, ask a health professional before using.
Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).


Directions
Use only with dosing cup provided.
Adults and children 6 years and over: 1 teaspoonful (5 mL) or 2 teaspoonfuls (10 mL) once daily depending upon severity or symptoms; do not take more than 2 teaspoonfuls (10 mL) in 24 hours.
Adults 65 years and over: 1 teaspoonful (5 mL) once daily, do not take more than 1 teaspoonful (5 mL) in 24 hours.
Children 2 to under 6 years of age: ½ teaspoonful (2.5 mL) once daily. If needed, dose can be increased to a maximum of 1 teaspoonful (5 mL) once daily or ½ teaspoonful (2.5 mL) every 12 hours. Do not give more than 1 teaspoonful (5 mL) in 24 hours.
Children under 2 years of age: Ask a doctor
Consumers with liver or kidney disease: Ask a doctor

Other information
- Store between 20° to 25° C (68° to 77° F)
- See the top panel for the batch number and the expiry date

Inactive ingredients
Citric acid monohydrate, sodium benzoate, sodium citrate tribasic dihydrate, FD C Red N° 40, disodium edetate, strawberry essence liquid, glycerol, saccharin sodium, sorbitol, purified water.

Questions or comments? ☎ 561 338 5221
If you have questions of a medical nature, please contact your pharmacist, doctor or health professional.
*This product is not manufactured or distributed by McNeil Consumer Healthcare, division of McNeil-PPC, Inc., distributor of Zyrtec®.

Manufactured by: **Corporación Infarmasa S.A.**, Lima, Peru
Distributor: **Global Corporation Inc.**, 621 N.W. 53rd St., Suite 240, Boca Raton, Florida 33487



alleroff
allergy

**Cetirizine HCl/
Antihistamine /
1 mg/mL
oral solution**

**24
horas** Relief of:

- Sneezing
- Runny nose
- Itchy, watery eyes
- Itchy nose or throat

2 yrs. & older



Strawberry Syrup

4 FL OZ (120 mL)

Datos de este medicamento

Ingrediente activo	Función
Cetirizina HCl (5 mg en cada 5 mL)	Antihistamínico

Usos
Alivia temporalmente los síntomas de la fiebre del heno u otras alergias respiratorias superiores como:
- Estornudo
- Escorzo, ojos llorosos
- Rinorrea
- Escorzo de la nariz o la garganta

Advertencias
No usar si alguna vez ha tenido reacción alérgica a este producto o a alguno de sus ingredientes o a un antihistamínico que contiene hidroxizina.
Si usted tiene una enfermedad del hígado o del riñón, consulte a su médico antes de usar este producto. Su médico debe determinar si necesita una dosis diferente.
Si usted está tomando tranquilizantes o sedantes, consulte a su médico o farmacéutico antes de usar este producto.
Al utilizar este producto:
- Puede ocurrir somnolencia.
- Evitar las bebidas alcohólicas.
- El alcohol, los sedantes y los tranquilizantes pueden aumentar la somnolencia.
- Tenga cuidado al conducir un vehículo motorizado o manejar maquinaria.
Suspenda el uso y consulte a su médico si se produce una reacción alérgica. Busque ayuda médica de inmediato.
- Si está dando de lactar: No se recomienda su uso.
- Si está embarazada: Consulte con su médico antes de usar este producto.
Mantener fuera del alcance de los niños. En caso de sobredosis, obtenga ayuda médica o póngase en contacto con un centro de toxicología de inmediato (1-800-222-1222).

Instrucciones Use el dosificador incluido.
Adultos y niños mayores de 6 años: 1 cucharadita (5 mL) ó 2 cucharaditas (10 mL) una vez al día dependiendo de la severidad de los síntomas, no tome más de 2 cucharaditas (10 mL) en 24 horas.
Adultos mayores de 65 años: 1 cucharadita (5 mL) una vez al día, no tome más de 1 cucharadita (5 mL) en 24 horas.
Niños de 2 a 6 años: ½ cucharadita (2.5 mL) una vez al día. Si es necesario, la dosis puede ser aumentada hasta un máximo de 1 cucharadita (5 mL) una vez al día o ½ cucharadita (2.5 mL) cada 12 horas. No le dé más de 1 cucharadita (5 mL) en 24 horas.
Menores de 2 años: Consulte a su médico.
Pacientes con enfermedad hepática o renal: Consulte a su médico.

Otra información
- Conservar entre 20° a 25° C (68° a 77° F)
- Revise el número de lote y la fecha de vencimiento.

Ingredientes inactivos
Ácido cítrico monohidratado, benato de sodio, citrato de sodio tribásico dihidratado, colorante rojo FD C N° 40, edetato disódico, esencia líquida de fresa, glicerol, sacarina sódica, sorbitol, agua purificada.

¿Preguntas o comentarios? ☎ 561 338 5221
O sinvas contactar con un profesional de la salud.

Fabricado por: **Corporación Infarmasa S.A.**, Lima, Peru
Distribuidor: **Global Corporation Inc.**, 621 N.W. 53rd St., Suite 240, Boca Raton, Florida 33487



alleroff
allergia

**Cetirizina HCl/
Antihistamínico/
1 mg/mL
solución oral**

**24
horas** Alivio de:

- Estornudo
- Rinorrea
- Escorzo, ojos llorosos
- Escorzo de la nariz o la garganta

2 años a más



Sabor a fresa

4 FL OZ (120 mL)

54 x 54 x 134

Enter section text here

ALLEROFF
cetirizine hydrochloride syrup

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:16853-126 1
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CETIRIZINE HYDROCHLORIDE (UNII: 64O047KTOA) (Cetirizine - UNII:YO7261ME24)	CETIRIZINE HYDROCHLORIDE	1 mg

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	0.080 mg
SODIUM BENZOATE (UNII: OJ245FE5EU)	0.100 mg
SODIUM CITRATE (UNII: 1Q73Q2JULR)	0.090 mg
FD&C RED NO. 40 (UNII: WZB9127XOA)	0.003 mg
EDETATE DISODIUM (UNII: 7FLD91C86K)	0.025 mg
GLYCERIN (UNII: PDC6A3C0OX)	25 mg
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	0.075 mg
SORBITOL (UNII: 506T60A25R)	30 mg
WATER (UNII: 059QF0KO0R)	100 mL
STRAWBERRY (UNII: 4J2TY8Y81V)	0.07 mg

Product Characteristics

Color	pink (pink)	Score	
Shape		Size	
Flavor	STRAWBERRY (Strawberry)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16853-126 1-1	1 in 1 CARTON		
2	NDC:16853-126 1-2	2 in 1 PACKAGE, COMBINATION		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA022155	01/15/20 10	

Labeler - Corporacion Infarmasa (934098294)

Establishment

Name	Address	ID/FEI	Business Operations
Corporacion Infarmasa		934098294	manufacture

