GERI-LANTA MAXIMUM STRENGTH- aluminum hydroxide, magnesium hydroxide, dimethicone liquid

Geri-Care Pharmaceuticals, Corp

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

gerilanta max

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 400mg (equivalent to dried gel, USP) Magnesium hydroxide 400 mg Simethicone 40mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if

symptoms last more than 2 weeks

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
- do not take more than 12 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks

• children under 12 years: ask a doctor

Other information

- each 5 mL teaspoonful contains: magnesium 165 mg, sodium 1 mg
- store at room temperature
- protect from freezing
- keep tightly closed
- TAMPER-EVIDENT: Do not use if breakaway band on bottle is missing or broken.

Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

package Label

NDC 57896-619-12

GERICARE

Maximum Strength GERI-LANTA

antacid/antigas

Alumina, Magnesia and Simethicone Oral Suspension For Relief of:

- HEARTBURN SOUR STOMACH
- ACID INDIGESTION
- PRESSURE AND BLOATING

compare to the active ingredients in Maximum Strength MYLANTA

12 FL OZ (355 mL)

Alcohol: 0.5%

GERICARE

Maximum Strength

GERI-LANTA

ANTACID • ANTIGAS

Alumina, Magnesia, and Simethicone Oral Suspension

For Relief of:

- HEARTBURN SOUR STOMACH
- ACID INDIGESTION
- PRESSURE AND BLOATING

Compare to the active ingredients in Maximum Strength MYLANTA®*

12 FL OZ (355 mL)

Alcohol: 0.5%

Drug Facts

Active ingredients (in each 5 mL teaspoonful)

Purposes

Uses relieves • heartburn • sour stomach
• acid indigestion • the symptoms referred to as gas

Warnings

Ask a doctor before use if you have · kidney disease · a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions • shake well before use • adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor • do not take more than 12 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks • children under 12 years: ask a doctor

Other information • each 5 mL teaspoonful contains:
magnesium 165 mg, sodium 1 mg • store at room temperature
• protect from freezing • keep tightly closed • TAMPER-EVIDENT:
Do not use if breakaway band on bottle cap is missing or broken.

Inactive ingredients benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

*This product is not manufactured or distributed by the owner of the registered trademark MYLANTA®.

Distributed by Geri-Care Pharmaceuticals 1650 63rd Street Brooklyn, NY 11204

REV 0111 2-910-06122-0



GERI-LANTA MAXIMUM STRENGTH

aluminum hydroxide, magnesium hydroxide, dimethicone liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:57896-619

210-06122-1 REV1113

Route of Administration ORAL

Active Ingredient/Active Moiety

ı	0	3		
		Ingredient Name	Basis of Strength	Strength
I			ALUMINUM	400 mg

ALUMINUM ITTOKO AIDE (UMI. 3QDU 12IUNU) (ALUMINUM ITTOKO AIDE - UMI.3QDU 12IUNU)	HYDRO XIDE	in 5 mL
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6 V3LHY838, HYDRO XIDE ION - UNII:9159 UV381P)	MAGNESIUM HYDRO XIDE	400 mg in 5 mL
DIMETHICO NE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 5 mL

Inactive Ingredients				
Ingredient Name	Strength			
BENZYL ALCOHOL (UNII: LKG8494WBH)				
BUTYLPARABEN (UNII: 3QPI1U3FV8)				
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679 OBS 311)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
WATER (UNII: 059QF0KO0R)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SORBITOL (UNII: 506T60A25R)				

Product Characteristics				
Color		Score		
Shape		Size		
Flavor	LEMON (Lemon)	Imprint Code		
Contains				

l	Packaging				
ı	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
ı	1	NDC:57896-619-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	0 1/0 1/20 11	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part331	0 1/0 1/20 0 0		

Labeler - Geri-Care Pharmaceuticals, Corp (611196254)

Registrant - GCP Laboratories (965480861)

Establishment			
Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(57896-619), pack(57896-619)